


Barlow Test



Ortolani Test

The Barlow test is an attempt to dislocate the hip. If positive, you will feel the hip sublux or dislocate. The Ortolani test is the maneuver to reduce a dislocated hip. If positive, you will feel a clunk.

**Orthopedics Consults & Referrals**

1. Consultation:
  - Beneficiary patients: contact ANMC orthopedic surgeon on call at (907) 563-2662 (\*97) or send message through Tiger Connect.
  - Non-beneficiary patients: contact Ken Thomas, MD at Anchorage Fracture & Orthopedics at (907) 563-3145.
2. Referral:
  - Place an order for "Refer to Orthopedics External" with brief history. Note the orthopedist who was consulted. Indicate where the referral should be sent.
  - Send a RAVEN Communication to Chronic Peds Case Manager Pool about the referral and level of importance.

**Imaging**  
Patient must have either ultrasound or X-ray, as below.

1. Hip ultrasound: 6 weeks to 4 months of age.
  - Performed at Alaska Regional Hospital.
  - Place order for "Refer to Pediatric Clinic External (MRI / EEG / VFSS / Hip US)" with brief history.
  - If patient is a beneficiary, request follow-up appointment at Southcentral Foundation Team B.
  - If patient is not a beneficiary, request follow-up appointment with a pediatric provider in Bethel.
  - Send a RAVEN Communication to Chronic Peds Case Manager Pool about the referral and level of importance.
2. X-ray, AP pelvis: over 4 months of age. (Note: in premature infants, ossification of femoral heads is delayed. May use corrected gestational age of 4 months or later.)
  - Performed at YKHC.
  - Place an order for "XR Pelvis (Pelvis AP only)" and put in comments "AP view with hips in neutral position to rule-out developmental dysplasia of the hip."
  - Send a RAVEN Communication to Chronic Peds Case Manager Pool stating the order was placed and requesting an appointment for this with a pediatric provider in Bethel.