



CLINICAL GUIDELINES

Arranged by system, and then alphabetical.

General

Consultations.....	3
Guideline Guideline.....	4
Process to Update the EHR to Match Guidelines.....	6

Critical Care & Emergency Medicine

Acetaminophen Overdose (Adult and Pediatric).....	7
Acute Coronary Syndrome (MI).....	8
Burns (Adult and Pediatric).....	12
Cerebrovascular Accident.....	13
Death Protocol.....	16
Frostbite (Adult and Pediatric).....	19
Head Injury (≥18 years).....	21
Head Injury/Concussion (<18 years).....	22
High-Flow Nasal Cannula (Pediatric).....	23
Hypothermia.....	24
Intubation (Adult and Pediatric).....	25
Intubated Pediatric Patients: ET CO ₂ Monitoring.....	29
Massive Transfusion Protocol (≥14 years).....	31
Medevac Activation: Village to YKHC.....	32
Medevac Activation: YKHC to Anchorage.....	33
Medevac/Transfer Process for Trauma.....	34
Military Transport for Emergencies.....	35
Pediatric Medevacs: Bethel to Anchorage.....	36
Procedural Sedation & Analgesia Outside the OR.....	37
Respiratory Distress & Bronchiolitis Management (<5 years).....	42
Seizure Evaluation (Pediatric).....	43
Sepsis (Adult).....	44
Sepsis Medications (Adult).....	45
Sepsis (Pediatric).....	47
Spinal Cord Injury Management.....	48
Status Epilepticus Treatment (Adult).....	50
Status Epilepticus Treatment (Pediatric).....	51
Strangulation.....	52
Trauma Outside Bethel.....	53
Villages without Health Aides.....	55

Adult Critical Care Guide:

https://yk-health.org/images/9/98/Adult_Critical_Care_Guide.pdf

Pediatric Critical Care Weight-Based Guide:

https://yk-health.org/images/5/5d/Pediatric_critical_care_guide.pdf

Abuse/Assault

Sexual Assault (≥18 Years).....	56
Suspected Physical Abuse Procedure (Pediatric).....	57
Suspected Sexual Abuse Procedure (Pediatric).....	58

Endocrine

Diabetes, Type 2.....	59
Pediatric Endocrine Protocols.....	61

Gastrointestinal

Congenital Sucrase-Isomaltase Deficiency Resource.....	62
Dyspepsia/H pylori (Adult and Pediatric).....	63

Hematologic

Anemia in Adults.....	64
Iron Infusion for Chronic Iron-Deficiency Anemia (Adult and Pediatrics).....	65

Infectious Disease

Amoxicillin Allergy Trials (Pediatric).....	66
Botulism.....	67
Bronchiectasis/Chronic Cough (<18 years).....	68
COVID: Molnupiravir, Emergency Use.....	69
COVID: Paxlovid.....	70
Croup/Stridor (6 months – 3 years).....	71
Fever (0-90 days).....	72
Fever in Underimmunized Children.....	73
Hepatitis C – click here for management recommendations.....	
Influenza (Adult and Pediatric).....	74
Lymphadenitis, Acute Cervical (Pediatric).....	75
Meningitis: Use of Dexamethasone.....	76
Mpox: Emergency Use of Tecovirimat.....	77
Multisystem Inflammatory Syndrome (MIS-C).....	78
Otitis Media, Acute (3 months – 12 years).....	79
Peritonsillar Abscess.....	80
Pertussis.....	81
Pharyngitis (Adults and Pediatric).....	82
Pneumonia (Adult).....	83
Pneumonia Treatment (3 months – 18 years).....	84
Rabies Prevention.....	85
Septic Arthritis and Osteomyelitis, Suspected (Pediatric).....	86
Sexually Transmitted Infections.....	87
Sinusitis (4 – 18 years).....	90
Skin and Soft Tissue Infection (Adult and Pediatric).....	91
Tuberculosis, Active Pulmonary (≥14 years).....	93
Tuberculosis, Latent (≥14 years).....	94
Tuberculosis Evaluation and Treatment (<14 years).....	95
Tuberculosis Testing: Induced Sputum Collection.....	96
UTI (Adult).....	97
UTI (3 months – 5 years).....	98
Varicella, Suspected.....	99



Neonatal/Pediatric Growth & Development

Admissions from Clinic (Pediatric).....	100
Caffeine Protocol, Post-NICU Discharge.....	101
Failure to Thrive in Children <24 Months.....	102
Hip Exam and Surveillance in Infants.....	104
Jaundice in a Baby <4 Weeks.....	105
Late Preterm & Low Birth Weight Infants, Care of.....	106
mPEWS Protocol for Pediatric Patients.....	107
Neonatal Nasal CPAP Set-Up Guide.....	108
Newborn Early-Onset Sepsis/GBS.....	110
Neonatal Glucose Screening.....	111
Neonatal Resuscitation Summary.....	112
Neopuff™ and ER Warmer Set-Up Guide.....	114
Surfactant Administration Protocol.....	118
Suspected Neonatal Withdrawal.....	119
Therapeutic Hypothermia for Neonatal Hypoxic-Ischemic Encephalopathy – PAMC NICU protocol	
Village Deliveries.....	120

Obstetrics

Anemia in Pregnancy.....	121
Aneuploidy.....	122
Diabetes, Gestational.....	123
Ectopic Pregnancy Treatment.....	124
First Trimester Vaginal Bleeding.....	125
Group B Streptococcus (Maternal).....	127
HIV Screening and Prenatal Care.....	128
Hypertension in Pregnancy, Chronic.....	129
Hypertension, Gestational/Preeclampsia.....	130
Hypertension in Pregnant and Postpartum Patients, Severe.....	131
Induction of Labor.....	133
Intrahepatic Cholestasis of Pregnancy.....	134
Intrauterine Growth Restriction.....	135
Labor Patient in a Village.....	136
Molar Pregnancy.....	137
Oligohydramnios.....	138
Post-Dates Pregnancy.....	139
Postpartum Hemorrhage.....	140
Prenatal Care.....	141
Preterm Labor.....	142
Preterm Premature Rupture of Membranes.....	144
Rhogam®.....	146
Trial of Labor after Caesarian Delivery.....	147

Preventative Health Care & Outpatient Protocols

Amoxicillin Allergy Trials (Pediatric).....	148
Breast Cancer Screening.....	149
Cervical Cancer Screening with hrHPV.....	150
Colorectal Cancer Screening.....	151
Hyperlipidemia.....	155
Hypertension.....	156
Lead Screening (Pediatric).....	157
Nirsevimab 2024-2025 Season.....	158
Osteoporosis Screening and Treatment.....	159
Pre-anesthesia Management.....	162
Primary Care for Ex-Premies – Checklist	164
Primary Prevention of Cardiovascular Disease.....	165
Sports Clearance for Pediatric Patients with History of COVID-19.....	166

Psychiatry

Alcohol Hangover/Withdrawal.....	167
Attention Deficit Hyperactivity Disorder (Pediatric).....	168
Care of an Agitated or Aggressive Patient on Inpatient or DES.....	169
Intoxicated Patient.....	170
Involuntary Psychiatric Admissions.....	171
Psychiatric Admissions (General).....	172

Documentation

Care Conference Checklist.....	173
Nutritional Supplements, Documentation Requirements.....	176



Yukon-Kuskokwim HEALTH CORPORATION

Treatment Protocol Consultations

Phone Numbers

ANMC: Consult: *97 or (907) 563-2662
 Transfer: (907) 729-2337
 PICU Cell for urgent consults: (907) 297-8809
 Providence: ED for on-call specialist: (907) 212-3111
 Trauma: (907) 212-2525
 Alaska Regional Hospital Access Center: (844) 880-5522
 VA/JBER: ED: MD consult number (907) 580-5556
 Transfer: (907) 580-6420
 Admissions 24/7: (907) 580-6423
 Operator: (907) 552-1110
 Harborview Seattle (burns): (888) 731-4791

Remember: Unless you transfer care of the patient, YOU are responsible for orders, documentation, and notifying the patient and family of the plan of care.

SRC and village itinerant providers do not have the luxury of paging the provider STAT to bedside. However, the SBAR case presentation and the documentation requirements listed on this protocol still apply.

Page the appropriate provider:

1. ANMC for beneficiaries.
2. Providence Hospital or Alaska Regional Hospital for non-beneficiaries.
3. Alaska Regional for prison inmate.
4. VA or JBER (Joint Base Elmendorf/Richardson) for veterans.

Provider needs consultation about patient at YKHC

Consulting provider located in Bethel?

No

Yes

Patient is critically ill and the consultant is required at bedside?

No

Yes

Page provider STAT to come to bedside and assist in management.

Be prepared with the following information:

1. State your name, title, and department (e.g. ED physician, outpatient NP, second year resident, etc.)
2. State purpose of call (e.g. quick question, possible admission, management advice, etc.)
3. Provide name, age, DOB, and location of patient. If the patient is pregnant, give gravity and parity and gestational age in initial sentence. If the patient is a child, give the age in the initial sentence.
4. Use SBAR (see box).
5. Ask a **specific question** about management.
6. If patient is to be transferred, state whether you think that the patient can travel by commercial flight or will require air medevac.
7. If there is a problem getting an accepting physician for a medevac/transfer or with patient management decisions, see NOTE below.

Document consultant advice in the medical record, include date, time, first and last name of consultant and a summary of the advice given.

If on-going management is required, a decision must be made immediately and communicated to the team about who will be the primary managing provider giving orders and documenting in the medical record.

Once patient is stabilized, discussion will occur between the primary provider and the consultant regarding further documentation and ongoing management.

Page the appropriate provider. Be prepared with the following information:

1. State your name, title, and department (e.g. ED physician, outpatient NP, second year resident, etc.)
2. State purpose of call, including if you want a formal consult (e.g. quick question, possible admission, management advice, etc.)
3. Name, MRN, age, DOB. If the patient is pregnant, give gravity and parity and gestational age in initial sentence. If the patient is a child, give the age in the initial sentence.
4. Use SBAR (see box).
5. Ask a **specific question** about management.

Provider requesting consult must document consultant's advice in the medical record. Include date, time, first and last name of consultant, and a summary of the advice given.

Note: consultants are encouraged to document their recommendations in a separate note or as an addendum to the provider note. If done, this note does not obviate the initial provider's documentation requirements.

At any time in the process, if the primary provider wants support at the bedside, page the consultant and ask them to come to bedside and provide support.

Clear role delineation must occur establishing who is the primary managing provider.

SBAR

Situation: a concise statement of the problem, a "one-liner"

"This is a 3 year old otherwise healthy girl with a fever..."
 "My patient is a 20 year old G3P2 at 26 weeks with vaginal bleeding..."
 "I'm taking care of a 21 year old male with fever and abdominal pain..."

Background: pertinent and brief information related to the situation

"The labs are normal and CXR shows no infiltrate but her pulse is elevated..."
 "I have performed a sterile speculum exam and there is frank blood in the vault..."
 "The patient's CT show appendicitis and the patient is vomiting all intake..."

Assessment: analysis and consideration of options, what you found/think

"I think she needs a fluid bolus but I am wondering if she also needs a UA..."
 "I think this patient might have an active abortion..."
 "I think this patient has appendicitis and needs to be transferred to ANMC..."

Recommendation: action requested, what you want

"I want your opinion on how much fluid and the need for a UA..."
 "I want you to come in and assess this patient in person..."
 "I would like to transfer this patient via medevac to ANMC..."

Note about Disagreements

If there is a disagreement regarding the management of a patient and a consensus cannot be reached, a third opinion shall be obtained. This can either be from another YKHC provider or from a provider from another facility. At any time, the Clinical Director on call can also be notified to assist.

This protocol is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
 Approved by Clinical Guideline Committee 8/23/23.
 Click [here](#) to see the supplemental resources for this guideline.
 If comments about this protocol, please contact
 Ellen_Hodges@ykhc.org.



EMERGENT Consults

- Need a call back immediately.
- Examples: Child is in status epilepticus or impending respiratory failure.
- Send priority message via Tiger Connect to Peds Wards on Duty using format below.

URGENT Consults

- Need a call back within one hour.
- Examples: Advice on antibiotic choice or questions about a rash.
- Send message via Tiger Connect to Peds Wards on Duty using format below.

NOT URGENT Consults

- Question can wait until the end of the day/next morning.
- Examples:
 - "Noted that weight percentile has decreased by >2 major percentiles on weight growth chart. Forwarding note to pediatrician for recommendations on further work-up and management for failure to thrive."
 - "During this WCC, reviewed PMH and noted child has not seen neurologist in several years and is off antiepileptics. Forwarding note to pediatrician for recommendations on further management of seizure disorder."
- Do not send a message via Tiger Connect.
- Complete note and forward to "Chronic Peds, RMT" box via Message Center. Note MUST include a specific question for the pediatrician in the plan.
- Note reviewed by inpatient pediatrician, who will addend the note with recommendations and send it back. It will be addressed with the same triage principles we use to prioritize RMT. Goal will be response by the end of the day, but if there is critical care, the night pediatrician will address it by the next morning.


Tiger Connect Message Format for **EMERGENT** and **URGENT** Consults

1. Urgency of consult: need call back ASAP or within one hour.
2. Name of provider, location, role, and phone number.
3. Name and MRN/DOB of patient.
4. One-liner about patient. Here are some examples:
 - "4 yo girl with h/o seizures here for prolonged seizure."
 - "3 month old boy with h/o respiratory failure requiring ICU care here with increased work of breathing."
5. Specific question. Here are some examples:
 - (**EMERGENT**) "The seizure is now >5 minutes and needs medication to stop it. What drug and dose should I give?"
 - (**EMERGENT**) "This child has a RR of 80 and hasn't improved with albuterol or nasal suction. I would like to discuss if a medevac is appropriate."
 - (**URGENT**) "I think this child needs antibiotics, and I'd like to discuss an appropriate choice."
 - (**URGENT**) "This child may require further evaluation in Bethel, and there is a commercial flight landing in two hours. I'd like to discuss whether the child should be sent to Bethel on that flight."

NOTE: If true emergency, limit message to #2 and #4.

How to Send a Priority Message in Tiger Connect

Click the exclamation point to the left of the message box. It will turn red.



This protocol is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 8/23/23.
Click [here](#) to see the supplemental resources for this guideline.
If comments about this protocol, please contact Leslie_Herrmann@ykhc.org.




Miscellaneous

- Goal is guidelines are to be reviewed every two years with revisions and updates as appropriate. Updates may happen sooner if needed.
- If a guideline has not been reviewed in the past five years, it will be decommissioned until it is revised.
- Deadlines for feedback will generally be a period of 1-2 weeks.
- At any time, anyone may send feedback on a guideline. This feedback will be saved for the next guideline revision.
- Minor changes including (but not limited to) correction of typos, changes in test names, small additions, updating hyperlinks, and changes in contact information may be made and published without formal committee vote.

Wiki Supplements

- The long-term goal for the guidelines is for every guideline to have a corresponding supplement page on the wiki.
- The guideline will be information needed to take care of a patient in the moment.
- The wiki supplement will include references, resources, historical background, past versions, and other information.



The Clinical Guideline Committee is a formal sub-committee of MSEC. The policy and procedure outlining its processes can be found [here](#).

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 8/2/24.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline

Process to Update the EHR to Match Guidelines

Contact

If any members of the medical staff identify orders that are discrepant with an approved clinical guideline, they should email Clinical_Guidelines@ykhc.org. The Clinical Guideline Committee will review the request and begin the process outlined here.

A clinical guideline is created or updated.

Point person for guideline reviews all related Power Plans and orders in the EHR.

Point person sends the guideline and relevant Power Plans and orders to Super User with summary of needed changes.

Super User makes screenshot of Power Plans and orders in CERT prior to making changes.

Super User makes requested changes in CERT.

RAVEN team reviews changes in CERT.

Point person for guideline validates changes.

RAVEN team moves changes to PROD.

RAST committee is notified of changes.

Rationale

- The YKHC Clinical Guidelines are the agreed-upon standard of care for the YKHC medical staff.
- This standard of care should be reflected in the available orders and Power Plans in the EHR.
- As such, if orders in the EHR do not match a clinical guideline, these orders may be changed without getting approval from the RAST committee. The RAST committee will be notified of these changes.
- This guideline outlines the process by which EHR changes based on updates in clinical guidelines may be made.

Definitions

- CERT: domain for testing changes to the EHR.
- PROD: Live domain used by staff to access the EHR.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/24/23.

If comments about this guideline, please contact Clinical_Guidelines@ykhc.org.



Contact Poison Control at 800-222-1222 for:

- Pediatric patients <12 years of age.
- Polysubstance or Tylenol PM overdose.
- Any questions or advice.

Patient presents with acute acetaminophen ingestion.

Intentional overdose

Notify BH on call.

Hours post-ingestion

< 4 hours

> 4 hours or unknown

- Administer activated charcoal 1 gram/kg (max 50 grams) if airway intact.
- If ingestion is suspected to be >150 mg/kg or >7.5 grams, start acetylcysteine 30-60 minutes after activated charcoal.

Draw labs
4 hours
post-ingestion.

- Draw acetaminophen level, CMP, and PT/INR.
- Consider starting acetylcysteine while awaiting results.
- If suspected intentional and/or polysubstance ingestion, expand work-up to include UDS, salicylate level, alcohol level, pregnancy test, VBG, lactate, and EKG (if appropriate).

Indications for Starting N-acetylcysteine (NAC)

- Acetaminophen level above treatment line on nomogram.
- Suspected ingestion >150 mg/kg or >7.5 grams total in patient who will not get labs within eight hours of ingestion.
- Acetaminophen level >10 mcg/mL and time of ingestion unknown.
- Possible acetaminophen ingestion and ANY evidence of liver injury.

Indications for Stopping N-acetylcysteine (NAC)

All must be true.

- 21 hour IV protocol is complete or patient has received at least 21 hours of oral therapy.
- Acetaminophen level undetectable.
- ALT trending down (less than 50% of peak level).
- Patient is asymptomatic (especially no RUQ pain).

Note: If polysubstance ingestion, other substances such as diphenhydramine may affect absorption, so this guideline may be less relevant. Consult Poison Control.

Behavioral Health
evaluation if
intentional; otherwise
discharge.

Does patient meet criteria for NAC? (See box.)

Yes

Admit to hospital for IV or PO acetylcysteine.
Consider consult with toxicologist/Poison Control.

Discontinue NAC.


Indications
to stop
NAC met

Recheck acetaminophen
level, CMP, and INR Q12h.

Indications to stop NAC
not met after 21 hours of therapy

- Continue NAC at 6.25 mg/kg/hour IV or 70 mg/kg PO Q4h.
- If ALT >1000 or trending up, INR >1.5 or trending up, or patient worsening, consult Poison Control and/or hepatologist via ANMC, SCH, or UW Medcon. Consider transfer.

Rumack-Matthew Nomogram for
Single Acute Acetaminophen Poisoning



Village Management

- Administer activated charcoal 1 gram/kg (max 50 grams) if airway intact and <4 hours since ingestion.
- If patient in village and toxicity is at all possible, start treatment with oral acetylcysteine and draw blood at 4 hours post ingestion. Instruct health aide to draw 2 mL (minimum 200 microliters) in a gold/SST or green top tube.
- Transport patient and blood work to Bethel on next available commercial flight, if stable.

For vomiting:

- If within one hour of NAC dose, repeat full dose.
- May give ondansetron or metoclopramide.

N-Acetylcysteine (NAC) Administration Protocols

- 1. IV 21 Hour Protocol:** Dose is 150 mg/kg (max 15 grams) over 60 minutes immediately followed by 50 mg/kg (max 5 grams) over 4 hours immediately followed by 100 mg/kg (max 10 grams) over 16 hours (6.25 mg/kg/hour). Dilute with D5W or ½ NS. See [Dose Calculator](#) for details on dose and dilution, especially in children under 40 kg. Note: calculator defaults to pounds.
- 2. PO 72 Hour Protocol:** Dilute with strongly-flavored juice or soda. Mix one part medication with three parts juice/soda. Loading dose is 140 mg/kg. Maintenance dose of 70 mg/kg Q4h for up to 72 hours. The villages carry vials of inhalation/oral solution that is 200 mg/mL in 30 mL vials. See [this resource](#) for details on dosing, including diluent and dosing volumes for weight.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 4/28/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



Box 1: Immediate Interventions

- Supplemental oxygen *pm* to maintain SpO₂ 90-96%.
 - Aspirin 324 mg PO chewed (four 81 mg "baby" aspirin).
 - Nitroglycerin 0.4 mg sublingual *pm* pain (up to three times as BP permits) unless contraindicated.
- Contraindications: recent phosphodiesterase use, sBP <90, right ventricular infarct (consider when evidence of inferior wall ischemia).

NOTE: pain relief with nitroglycerin (or lack thereof) is not diagnostic of cardiac ischemia.

Symptoms suggestive of acute coronary syndrome

Perform 12 lead EKG.
If patient in a village, see Box 3.

Perform immediate interventions. See Box 1.

Tiger Text
ANMC
Cardiology On-
Call with picture
of EKG

STEMI?
See Box 2.

No

- HS-cTnT (high sensitivity troponin), serial EKGs.
- Consider critical diagnoses. See Box 5.
- Consider P2Y12 inhibitor
- Consider morphine if pain not relieved by nitro and no contraindications

Diagnostic EKG or
HS-cTnT findings?
(Box 2 & 4)

No

Tiger Text ANMC
Cardiology On-Call
with picture of EKG.

Consulting Cardiology

- For all STEMI patients, consult PAMC Cardiology by calling the PAMC ED at (907) 212-3433 and asking for the cardiologist on call. For beneficiary patients, ANMC Cardiology should be made aware of the transfer on a non-urgent basis.
- For NSTEMI-ACS patients, consult ANMC Cardiology for beneficiary patients and PAMC Cardiology for non-beneficiary patients.

Box 2: EKG Criteria

STEMI:

- ST elevation in 2 contiguous leads of >0.2mV in V2-V3 OR >0.1mV in all other leads
- New or presumably new LBBB
- Positive Sgarbossa criteria for pre-existing LBBB

High risk Non-ST elevation ACS (NSTEMI-ACS):

- Dynamic T wave inversions
- Transient ST elevation

Box 4: HS-cTnT Evaluation for Acute Cardiac Injury

The lowest reported value is "<6 ng/L," which equates to "undetectable." FDA-approved normal values (99th percentiles in healthy subjects) are:

- Men: <22
- Women: <14
- Change in one hour ($\Delta 1h$): <3

Cutoffs are arbitrary and do not correspond to any evidence-based positive-predictive value for ACS.

For patients with elevated troponins and clinical history consistent with ACS, consult cardiology. This information is from data available February 2020. Please see [wiki page](#) for further information.

Disclaimer

- This algorithm is not intended for undifferentiated chest pain without an apparent cause.
- Acute coronary syndrome is defined as acute occlusion of a coronary artery and does not include type 2 MI/demand ischemia.

<12 hours
from symptom
onset?

Yes

Complete [Fibrinolytic Checklist](#).
Contraindications to fibrinolytics?

No

**Initiate fibrinolytic therapy.
Do not delay fibrinolytics while
awaiting troponin in STEMI.
See next page for dosing.**

Yes

- Administer additional medications. See table on [next page](#).
- Activate medevac if appropriate.

High risk NSTEMI-ACS

Low/Intermediate risk for NSTEMI-ACS

- Broaden differential diagnosis.
- Consider a validated risk-stratification scoring tool (like [GRACE](#) or [TIMI](#)).
- If patient is high-risk for coronary disease, consult cardiologist for discharge and follow up recs, including timing and location of stress testing.
- If patient is considered low-risk for coronary disease, secure outpatient follow up to re-evaluate symptoms and optimize primary prevention (i.e. lipid/A1c testing, aspirin).

Box 3: Village Management

- If EKG meets high risk criteria in Box 2, review with ED Physician and activate medevac. Perform interventions in Box 1.
- ED physician coordinates with ANMC/PAMC regarding whether to have LifeMed give lytics and whether to stop in Bethel or ramp transfer to Anchorage.
- If EKG or health aide not available, use clinical history and validated tool such as [EDACS](#) to stratify risk for ACS. Consult with ED Physician and/or CD on call regarding appropriateness of medevac for risk factors alone.

Box 5: Critical Differential Diagnosis

- Aortic dissection
- Tension pneumothorax
- Pulmonary embolism
- Perforated peptic ulcer

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 3/24/23.
Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact clinical_guidelines@ykhc.org.



Nitroglycerin (NTG)
 • **Contraindications:** PDE-inhibitor use, cardiogenic shock, RV infarct, sBP<90, marked tachycardia or bradycardia.
 • **Sublingual dosing:** 0.4 mg SL Q5 minutes up to three doses
 • **IV dosing:** start at 10-20 mcg/min, titrate Q3-4 minutes to typical range 60-100 mcg/min

Beta-Blockers
 • No evidence of benefit from routine immediate beta-blocker.
 • Indicated for HTN and/or ongoing ischemia refractory to NTG.
 • **Contraindications:** cardiogenic shock, RV infarct, symptomatic asthma.
 • **Cautions:** risk for cardiogenic shock (bradycardia, HR>110, sBP<120, age>70, increased time since STEMI onset), inferior MI, controlled asthma.

Emergency Department Medication Summary				
		STEMI <12 hours	STEMI >12 hours	NSTE-ACS
Oxygen		Maintain SpO ₂ 90-96%	Maintain SpO ₂ 90-96%	Maintain SpO ₂ 90-96%
Nitrates (<i>prn</i> pain, HTN)		Sublingual or drip	Sublingual or drip	Sublingual or drip
Fibrinolytic		Tenecteplase See below.	Not indicated	Not indicated
Antiplatelet agents	Aspirin	Aspirin 324 mg PO chewed (four 81 mg "baby" aspirin)	Aspirin 324 mg PO chewed (four 81 mg "baby" aspirin)	Aspirin 324 mg PO chewed (four 81 mg "baby" aspirin)
	P2Y ₁₂ receptor blocker	Clopidogrel Age ≤75: 300 mg PO Age >75: 75 mg PO	Clopidogrel 600 mg PO	Consult cardiology.
Anticoagulation		Enoxaparin (see table for dose)	Enoxaparin (see table for dose)	Enoxaparin (see table for dose)
Beta-blocker		Metoprolol 5 mg IV <i>pm</i> Q5 minutes (max 15 mg)	Metoprolol 5 mg IV <i>pm</i> Q5 minutes (max 15 mg)	Metoprolol 5 mg IV <i>pm</i> Q5 minutes (max 15 mg)
Morphine		No longer routinely given; associated with increased mortality. Reserve for significant pain refractory to NTG and beta-blocker.		

At time of Dx unless contraindicated

Fibrinolytic Therapy (Tenecteplase)

Goal: administer ≤ 30 minutes from arrival.
 Rapidly complete the fibrinolytic checklist and consent.

Dosing:

- <60 kg: tenecteplase 30 mg IV bolus
- ≥60 kg to <70 kg: tenecteplase 35 mg IV bolus
- ≥70 kg to <80 kg: tenecteplase 40 mg IV bolus
- ≥80 kg to <90 kg: tenecteplase 45 mg IV bolus
- ≥90 kg: tenecteplase 50 mg IV bolus

Administer concurrent aspirin, clopidogrel, and anticoagulant therapy, per table above.

Enoxaparin Dosing

	Age <75 years and STEMI	Age ≥75 years and STEMI	Any age and NSTE-ACS
Creatinine clearance ≥30 mL/min	30 mg IV + (1 mg/kg SC now then Q12h) Max dose 100 mg	0.75 mg/kg SC Q12h Max dose 75 mg	1 mg/kg SC now then Q12h
Creatinine clearance <30 mL/min	30 mg IV + (1 mg/kg SC now then Q24h) Max dose 100 mg	1 mg/kg SC Q24h Max dose 100 mg	1 mg/kg SC now then Q24h

NOTE: Enoxaparin and unfractionated heparin are NOT dialyzable; ESRD/dialysis patients should receive fondaparinux, which is not on the YKHC formulary. Discuss with cardiologist if appropriate.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/24/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Megan_Young@ykhc.org and Andrew_Swartz@ykhc.org.



Fibrinolytic Checklist

INDICATIONS (initial yes or no)

YES	NO	
		Presentation consistent with acute coronary syndrome (coronary artery occlusion)
		<p><u>AND</u> at least one of the following:</p> <ul style="list-style-type: none"> • 1 mm J-point elevation in two contiguous leads (other than V₂-V₃) • In leads V₂-V₃ <ul style="list-style-type: none"> Men ≥ 40 years: ≥ 2 mm J-point elevation Men <40: ≥ 2.5 mm J-point elevation Women: ≥ 1.5 mm J-point elevation

ABSOLUTE CONTRAINDICATIONS (initial yes or no)

YES	NO	
		History of <u>any</u> intracranial hemorrhage
		History of prior ischemic stroke, significant closed head injury or facial trauma, or intracranial or spinal surgery in the previous three months
		Presence of a cerebral vascular malformation
		Presence of a primary or metastatic intracranial malignancy
		Symptoms or signs suggestive of an aortic dissection
		Any bleeding diathesis
		Any active bleeding that is severe or has high potential for life-threatening blood loss; this does not include menstrual bleeding
		sBP > 180 and/or dBP >110 at presentation in patient at low risk of cardiac death (age < 55, no prior MI, and Killip class I).
		Terminal illness, defined as end of life care or documented/expressed patient wish to abstain from high risk or invasive procedures

RELATIVE CONTRAINDICATIONS (initial yes or no) – If any of below are present, used shared decision making with patient.

YES	NO	
		Age 65-74 (ICH relative risk 3.12 [2.54-3.83]); Age ≥ 75 years (ICH relative risk 5.40 [4.40-6.63])
		History of chronic severe poorly controlled HTN
		sBP > 180 and/or dBP >110 at presentation in patient at high risk of cardiac death (age ≥ 55, Hx prior MI, or Killip class ≥ II).
		History of ischemic stroke more than three months ago
		Dementia OR any known intracranial disease that is not an absolute contraindication
		Traumatic or prolonged (>10 minutes) cardiopulmonary resuscitation
		Major surgery in the previous three weeks
		Internal bleeding in the previous 2-4 weeks
		Active peptic ulcer
		Non-compressible vascular punctures
		Pregnancy
		Current warfarin therapy (the risk of bleeding increases as the INR increases)

This checklist is advisory for clinical decision-making and may not be all-inclusive. Risks and benefits will need to be assessed individually.

Physician signature: _____

Printed name: _____ Date and time: _____

Place patient ID sticker here.



PROCEDURE CONSENT

I hereby authorize _____ and such assistants as he/she may designate, to perform the following operation or procedure:

TECHNICAL DESCRIPTION

Intravenous thrombolytic therapy for acute STEMI (ST-elevation myocardial infarction).

LAY DESCRIPTION

Give clot-dissolving medication through an IV to dissolve the clot which is causing a heart attack.

_____ has discussed with me the information briefly summarized below:

BENEFITS

- When PCI is not available within two hours, thrombolytic medication is the "standard of care" for achieving coronary reperfusion within 12 hours of acute STEMI onset.
- When administered within 6 hours of pain onset, about 1 in 40 persons will have their life saved.
- When administered between 6-12 hours after pain onset, about 1 in 60 persons will have their life saved.
- Decreased risk of developing heart failure.
- A STEMI patient who receives thrombolytic medication is about 3-5 times more likely to have their life saved than to have brain bleeding (see below).

RISKS (some, but not all)

- About 1 in 100 persons will experience non-life-threatening bleeding.
- About 1 in 100-250 persons will experience bleeding into the brain which usually results in either death or significant disability.

RISKS OF NOT HAVING THE PROCEDURE

- Higher risk of death.
- Higher risk of developing heart failure.

ALTERNATIVE TREATMENTS

None are available at this facility.

Patient signature: _____

Printed name: _____ Date and time: _____

Witness signature: _____

Printed name: _____ Date and time: _____


Physician signature: _____

Printed name: _____ Date and time: _____


Witness signature: _____

Printed name: _____ Date and time: _____

Place patient ID sticker here.



Lund and Browder Chart to Estimate TBSA in Adults and Children



Age		0	I	5	10	15	Adult
Front or back half	(%)	(%)	(%)	(%)	(%)	(%)	(%)
I	(Head)	9½	8½	6½	5½	4½	3½
II	(Thigh)	2¼	3¼	4	4¼	4½	4¼
III	(Leg)	2½	2½	2¼	3	3¼	3½

Consensus Formula (Brooke/Parkland)

Only used if TBSA >15%.

(weight in kg) x 2-4 mL x %TBSA = total fluid to be given over 24 hours
Do not convert %TBSA to a decimal. For example, 15% TBSA would be 15.

Give half in first eight hours from time of burn. Give other half over the next sixteen hours.
If delayed presentation, begin at initial calculated 8 hour rate; do not "catch up" with fluid boluses.
Goal UOP 0.5-1 mL/kg/hour.


Use LR used for adults unless mitigating circumstances. For pediatric patients <30 kg, add D5.

Classification of Burns by Depth

Burns evolve over time; initial TBSA and depth classification can change and often the difference between deep partial thickness and full thickness can only be determined operatively.

- Superficial (1st degree): epidermis only, dry, red, blanches with pressure, no blisters, painful.
- Superficial partial-thickness (2nd degree): epidermis and part of dermis, blisters, moist, red, weeping, blanches with pressure, painful.
- Deep partial-thickness (2nd degree): epidermis and deep dermis, blisters, wet or waxy dry, patchy white to red, does not blanch, pressure sensation only.
- Full-thickness (3rd degree): epidermis and entire dermis, waxy white to leathery gray to charred/black, dry and inelastic, does not blanch, sensation to deep pressure only, may be defined as 4th degree with extension into underlying fascia, muscle, or bone.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 4/28/23. Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact Travis_Nelson@ykhc.org.





Thrombolytic Checklist

INDICATIONS (initial yes or no)

YES	NO	
		Less than 4.5 hours since onset of symptoms or last known normal.
		NIHSS greater than 5 (or less than 5 with disabling symptoms).
		Symptoms are NOT rapidly improving.
		Symptoms are NOT due to untreated hypoglycemia (BG<50).

ABSOLUTE CONTRAINDICATIONS (initial yes or no)

YES	NO	
		CT evidence of hemorrhage OR extensive area of hypodensity (irreversible injury).
		GI/GU bleed in the last 21 days.
		Severe, uncontrolled, hypertension >185/110.
		Current intracranial neoplasm.
		Active internal bleeding or known aortic dissection.
		Any bleeding diathesis.
		Presentation suggestive of SAH or endocarditis (not septic emboli).
		History of intracranial hemorrhage.
		Anticoagulation (warfarin or DOAC in the last 48 hours or therapeutic-dosed heparinoids).
		Any of the following in the last three months: ischemic stroke, intracranial surgery, intraspinal surgery, or serious head trauma.

RELATIVE CONTRAINDICATIONS (initial yes or no) – If any of the following relative contraindications are present, consider expert consultation prior to giving thrombolytic and/or consider these with consent and shared decision-making.

YES	NO	
		History of GI or GU hemorrhage.
		Arterial puncture in a non-compressible site in the last seven days.
		Seizure at onset with postictal neurologic impairment.
		Major surgery in the last 14 days.
		Pregnancy.
		Onset 3-4.5 hours with NIHSS >25 (higher bleeding risk) or age >80 (higher bleeding risk).
		Untreated AVM or aneurysm.
		Systemic malignancy.
		History of arterial dissections.
		Blood glucose greater than 400 (associated with worse outcomes).

This checklist is advisory for clinical decision-making and may not be all-inclusive. Risks and benefits will need to be assessed individually.

Physician signature: _____

Printed name: _____ Date and time: _____

Place patient ID sticker here.



PROCEDURE CONSENT

I hereby authorize _____ and such assistants as he/she may designate, to perform the following operation or procedure:

TECHNICAL DESCRIPTION	Intravenous thrombolytic therapy for acute ischemic stroke.
LAY DESCRIPTION	Give clot-dissolving medication through an IV to dissolve the clot which is causing a stroke.
_____ has discussed with me the information briefly summarized below:	
BENEFITS	<ul style="list-style-type: none"> • Thrombolytic medication is a treatment that may restore blood flow to the brain. • In studies, if these drugs were given less than three hours after the stroke started, 33% of patients given thrombolytic drugs had a good outcome. In patients who did not get thrombolytic drugs, 23% got better. Ten people would have to get the drug to help one person have a better outcome. • If these drugs were given between three and four and a half hours after the stroke started, 35% of patients given thrombolytic drugs had a good outcome, and 30% of patients who didn't get the drug also got better. Twenty people would have to get the drug to help one person have a better outcome. • Patients who receive this drug within three hours of the stroke starting have a 10% increase in chance of improved neurologic function. • Patients who receive this drug between three and four and a half hours from the stroke starting have a 5% increase in chance of improved neurologic function.
RISKS (some, but not all)	<ul style="list-style-type: none"> • In a large study of stroke patients, 6.8% of them had bleeding in their brain after receiving thrombolytic drugs for stroke, compared to 1.3% of those stroke patients who did not receive the drug. If we give this drug 18 times, it will probably make one person have bleeding in their brain. • Among all people given this drug, 2% die from a hemorrhage.
RISKS OF NOT HAVING THE PROCEDURE	<ul style="list-style-type: none"> • Higher risk of developing permanent, disabling stroke symptoms.
ALTERNATIVE TREATMENTS	No other treatments available at this facility. Only monitoring symptoms and rehabilitation.

Patient signature: _____

Printed name: _____ Date and time: _____

Witness signature: _____

Printed name: _____ Date and time: _____

Physician signature: _____

Printed name: _____ Date and time: _____

Witness signature: _____

Printed name: _____ Date and time: _____

Place patient ID sticker here.



Patient with serious illness with expected death.

Preparation

In hospital:

- Complete [Physician Orders for Life-Sustaining Treatment \(POLST\)](#) order form. Review with patient and family regularly.
- Review DNR/DNI status at least once an admission. Remember, all decisions regarding end-of-life care may be modified at any time per patient and family wishes.
- Place DNR/DNI order in RAVEN. Update code status on RAVEN banner by going to Ad hoc → Code Status form.
- When discharging home, ensure all support is in place, including family care plan, comfort meds (consider sublingual morphine and lorazepam), incontinence supplies, etc.

In village:

- Discuss and document goals of care, code status, wishes for medevac/hospitalization with patient and family. Update code status in RAVEN as above.
- Complete [Expected Home Death](#) form and send to AST/BPD.
- Place on RAVEN banner by going to AdHoc → Patient Registries and check off "Expected Home Death."
- Communicate with village health aides.

After a home death has occurred

- Medical providers can pronounce death remotely after speaking with a qualified representative, which includes health aides. Representative must ascertain that there is no heart beat or spontaneous breathing.
- Send [Expected Home Death](#) form to the State Medical Examiner and AST/BPD. If this form was not completed prior to death but would have been indicated, it is acceptable to fill it out after death. This will expedite things for the family.

If this is an expected neonatal death, go to [page 3](#).

Required Notifications

Ensure you have next of kin's name and phone number prior to making these calls.

In hospital:

- Bethel Police Dept 907-543-3781 Even if Expected Death form has been completed.
- Life Alaska 888 543-3287. Required by CMS for all hospital deaths.
- State Medical Examiner 888 332-3273. Please review page 3 for ME notification requirements.

In village:

- Alaska State Troopers 800 478-9112
- State Medical Examiner 888 332-3273 Please review page 3 for ME notification requirements.
- Optional: Life Alaska 888 543-3287. Deceased individuals in villages may still be candidate for tissue donation.

Documentation

- Death Note in RAVEN should be an Alert Note using autotext ".death" which fulfills all documentation requirements.
- Forward death note to Chief of Staff and designated Medical Records representative.
- Complete highlighted portions of Death Certificate and place in Medical Records basket.
- If death occurred in the hospital, complete [Notification of Death](#) form.

Helpful Forms

Note: Copies of the death packet are also kept on the inpatient unit.


- [Physician Orders for Life-Sustaining Treatment \(POLST\)](#)
- [Expected Home Death](#)
- [Death Certificate Worksheet](#)
- [Notification of Death](#)

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/14/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact clinical_guidelines@ykhc.org.



Regarding Life Alaska

It is a CMS and TJC requirement to notify designated organ donation organization (Life Alaska) for all in hospital deaths. We are not mandated to contact Life Alaska for village deaths; however, individuals who die in villages may still be candidates for tissue donation. Additionally, if the death will become an ME case, Life Alaska must be notified.

Helpful Phone Numbers

- Alaska State Medical Examiner: 888 332-3273
- Life Alaska: 907 562-5433
- Alaska State Troopers (AST): 800 478-9112
- Bethel Police Department (BPD): 543-3781
- State Epidemiology: 907 269-8000
- OCS Intake (for reports): 800 478-4444
- APS Intake (for reports): 800 478-9996

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.


Approved by Clinical Guideline Committee 7/14/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact clinical_guidelines@ykhc.org.



Note: This guideline applies to fetuses ≥ 21 weeks. Do not follow this guideline for miscarriage prior to 21 weeks.



Medical Examiner Notification

Per YKHC and State of AK policy, the Medical Examiner must be notified when the death appears to have:


1. Been caused by unknown or criminal means, during the commission of a crime, or by suicide, accident, or poisoning.
2. Occurred under suspicious or unusual circumstances or occurred suddenly when the decedent was in apparent good health.
3. Been unattended by a practicing physician or occurred less than 24 hours after the deceased was admitted to a medical facility.
4. Been associated with a diagnostic or therapeutic procedure.
5. Resulted from a disease that constitutes a threat to public health.
6. Been caused by a disease, injury, or toxic agent resulting from employment.
7. Occurred in a jail or corrections facility owned or operated by the state or a political subdivision of the state or in a facility for the placement of persons in the custody or under the supervision of the state.
8. Occurred in a foster home.
9. Occurred in a mental institution or mental health treatment facility.
10. Occurred while the deceased was in the custody of, or was being taken into the custody of, the state or a political subdivision of the state or a public officer or agent of the state or a political subdivision of the state
11. Been of a child under 18 years of age or under the legal custody of the Department of Health and Human Services, unless the child's death resulted from a natural disease process and was medically expected and the child was under supervised medical care during the 24 hours before the death.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/14/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact clinical_guidelines@ykhc.org.





Alteplase Checklist

INDICATIONS (initial yes or no)

YES	NO	
		Grade 3 or 4 frostbite.
		High risk for life-altering disability.
		Patient able to arrive in Bethel <24 hours from time rewarming complete.
		Patient or guardian able to give informed consent.

ABSOLUTE CONTRAINDICATIONS (initial yes or no)

YES	NO	
		Prior intracranial hemorrhage.
		Known structural cerebral vascular lesion.
		Known malignant intracranial neoplasm.
		Ischemic stroke within three months.
		Suspected aortic dissection.
		Active bleeding or bleeding diathesis (excluding menses).
		Significant closed-head trauma or facial trauma within three months.

RELATIVE CONTRAINDICATIONS (initial yes or no) – If any of the following relative contraindications are present, consider expert consultation prior to giving thrombolytic and/or consider these with consent and shared decision-making.

YES	NO	
		History of chronic, severe, poorly controlled hypertension.
		Severe uncontrolled hypertension on presentation (SBP >180 mmHg or DBP >110 mmHg)
		History of ischemic stroke more than three months prior
		Traumatic or prolonged (>10 minute) CPR or major surgery less than three weeks
		Recent (within two to four weeks) internal bleeding or recent invasive procedure or serious trauma.
		Noncompressible vascular punctures.
		Pregnancy.
		Active peptic ulcer GI malignancy, GI hemorrhage in previous 21 days, h/o GI bleed.
		Pericarditis or pericardial fluid.
		<ul style="list-style-type: none"> • Therapeutic LMWH. • Current use of any anticoagulant that has produced an elevated INR >1.7 or PT >15 seconds or abnormal PTT.
		Age >75 years.
		Diabetic retinopathy.
		Platelet count <100,000.

This checklist is advisory for clinical decision-making and may not be all-inclusive. Risks and benefits will need to be assessed individually.

Physician signature: _____

Printed name: _____ Date and time: _____

Place patient ID sticker here.



YKHC follows the statewide “Guidelines for the Management of Acute Blunt Head Trauma in Alaska,” found [here](#).

If concern for spinal injury, see YKHC [Spinal Cord Injury Management](#) guideline.

Village Management of Isolated Head Trauma


- See [Canadian CT Head Injury/Trauma Rule](#).
- If GCS 14-15 and no high risk criteria met, observe in clinic for two hours. If worsening or no improvement, consider transfer to Bethel in consultation with ED physician.
- If GCS 9-13 or 14-15 with high-risk criteria, consult Emergency RMT provider or ED physician.
- If GCS ≤ 8 , activate medevac and support airway per ATLS.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 11/21/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Clinical_Guidelines@ykhc.org.



Village Management

- If Low Risk: Discharge with competent caregiver with clear return precautions. Do not send to Bethel unless otherwise indicated.
- If Intermediate Risk: Consider medevac vs observation with Q1h VS and neuro checks. If any worsening or no improvement over 4 hours, activate medevac.
- If High Risk: Activate medevac.

Plain films of the skull are not recommended.

C-spine Injury

Please see the [YKHC Spinal Cord Injury Management guideline](#) for pediatric C-spine resources.

Concussion Management

- Complete [Acute Concussion Evaluation](#) at every visit.
- Follow-up in outpatient clinic in 1-2 weeks.
- Consider [Sport Concussion Assessment Tool \(SCAT\)](#) at follow-up.
- Consider balance testing.
- Avoid medications that can worsen somnolence.
- If symptoms persist >3-4 weeks, consider referral to neurologist, psychologist, physical therapy, etc.
- Return to school per [CDC Heads Up Protocol](#).
- Return to play per [ASAA Guidelines](#).

Pediatric Glasgow Coma Scale (GCS)

	Infant	Child	
Eye opening	Spontaneous	Spontaneous	4
	To speech	To speech	3
	To pain	To pain	2
	No response	No response	1
Best verbal response	Coos, babbles	Orientated, appropriate	5
	Irritable cry	Confused	4
	Cries to pain	Inappropriate words	3
	Moans to pain	Incomprehensible sounds	2
Best motor response	No response	No response	1
	Moves spontaneously	Obeys commands	6
	Withdraws to touch	Localizes painful stimulus	5
	Withdraws to pain	Withdraws to pain	4
Best motor response	Flexion to pain	Flexion to pain	3
	Extension to pain	Extension to pain	2
	No response	No response	1

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 5/15/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Clinical_Guidelines@ykhc.org.

**REMEMBER:**

- No pediatric patient may be kept at YKDRH on HFNC unless medevac is on weather-hold.
- Maintain patient on HFNC until medevac arrival.
- Requirements for HFNC:
 - The patient must have 1:1 nursing care until stabilized. After stabilization, nursing care may be 2:1 until medevac arrival.
 - The patient must have a respiratory therapist at bedside until stabilized.
- All newborns on HFNC must remain in the nursery.

Apnea

If patient has apnea with poor or worsening response to stimulation, prepare for intubation.

Flow Rates

- Titrate flow to 0.5-2 LPM/kg.
- Younger patients often require higher flow rates per kilogram.
- Consult the PICU for any patient requiring > 1.5 LPM/kg.
- Listen to lungs with each adjustment. If child is unable to easily exhale or complete an exhalation, decrease flow rate until exhalation is adequate.

Troubleshooting

- Consider NG/OG-tube for decompression.
- Use a pacifier to keep the patient's mouth closed and prevent loss of pressure. Consider Sweet-Ease.
- Try environmental changes to comfort a fussy baby: caregiver may hold patient in semi-recumbent position, patient may be swaddled, patient may be fanned if hot, lights may be dimmed, etc.
- Consider mild anxiolysis in consultation with medical control.
- Consider higher levels of flow to improve washout.

Patient with moderate to severe sustained retractions or sustained hypoxia <88% not improved with **SUPPORTIVE MEASURES** (see box) and 2 LPM conventional nasal cannula or infant with apnea responsive to stimulation. (See box.)

- Page respiratory therapist.
- Page pediatric hospitalist.

- Activate medevac.
- **PREPARE PATIENT.** (See box.)

- RT to start high-flow nasal cannula with pediatrician consultation.
- **Low-flow cartridge** to be used with neonatal/infant cannula and produces flow rates of 1-8 LPM. This should only be used in patients ≤ 4 kg.
- **High-flow cartridge** to be used with larger cannula and produces flow rates of 5-40 LPM.

Initial Settings

See Flow Rates box to left.

FiO2 50%, 37°C.

For newborns, consult neonatologist.

Titrate flow by 1 LPM increments over first 3 minutes until improvement.
If patient is worsening on high flow rates, consider a lower flow rate.

Titrate FiO2 to maintain sats 94-99%.

Frequent gentle nasal suction as needed.

Reassess at least every 20-30 minutes.

Signs of Clinical Improvement

- ↓HR (often noted first)
- ↓RR
- ↓retractions
- ↓irritability
- improved air movement
- decreased apnea

Maintain current settings until medevac arrives.

If no improvement, consider obtaining ANMC PICU consult, checking blood gas, increasing supportive measures, intubation, etc.

SUPPORTIVE MEASURES

- Control fever, as it can be an independent cause of respiratory distress.
- Nasal suction ± nasal saline or saline nebs.
- IV hydration.
- Consider back-to-back or continuous albuterol.
- Consider phenylephrine 0.25%, 1 spray to each nostril once.

PREPARE PATIENT

- Make patient NPO.
- Ensure reliable IV access.
- Suction nares well.
- Choose a nasal cannula with prongs that do not occlude more than 50% of the nares.
- Position patient: optimal patient position is semi-recumbent, not supine or upright. Consider using blue seat (stored in the ED) with adjustable angle. Use blankets and towels for shoulder rolls and to support position and ensure patient is not slumping over. Caregivers may hold the child if it helps keep him/her calm as long as the child is at a ~45 degree angle.
- To prevent condensation causing problems, place patient at a higher level than unit and clip tubing to patient's clothing.
- Consider transcutaneous CO2 monitoring.

Conditions in Which to Use with Caution

- Hypercapnic respiratory failure
- Facial anomalies (including choanal atresia) or injuries that preclude appropriate NC fit
- Active vomiting
- Bowel Obstruction
- Air leak (including pneumothorax or pneumomediastinum)


Use care in patients with excessive secretions; ensure good suction during use.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Click [here](#) to see the supplemental resources for this guideline.

Approved by Clinic Guidelines Committee 7/23/25.

If comments about this guideline, please contact
Amy_Carson-Strnad@ykhc.org.





Intubation (Adult and Pediatric)

Predict difficult airway and post-intubation hypotension (See right for Push-Dose Epinephrine):

1. Shock Index (SI) = HR/sBP. If >0.8, high likelihood of post-intubation hypotension. Optimize resuscitation with fluids/blood prior to intubation if possible.
2. Have at bedside: BVM with correct size mask, OPA/NPA, suction, LMA/i-Gel, bougie, push-dose epinephrine (see below).
3. Predictors of difficult intubation: see next page. **Consider contacting anesthesiology if difficult airway is anticipated!**
4. Prepare several doses of all medications, including an extra dose of paralytic and boluses of post-intubation sedation.
5. Optimize positioning: shoulder roll for infants and children, ramp positioning for obese patients.

Use [Adult Critical Care Guide](#) for more information about medications.

ADULTS, may choose one or more:

1. Fentanyl 2 mcg/kg
2. Ketamine 0.1-0.3 mg/kg
3. Rocuronium 0.1 mg/kg

ADULTS, choose one agent. Reduce doses if risk of hypotension.

1. Etomidate 0.3 mg/kg; first line for most patients.
2. Ketamine 1-2 mg/kg.
3. Propofol 2 mg/kg.
4. Midazolam 0.1-0.3 mg/kg.

ADULTS, choose one:

1. Rocuronium 1-2 mg/kg; duration 30-40 minutes.
2. Succinylcholine 1-2 mg/kg; duration 3-5 minutes; see warning on next page.
3. Vecuronium 0.1 mg/kg; duration 45-65 minutes.

Prepare

PUSH-DOSE EPINEPHRINE

- Mix 9 mL of NS with 1 mL of epinephrine 1:10,000 (or 0.1 mg/mL) to make epinephrine 10 mcg/mL.
- Adult dose is 0.5-2 mL Q 2-5 minutes (5-20 mcg).
- Pediatric dose is 0.1 mL/kg, max 2 mL.

Pre-oxygenate

Goal SpO₂ > 93% for three minutes. Mask and NC in all patients. Non-rebreather with "flush rate" (as high as possible) O₂ if spontaneously breathing. Bag-mask ventilation if no spontaneous breathing. Non-invasive PPV if in ARDS.

Pre-medicate (optional)

PEDIATRICS (<36 kg):

Atropine – onset <1 minute. Use for <1 year or if using ketamine. Doses on [Pediatric Critical Care Guide](#) (code sheets). Consider if at risk for bradycardia with airway manipulation, shock, hypoxia, acidosis, or severe electrolyte abnormalities.

Induce

PEDIATRICS (<36 kg):

Midazolam AND fentanyl, using doses on [Pediatric Critical Care Guide](#) (code sheets). May need to repeat fentanyl dose.

Paralyze

PEDIATRICS (<36 kg):

Rocuronium, using doses on [Pediatric Critical Care Guide](#) (code sheets). **DO NOT USE SUCCINYLCHOLINE.**

Perform video or direct laryngoscopy

Options:

- Bag-mask ventilation ± OPA or supraglottic airway (LMA/i-Gel)
- Bougie.
- Other form of laryngoscopy, including intubating bronchoscopy.
- Change provider doing the procedure.
- Contact anesthesia.
- Change patient position.
- Cricothyrotomy (last resort and should be avoided <10 years unless no other options).

Succeeds

Confirm placement:

1. Visualize tube through cords.
2. End tidal CO₂ detector turns and remains yellow after 6 breaths.
3. Bilateral breath sounds.
4. No noise over epigastrium.
5. Normal waveform on EtCO₂ capnography (gold standard).
6. Confirmation on CXR.

PEDIATRICS:

- Secure tube with silk tape. Do not use a commercial tube holder device for tubes 5.0 and smaller.
- Always place NG/OG tube for decompression.
- Pre-oxygenate for full two minutes between attempts.

Continuing sedation/analgesia

ADULTS:

1. Fentanyl 1-4 mcg/kg/hour, bolus 50-100 mcg.
2. Midazolam 0.02-0.1 mg/kg/hour, bolus 1-5 mg.
3. Propofol 5-80 mcg/kg/minute, bolus 1-2 mg/kg.
4. Ketamine 0.5 mg/kg/hour, bolus 1-2 mg/kg.

PEDIATRICS:

- Fentanyl continuous infusion with boluses Q15minutes *pm*, using doses on Critical Care Guide. Give bolus prior to starting drip.
- Consider midazolam boluses if needed. Consider continuous infusion if needed. Watch for hypotension.

Post-intubation care

- Obtain ABG.
- Place NG/OG tube.
- Elevate head of bed 30-45°.
- BP Q3 minutes x30 minutes.
- Place in-line suction.
- Consider nebulized treatments.
- Consider C-collar.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 10/23/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact

Travis_Nelson@ykhc.org or Leslie_Hermann@ykhc.org.



Predictors of Difficult Intubation

Predictors of Difficult Intubation

- Mallampati grade 3 or 4
- Cormack & Lehane grade 3 or 4
- Wilson score of > 2
- LEMON system; objective/subjective scoring

Wilson Score

	0	1	2
Weight (kg)	< 90	90-110	> 110
Head and neck movement	> 90°	~ 90°	< 90°
• Inter-incisor gap (cm) • SL (maximum forward protrusion of lower incisors beyond uppers)	> 5 > 0	= 5 = 0	< 5 < 0
Receding mandible	None	Moderate	Severe
Buck teeth	None	Moderate	Severe

LEMON System

L	Look: trauma, large tongue
E	Evaluate 3:3:2 rule.
M	Mallampati score ≥3
O	Obstruction
N	Neck mobility (limited)

Helpful Resource: [the Difficult Airway App](#)


Difficulty with BVM

Predictors of Difficulty with BVM

R	Radiation/Restriction
O	Obstruction/Obesity/OSA
M	Mask seal/Male/Mallampati ≥3
A	Aged
N	No teeth

Options if having difficulty with BVM

- 2-hand technique with 2 providers
- Oral/nasal airways
- Positioning
- Consider no paralytics



Paralytics

Succinylcholine

Absolute contraindications:

- Family / personal history of malignant hyperthermia
- Hyperkalemia; if unknown K, obtain EKG for peaked T's
- Upper motor neuron injury, denervating neuromuscular disease
- Use after acute phase of burns, major trauma, crush injury

Relative contraindications:

- Elevated ICP
- Pseudocholinesterase deficiency

Treatment of malignant hyperthermia:

Dantrolene 2.5 mg/kg IV, redosing based on expert guidance

Avoid in pediatric patients.

Rocuronium

Note: Incidence of rocuronium IgE-induced anaphylaxis is estimated at 1:2500. Consider if sudden cardiovascular collapse after giving rocuronium.



ADULTS: ARDS/Protective Ventilation Protocol (appropriate for most patients without indication for alternate ventilation)

Initial Ventilator Settings:

1. Set Tidal volume (Vt) = 6-8 mL/kg using Ideal Body Weight. See [MDCalc Tidal Volume Calculator](#).
2. Reduce Vt by 1 mL/kg every 1-2 hours until Vt 6 mL/kg.
3. Set initial rate to 18-35 bpm based on pre-intubation rate.
Obstructive lung disease: Consider lower RR to maximize expiratory phase.
4. Set initial PEEP at 5 cm H₂O.
• If BMI > 30, set PEEP to 8 cm H₂O.
• If BMI > 40, set PEEP to 10 cm H₂O.
5. Set initial FiO₂ at 30-40%; adjust to SpO₂ 88-95%.
6. Set inspiratory flow rate 60-80 lpm.
Obstructive lung disease: Consider inspiratory flow rate 80-100 lpm

Check BP immediately after any major changes in vent settings.

Adjust settings based on patient status, blood gases, CXR, and expert consultation.

Oxygenation goal: PaO₂ 55-80 mmHg or SpO₂ 88-95%.

Use a minimum PEEP of 5 cm H₂O. Consider use of incremental FiO₂/PEEP combinations such as shown below (not required) to achieve goal.

PEDIATRICS: Suggested Starting Ventilator Settings

1. Choose mode SIMV.
2. Set FiO₂ to 1.0 and titrate to maintain SpO₂ 92-94%. Goal is to decrease FiO₂ to <0.5 if possible.
3. Set Tidal Volume (Vt) at 8-10 mL/kg. If concern for ARDS, set Vt to 6-8 mL/kg.
4. Goal is inspiratory plateau pressures <30 cm H₂O.
5. Set respiratory rate by age, increasing or decreasing based on disease process:
Adolescents 12-15 breaths/minute
Children 15-20 breaths/minute
Infants 20-25 breaths/minute
Neonates 25-30 breaths/minute
6. Set PEEP to 5 cm H₂O to optimize alveolar recruitment.
7. Set inspiratory time by age:
Adolescents 1.0 second
Children 0.7 second
Infants/neonates 0.5 second
8. If using pressure support, set at 5-10 cm H₂O.
9. Get a blood gas ~30 minutes after any changes to ventilator settings.

Check BP immediately after any major changes in vent settings.

Call PICU at (907) 297-8809 immediately to help troubleshoot any problems. Low threshold to use Zoom.

For All Modes of Ventilation

- Initial vent setting are based on patient presentation.
- Vent settings are adjusted based on patient tolerance of mechanical ventilation and ABG results.
For high PCO₂: increase rate and Tidal Volume
For low PO₂: increase FiO₂ and PEEP
- Obtain ABG prior to intubation, 30 minutes following intubation, and 30 minutes after vent changes.
- Goal plateau pressure < 30 cm H₂O; decrease Vt to lower plateau pressure.
Obese patients may require higher plateau pressure.
- Target pH > 7.30; increase RR to control hypercapnia.
- Avoid intubation if possible in patients with obstructive lung disease; maximize use of NIPPV.

Check BP immediately after any major changes in vent settings.

Extubation

If considering extubation in the Emergency Department, see [this algorithm](#) and [this resource](#).

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 10/23/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact

Travis_Nelson@ykhc.org or Leslie_Herrmann@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline

Intubation Preparation Checklist

(for use outside the OR)

Risk Assessment

- ☐ Assess risk of difficult airway.
- ☐ Assess risk of physiologic compromise.
- ☐ Discuss if CRNA is needed.

Equipment

Monitoring

- ☐ Pulse-oximetry
- ☐ ET CO₂
- ☐ BP set to cycle Q1min
- ☐ Cardiac leads

Equipment

(All located in the code cart.)

- ☐ Wall: Suction, BVM with O₂ flowing, NC turned as high as possible ("flush rate")
- ☐ Intubation tray:
 - ETT x2 (including anticipated size and one smaller)
 - Ensure balloon has been tested.
 - 10 mL syringe
 - Stylet
 - Laryngoscopes x2 (direct and video)
 - ET CO₂ detector
 - Stethoscope
- ☐ NG/OG tray:
 - NG/OG tube
 - Toomey syringe
 - Anti-reflux valve
- ☐ Know locations of adjuncts and rescue equipment:
 - OPA/NPA
 - Supraglottic airway device (LMA, etc.)
 - Bougie
 - Needle crich
- ☐ Ventilator

Drugs & Lines

- ☐ Two functioning IVs
- ☐ Induction drug doses
- ☐ Paralytic drug doses
- ☐ ± Push-dose epi.
- ☐ ± Pressors

Post-Intubation Planning

- ☐ Post-intubation sedation
- ☐ Initial ventilator settings
- ☐ Medevac activation

Team Roles

- ☐ Intubator
- ☐ Team lead (may be same as intubator)
- ☐ Respiratory therapist
- ☐ Primary nurse (recorder)
- ☐ Medication nurse
- ☐ Float nurse/tech
- ☐ X-ray tech on stand-by
- ☐ ± CRNA

Plan

Optimize Patient

- ☐ Position
- ☐ Oxygenation with NC as high as possible ("flush rate") AND mask/BVM at max rate
- ☐ Optimize hemodynamics
- ☐ Consider NG tube to suction in infants

Contingencies

- ☐ Secondary airway plan
- ☐ Plan for post-intubation hypotension

Is everyone ready?

Does anyone have any questions or concerns?



What You Need


SpaceLab™ Monitor

Masimo Airway Adaptor
(front and back of infant/neonatal
package shown here)Heated Moisture
Exchanger (HME)

Vent Circuit



How to Set it Up

Plug into cartridge on
back of monitor.

Troubleshooting: Things to Try if Unable to Get Reading

- Swap the cartridge on the back of the monitor with one from another room. (See photo to right.) Some monitors are not defaulted to monitor CO₂ and must be set up: (1) After plugging cartridge in, screen will show "NO SAMPLING LINE – Check system." (2) Press "GAS." (3) Press "SETUP." (4) Press "RESUME CO₂."
- Try new Masimo Airway Adaptor.
- Calibrate the monitor by pressing "cal" → "gas."
- Make sure there is no moisture in the adaptor.
- Check that all connections fit tightly.

Cartridge



This resource is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guidelines Committee 7/23/25.

If comments about this resource, please contact Leslie_Herrmann@ykhc.org.



What You Need

Zoll™ Monitor with this cable



Zoll Airway Adaptor

- Neonatal/Pediatric adaptor (shown) is purple and is for ETT sizes 4.0 or smaller.
- Pediatric/Adult adaptor is clear and is for ETT sizes larger than 4.0.

Heated Moisture
Exchanger (HME)

Vent Circuit



Plug cable into back of monitor.



Allow two minutes for
monitor to warm up.

How to Set it Up



Y-connector
to ET tube

Inline suction
catheter (Ballard)

Zoll Airway
Adaptor

HME

Vent circuit



Troubleshooting: Things to Try if Unable to Get Reading

- Make sure the Zoll has had two minutes to warm up.
- Try new Zoll Airway Adaptor.
- Make sure there is no moisture in the adaptor.
- Check that all connections fit tightly.

This resource is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guidelines Committee 7/23/25.


If comments about this resource, please contact Leslie_Herrmann@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Treatment Protocol

Massive Transfusion Protocol (≥14 years)



Criteria for Activation (any)

- Patient has required ≥3 units pRBC in 1 hour
- Shock Index of HR/sBP > 1
- pH < 7.24 or base deficit ≤ -10
- ABC Score: ≥2 of the following
 - Penetrating mechanism
 - (+) FAST exam
 - Systolic BP < 90
 - Heart rate > 120
- RABT Score: ≥2 of the following
 - Penetrating mechanism
 - (+) FAST exam
 - Shock index > 1
 - Pelvic fracture

Factors to Consider for Cancellation of the MTP

- Bleeding is controlled or significantly improved.
- Hemodynamic status is improving
 - Stable or decreasing HR
 - Stable or increasing BP
 - Stable or increasing UOP
 - Decreasing vasopressor requirement
- Labs improving:
 - Hgb > 8
 - PT < 18
 - PTT < 42
 - Platelets > 50
 - Fibrinogen > 150
 - Lactate < 3
- Recognition of medical futility of further resuscitative efforts.

Risks of Massive Transfusion

Non-fatal complications in 50% of patients transfused >5 units of blood:

- Coagulopathy
- Thrombosis
- ARDS: adult respiratory distress syndrome
- TACO: transfusion-associated circulatory overload
- TRALI: transfusion-related acute lung injury
- Hemolytic reaction

Tips

- Avoid crystalloids to prevent dilution
- Acidosis predicts mortality. Treatment is to optimize resuscitation. No clear benefit to bicarb but may be considered if pH persistently < 7.2 despite resuscitation.
- Avoid hypothermia; keep core temperature > 36°C.
- pRBC should be transfused with a mass transfuser and blood warmer.
- Utilize ED technician to transport blood products during an MTP.

Availability of Blood Products at YKHC

Blood type	Quantity (units)	Rounds**
O negative	10	2-3
O positive	16	4-5
A positive	10	6-7
B positive	6	5-6
AB	none	7-8

Anticoagulation Reversal Agents at YKHC

- DOA/warfarin: Kcentra 2000 units or 25-50 units/kg at ~3 units/kg/minute
- Warfarin: Vitamin K 2.5-10 mg IV over 10-20 minutes
- Dabigatran: PraxBind® 2.5 grams IV Q5 minutes x2
- Heparin: Protamine 0.5-1 mg IV over 10 minutes

**Per lab policy, the blood bank cannot dispense all units to a single patient in case of other emergencies. May discuss with Clinical Director on call.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 11/21/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Travis_Nelson@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Treatment Protocol

Medevac Activation: Village to YKHC

Indications for medevac:

- Patient is in danger of losing:
 - Life
 - Limb
 - Eyesight
- Preterm labor

NOTE: In the event of multiple medevacs, the ED physician in collaboration with medevac dispatch prioritizes the medevacs.

Occasionally, a charter may be able to fly when a medevac cannot. Consider this option if on weather-hold.

May also consult [Military Transport for Emergencies](#) guideline.

Make every effort for the patient to be transferred to the village clinic to meet LifeMed. This is for the safety of the LifeMed crew. If there is no health aide in the village, ask for an Office Assistant to open the clinic for the patient to wait there.

Preterm Labor

1. See the [Labor Patient in a Village](#) and [Village Deliveries](#) guidelines.
2. Notify pediatrician. Take "go bag" from L&D with surfactant.
3. Remember to notify ED physician and OB charge RN.
4. If appropriate, consider contacting facilities in Anchorage to discuss suitability of ramp transfer.
5. Hospitalist remaining at YK will cover all emergency RMT for adults and peds, AND continue managing the preterm labor patient. Ask for help if needed (E1/E2, experienced clinic providers, CD on call, etc.).

In the event that a medevac is cancelled (patient deemed stable to come in on scheduled flight) medevac dispatch and receiving department must be notified by the managing physician immediately.

Health Aide or Provider in village consults Wards Hospitalist/Emergency RMT for initial management and possible medevac of critically ill patient.

Hospitalist consults ED Doctor on Duty to confirm appropriateness.

Notify pediatric hospitalist when activating a medevac for any child <12 years old.

If patient is NOT a beneficiary, ask if they have a preferred medevac company. If not, suggest they [register for LifeMed insurance online](#).

Activation of Medevac

Activating provider calls medevac dispatch with patient's name, DOB, village, and diagnosis. If applicable, dispatch will ask for escort's name and weight.

LifeMed Dispatch 1-800-478-5433

Complete the Patient Transport Order (PTO) and ensure it is faxed to **5-543-1262** and **x6099**.

Village Management:

Explicitly clarify whether ED Physician or Hospitalist will continue managing the patient with the health aide. (Typically this will be the ED physician)

Managing physician calls village Health Aide for updates, continues active management of the patient, and documents in EMR.

Managing physician updates ED physician & charge RN.

Blood Products

If appropriate, consider sending LifeMed crew with blood products. If this is anticipated:

1. Notify dispatch of plan, confirm whether LifeMed has available blood at hangar.
2. If no blood at hangar, contact YK bloodbank to request 2 units of "emergency release" blood to be prepared immediately.

Dispatch Process

1. Selected medevac dispatch notifies their medevac team.
If medevac cannot launch (weather, runway lights, etc.) dispatch will notify managing physician. Pilot will continue to check weather.
2. Receiving unit clerk faxes PTO and face sheet to medevac crew.
3. Medevac crew contacts health aide and managing physician as needed.
4. If there is a prolonged delay, medevac crew will contact the managing physician and health aide.

Medevac launches

1. Once in village, medevac crew calls managing physician to give report, establish treatment plan, and give ETA in Bethel.
2. Managing physician keeps receiving charge nurse informed of patient status/ETA of medevac.

Arrival in Bethel

Patient care is transferred to receiving unit and medevac crew gives report to staff.

Consider Medevac Direct to Anchorage

Indications:

1. Obvious need for acute surgical intervention (e.g. hip fracture)
 2. STEMI, obvious acute CVA
 3. Intubated in field
- MUST also be hemodynamically stable (not require stabilization at YK before transfer)

Notify LifeMed Dispatch immediately if considering.

Discuss with receiving facility specialist (e.g. orthopedics, ICU) and ER if needed.

Consult LifeMed regarding logistics of ramp transfer.


This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 12/12/24.
If comments about this guideline, please contact clinical_guidelines@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Treatment Protocol

Medevac Activation: YKHC to Anchorage



Phone Numbers

- LifeMed Dispatch: *96 or (800) 478-5433
- Guardian Dispatch: 888-997-3822
- Alaska Native Medical Center: Main operator: *97 or (907) 563-2662
Transfer Center: *58 or (907) 729-2337 or Tiger Text ANMC Transfer Center Coordinator
ED: (907) 729-1729
- Providence Alaska Medical Center: Main operator: (907) 562-2211
Transfer Center: (907) 212-7363
Trauma on call: (907) 212-2525
ED: (907) 212-3111
- Alaska Regional Hospital: Main operator: (907) 276-1131
Transfer Center: (844) 880-5522
- Fairbanks Memorial Hospital: Main operator: (907) 452-8181, House supervisor pager: (800) 607-3974
- Mat-Su Regional Medical Center: Main operator: (907) 861-6000
Transfer Center: (907) 861-6440
- Joint Base Elmendorf Richardson Hospital:
ED: (907) 580-5556
House supervisor: (907) 580-6413
- Department of Corrections On Call: (844) 751-4588

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 12/12/24.
If comments about this guideline, please contact clinical_guidelines@ykhc.org.



Trauma Transfer Criteria (TTC)

1. **CNS:**
 - Open skull fracture
 - ICH with mass effect or increase in size
 - GCS <10 or decreasing
 - Lateralizing or focal neurologic signs
 - Spinal cord injury
 - Vascular injury
2. **Chest:**
 - Requirement for ventilator
 - Hemothorax or pneumothorax not resolved by chest tube
 - Great vessel injury
 - Flail chest
 - Cardiac or pericardial injury
3. **Pelvis:**
 - Open pelvic fracture
 - Unstable pelvic ring fracture
4. **Abdomen:**
 - Solid organ injury
 - Viscous perforation
 - Hemo- or pneumoperitoneum
 - Major vessel injury
5. **Extremity:**
 - Fracture or dislocation with neurovascular compromise/loss of perfusion
 - Suspected compartment syndrome
 - Dislocation unable to be reduced
 - Threatened limb (ischemia, crush, shock, etc.)
 - Open long bone fracture
 - Femur fracture
6. **Shock persisting despite resuscitation**
7. **Multisystem trauma**
8. **Burns that exceed admission criteria at YKHC** (see [guideline](#))
9. **Frostbite that meets criteria for tPA** (see [guideline](#))
10. **Hanging/strangulation with CT findings and/or airway compromise** (see [guideline](#))

YKHC Trauma Admission Criteria

The following diagnoses meet the threshold to be admitted at YKHC.

1. Non-operative hip fracture
2. Chest tube placement not requiring surgery
3. >3 rib fractures
4. Non-accidental trauma of child or vulnerable adult
5. Pain not controlled by oral medications
6. Need for serial exams
7. Need for inpatient wound care
8. Hanging/strangulation with no CT findings or airway compromise (see [guideline](#))
9. Trauma patients with comorbidities, especially children, elders, and pregnant people, should prompt a lower threshold for admission.

Phone Numbers

- LifeMed Dispatch: *96 or (800) 478-5433
- Alaska Native Medical Center: Main operator: *97 or (907) 563-2662
Transfer Center: (907) 729-2337 or Tiger Text ANMC Transfer Center Coordinator
ED: (907) 729-1729
- Providence Alaska Medical Center: Main operator: (907) 562-2211
Transfer Center: (907) 212-7363
ED: (907) 212-3111
- Alaska Regional Hospital: Main operator: (907) 276-1131
Transfer Center: (844) 880-5522
- Fairbanks Memorial Hospital: Main operator: (907) 452-8181
House supervisor pager: (800) 607-3974
- Mat-Su Regional Medical Center: Main operator: (907) 861-6000
Transfer Center: (907) 861-6440
- Joint Base Elmendorf Richardson Hospital:
ED: (907) 580-5556
House supervisor: (907) 580-6413
- Department of Corrections On Call: (844) 751-4588

Patient with trauma is seen in a YKHC facility.

In the pre-hospital setting, if patient meets TTC AND is stable, consider ramp transfer. Otherwise, transfer to YKHC ED via most appropriate method.

Patient with trauma arrives in ED.

Follow ATLS protocols.

Stabilize patient and perform immediate life-saving interventions. If TTC are met, initiate transfer process as soon as possible. May delegate to another provider.

See [Medevac: Bethel to Anchorage Guideline](#) for transfer process.

- If patient meets TTC, contact Transfer Center of expected accepting institution. The Transfer Center will connect the YKHC provider with the trauma surgeon, who must be the first point of contact. The Transfer Center will contact other specialists as needed.

- If patient does not meet TTC but YKHC provider has questions about management or disposition, contact trauma surgeon via Tiger Connect or operator.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 9/24/24.


Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Megan_Young@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline Military Transport for Emergencies



Things to Consider

The local LifeMed team can sometimes go on a military flight. This decision is up to the local team and their administration and depends on many factors.

If the transport team is all military:

- Will military transport inappropriately lower the level of care the patient is receiving?
- What are the capabilities of the military team? Are they pararescue jumpers (PJs), paramedics, EMTs, etc.?
- What kind of equipment will the military team have?
- Does the military team have pediatric experience and equipment, if applicable?

If you are sending a team from YK:

- Will sending a team of YK employees impact the normal operations of the hospital? (You should avoid sending anyone scheduled to work the current or next shift.)
- An ideal YK team includes an ER RN and/or paramedic. Transport/EMS experience is a must.
- **A YK team must be entirely voluntary.**
- Ensure the team will have all appropriate drugs, weight-based equipment, monitors, pumps, stretchers/backboards, etc.
- Make a plan to keep the patient warm – the military will usually not supply blankets, Doctor Downs, etc.
- **If military transport is used, no YK trainees (residents, students, visitors, etc.) or other “ride-alongs” are allowed to go.** Ride-alongs may only go on LifeMed transports with the local team on their fixed wing aircraft.

Things to Know

- The RCC coordinates military missions. They will connect you with the appropriate people from the branch responding, which may be the National Guard, the Coast Guard, or the Air Force.
- You may have to retell the story to several people, including people with minimal medical knowledge. It helps to involve another provider to help coordinate the many phone calls without negatively impacting patient care.
- **The process often takes 6-8 hours or more.** If the Blackhawk and a full crew are not physically in Bethel, the military may have to send aircrafts from elsewhere in Alaska, which can lengthen the process to 10-12 hours.

Definitions

LLE: life, limb, or eyesight in danger
 CD: clinical director
 RCC: Rescue Coordination Center
 PJ: pararescue jumpers. These are military medics with ACLS and ATLS training who are not trained to provide further critical care. (For example, neonatal care, ventilator management, and infusion of medications are not typically part of their scope of practice.)

The form to activate the RCC can be found [here](#).

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 1/19/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Ellen_Hodges@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline

Pediatric Medevacs: Bethel to Anchorage

Pediatric patient in Bethel
requires medevac.

Critical pediatric patient (including patients
in respiratory distress, on HFNC, with
suspected sepsis, in status epilepticus, etc.)

- Consult YKHC pediatric hospitalist as soon as possible.
- YKHC pediatric hospitalist (or other provider in consultation) can call ANMC PICU directly for advice, management, and accepting physician.

YKHC provider remains in control of
management until patient leaves facility.

- Trauma, surgical, or orthopedic emergency
- Noncritical: for bed space, diagnostic or specialty services not available at YKHC, etc.

- Follow usual workflow.
- Pediatric consult optional.
- Please notify the YKHC pediatric hospitalist for any Chronic Peds Patient (CPP) being transferred.

**Contact YKHC pediatric hospitalist
via Tiger Connect “Peds Wards On
Duty” for all potentially critical
pediatric patients.**

Use the [Pediatric Critical Care Guide](#) and
ED Peds Critical Care PowerPlan.

Surgical Patients

- For all pediatric trauma cases and surgical cases in children >4 years, call the ANMC general surgeon on call.
- For surgical cases in children ≤4 years, call the pediatric surgeon on call through the ANMC operator or their office at (907) 929-7337.

Non-beneficiary Patients

- Non-beneficiary patients are transferred to Providence Alaska Medical Center via the PAMC Transfer Center. If you are told there is no bed, ask to speak to the physician (hospitalist or PICU). Arrangements can often be made to accept a patient even if a bed is not immediately available.
- Ask about medevac insurance coverage. May suggest family [register for LifeMed insurance](#) online, which can be done just prior to activation.

Neonatal Transfers

Contact PAMC neonatologist at (907) 212-3614 for advice, management recommendations, etc.

Notify ANMC pediatric hospitalist on-call for any beneficiary infant transferred to PAMC NICU.

After obtaining accepting physician, YKHC physician is responsible for activating Lifemed and discussing patient with neonatologist, if needed.

When to Transfer to PAMC NICU:

- GA <32 weeks
- BW <1500 grams
- Any newborn who required intubation
- Newborns requiring prompt surgical or medical subspecialty care
- No beds available at ANMC or non-beneficiary infant requiring transfer
- Discretion of NNP

When to Transfer to ANMC NICU:

- GA ≥32 weeks
- BW ≥1500 grams
- Any baby who meets criteria for transfer per the [Late Preterm guideline](#)
- Term or early term babies with temperature instability, respiratory distress, supplemental O₂ requirement, hypoglycemia requiring IV treatment, need for IV antibiotics, etc.

Contact

- ANMC PICU (physician or NP): (907) 297-8809 – may request to speak with physician.
- ANMC Transfer Center: (907) 729-2337 or Tiger Connect Transfer Center Coordinator.
- LifeMed: *96 or (800) 478-5433
- PAMC Transfer Center: (907) 212-7363
- PAMC PICU: 212-3133
- PAMC NICU: (907) 212-3614
- Alaska Pediatric Surgery: (907) 929-7337

LifeMed is the preferred medevac company for children younger than 3 years old.
If any difficulty, call CD on call to discuss.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 6/1/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



Indications for Procedural Sedation

Any procedure that cannot be accomplished with patient's current level of cooperation or pain tolerance.

Examples:

- Nonemergent chest tube placement
- Cardioversion
- I&D
- Laceration repair
- Fracture or joint dislocation reduction
- Pediatric foreign body removal
- Imaging

Airway Risk Assessment

See [Intubation guideline](#) for resources.

High-Risk History

- Stridor
- Obstructive sleep apnea
- Hx Trisomy 21
- Dysmorphic facial features
- Active respiratory tract infection
- Hx of difficult intubation
- Hx of cervical spine pathology

Exam

- Check that patient can open mouth fully and that TMJ function is normal.
- Look for micrognathia, loose teeth, dental appliance, and craniofacial abnormalities.
- Check that patient is able to extend neck >70°.
- Determine Mallampati Score and check 3-3-2 rule (in adults).

Expected Sedation Risk Level

Airway Risk Assessment combined with expected depth of sedation should guide level of rescue preparation.

- 1. No risk factors present:** No high risk findings in airway assessment and exam, ASA I-II.
Plan: standard monitoring and equipment, including:
 - Cardiopulmonary monitor (three lead ECG & RR)
 - Pulse-oximetry
 - Supplemental oxygen should be prepared but not given unless otherwise indicated.
 - BVM in room.
 - Suction.
 - End-tidal CO₂ monitor
 - Reversal agents in room.

- 2. Risk factors present:** some concern for airway status based on airway assessment and exam, but patient not expected to decompensate, and benefits of sedation outweigh risks.

Plan: discuss risks with patient/caregiver.

- In addition to standard monitoring and personnel, the following must also be present:
- A healthcare provider dedicated to airway management (preferably an RT)
 - Oral airway – correct size open and at bedside
 - Nasal trumpet – correct size open and at bedside
 - BVM with appropriately-sized mask should be open and prepared at bedside

Note: Consider CRNA at bedside.

Does provider have privileges for procedural sedation?

—No—
Seek out colleague with privileges
or
Page CRNA on call.

Yes

Patient has a condition that meets indications.

There are no contraindications.

Perform Airway Risk Assessment. See box.
Determine Expected Sedation Risk Level and plan accordingly. See box.

Discuss risks and benefits of procedural sedation with patient/guardian.
Obtain written consent.

Ensure all equipment is present
for Expected Sedation Risk Level.
(See box.)

Choose appropriate sedation medication. See page 2.
Calculate appropriate dose and confirm with team.

Perform time out:
• Correct patient
• Correct procedure
• Correct site

Administer medication.

RN documents medication
administration time and effect and
patient assessments on flowsheet.

Perform procedure.

- Monitor patient through recovery.
- RN remains at bedside until patient is fully alert.
- Provider documents in note using autotext "...procedsedationoutsideOR."

Contraindications

- Patient or caregiver refusal
- Hemodynamic instability
- Respiratory distress

Note: if benefits of procedure outweigh risk despite contraindications, refer to higher level of care or consult a local airway expert and discuss risks with patient/caregiver.

Note: Per ACEP, NPO times should not affect decision-making about moderate sedation. If deep sedation is required, non-emergent procedures may need to be delayed. Consider consult to CRNA to discuss further.

Levels of Sedation

- **Moderate sedation:** patient responds to light tactile stimulation. Spontaneous ventilation is adequate. An IV is required only if IV medication is to be given. (For example, if ketamine is to be given IM with no IV agents planned, an IV is not necessary.)
- **Deep sedation:** noxious stimuli are required to elicit a response. Spontaneous ventilation and ability to maintain airway may be impaired. Not routinely performed in the ED outside of life-threatening situations.

Other Scenarios

Use of a single agent is not always considered sedation. This protocol does not apply to the following:

- Anxiolysis with a benzodiazepine: patient may be drowsy but responds appropriately to verbal commands. Example: midazolam 0.2 mg/kg IN up to max dose 6 mg in <5 years and 10 mg in >5 years.
- Analgesia with opioids: pain control with intact decision-making.
- Ketamine at analgesic dosing (see next page).



Agent	Bolus Dose	Titration Dose	Onset	Duration	Reversal Agent	Comments
Etomidate	<u>Patients >10 years:</u> 0.2 mg/kg	0.05 mg/kg Q3-5 min	30-60 seconds	3-5 minutes	Time	<ul style="list-style-type: none"> No analgesic effect. Use IBW if BMI>30. Consider lower dose (0.1 mg/kg) for age >60 years, concurrent opioids, or if recent alcohol use. Administer via larger vessel. (antecubital or larger). Precautions: 30% have myoclonus with transient skeletal/eye movements.
	<u>Patients ≤10 years:</u> 0.2 mg/kg (0.1-0.3 mg/kg) Slow IV push over 30-60 seconds.	0.05 mg/kg Q3-5 min	30 seconds	2-10 minutes		
Ketamine for sedation	<u>Adults:</u> 1-2 mg/kg IV over 1-2 min		30 seconds	10-20 min	<ul style="list-style-type: none"> Time For laryngospasm: Succinylcholine 0.25-0.5 mg/kg IV or 3-4 mg/kg IM 	<ul style="list-style-type: none"> Local anesthetic (eg. lidocaine) can increase effective duration. Consider lower dose range for >60 years, concurrent opioids/alcohol. Consider dosing by adjusted body weight if BMI>30. Precautions: emergence reactions (treat with benzodiazepines), nausea/vomiting (pre-treat with ondansetron), transient increase in salivation. Contraindications: pregnancy, age <3 months.
	4-5 mg/kg IM		3-4 min	20-30 min		
	<u>Children >3 mo:</u> 1-2 mg/kg IV over 1 min		30-120 seconds	20-60 min		
	4-5 mg/kg IM		5-10 min	30-90 min		
Ketamine for analgesia	5 mg/kg PO		20-45 min	60-120 min	<ul style="list-style-type: none"> Time For laryngospasm: Succinylcholine 0.25-0.5 mg/kg IV or 3-4 mg/kg IM 	<ul style="list-style-type: none"> Local anesthetic (eg. lidocaine) can increase effective duration. Consider lower dose range for >60 years, concurrent opioids/alcohol. Consider dosing by adjusted body weight if BMI>30. Precautions: emergence reactions (treat with benzodiazepines), nausea/vomiting (pre-treat with ondansetron), transient increase in salivation. Contraindications: pregnancy, age <3 months.
	0.1-0.4 mg/kg IV 0.4-1.0 mg/kg IM		30 seconds 3-4 min	10-20 min 20-30 min		
Propofol	<u>Patients >2 yrs:</u> IV load 0.5-1 mg/kg	Repeat 0.1-0.3 mg/kg Q30-60 seconds	30-60 seconds	3-10 min	Time	<ul style="list-style-type: none"> No analgesia. Consider low dose for age >60, concurrent opioids/alcohol. Consider dosing by adjusted body weight if BMI>30. Separate administration of opioid and propofol by >20 minutes to decrease respiratory depression. Pre-oxygenate with high flow supplemental oxygen at least 3 minutes prior to procedure. Precautions: burning sensation during administration, hypotension, ↓CO, or bradyarrhythmias. High risk of respiratory depression/failure. Contraindications: allergies to egg, soybean, fat emulsion.
	<u>Children 6 mos – 2 yrs:</u> IV load 1-2 mg/kg	Repeat 0.1-0.3 mg/kg Q30-60 seconds				
		Max cumulative dose 3 mg/kg				
Morphine	<u>Adults:</u> 1-4 mg IV		5-10 min IV	3-5 hours	Naloxone 0.1 mg/kg IV. May repeat Q2 minutes.	<ul style="list-style-type: none"> Reduce dose when combining with a benzodiazepine. As opioids provide sedation and analgesia, administer them prior to benzodiazepines.
	10 mg PO		30 min PO			
	<u>Pediatrics:</u> 0.05-0.1 mg/kg IV Max 4 mg		5-10 min	2-3 hours		
Fentanyl	<u>Adults:</u> 0.5 mcg/kg if given with other sedatives	May repeat dose Q2min until desired sedation and analgesia achieved	<1 min		Naloxone 0.1 mg/kg IV. May repeat Q2 minutes.	<ul style="list-style-type: none"> Reduce dose when combining with a benzodiazepine. As opioids provide sedation and analgesia, administer them prior to benzodiazepines.
	0.5-1 mcg/kg Max 100 mg					
	<u>Pediatrics:</u> 1 mcg/kg IV up to 50 mcg/dose		3-5 min	30-60 min		



Agent	Bolus Dose	Titration Dose	Onset	Duration	Reversal Agent	Comments
Midazolam	<p><u>Adults:</u> 2-5 mg IV</p> <p><u>Pediatrics (6 mos - 12 yrs):</u> 0.2-0.3 mg/kg/dose IN</p> <p>0.05 mg/kg IV</p>	<p>May repeat dose Q2min until adequate sedation. Max 0.3 mg/kg.</p> <p>May repeat dose Q5min until max dose of 0.5 mg/kg is reached. Age <5 max 6 mg; age >5 max 10 mg.</p>	3-5 min	15-20 min	<p>Flumazenil 0.01 mg/kg (up to 0.2 mg) IV over 15 seconds. May repeat Q1 minute.</p>	<ul style="list-style-type: none"> • No analgesia. • Consider lower dose range for >60 years, concurrent opioids/alcohol. • Watch for dose-related hypotension.
Dexmedetomidine (Precedex™)	<p><u>Adults:</u> Bolus 0.5-1 mcg/kg – infuse over 10 minutes.</p> <p><u>2-18 years:</u> Bolus 2 mcg/kg -infuse over 10 minutes.</p> <p><u>1 month to < 2 years:</u> Bolus 1.5 mcg/kg – infuse over 10 minutes.</p> <p><u>Intranasal for <10 years</u> 2 mcg/kg IN x1, max dose 200 mcg.</p>	<p>Infusion 0.2-1 mcg/kg/hour.</p> <p>Infusion 1.5 mcg/kg/hour (titrate up to 2 mcg/kg/hour).</p> <p>Infusion 1.5 mcg/kg/hour (may titrate up to 2 mcg/kg/hour).</p>	Onset 5-10 minutes.	<p>Duration 60-240 minutes post discontinuation of infusion</p> <p>Duration 30-70 minutes post discontinuation of infusion</p> <p>Duration 30-70 minutes post discontinuation of infusion.</p>		<ul style="list-style-type: none"> • Sedative with modest analgesia and minimal respiratory depression. • No amnestic properties – consider midazolam if amnesia desired. • Biggest side effects: bradycardia and hypotension – generally dose/rate dependent. • Relative contraindications: inadequate hydration, reduced cardiac output, elevated LFTs. • Absolute contraindications: digoxin, cardiac conduction abnormalities.



PROCEDURE MONITORING

HR, RR, SpO₂, LOC (level of consciousness), and Modified Aldrete Score to be monitored and recorded Q5 minutes until fifteen minutes after last administration of sedating medication, then Q15 minutes x1 hour, then Q1h until returned to pre-sedation baseline. Respiratory status should be monitored continuously.

TIME OUT PERFORMED

- ☐ Correct patient
- ☐ Correct procedure
- ☐ Correct site

Time _____ Initials _____

EQUIPMENT READINESS

- | | |
|---|--|
| In room: | Readily accessible: |
| <input type="checkbox"/> Cardiopulmonary monitor with three lead ECG, RR, and BP cuff | <input type="checkbox"/> Crash cart |
| <input type="checkbox"/> Pulse-oximeter | <input type="checkbox"/> Reversal agents |
| <input type="checkbox"/> Supplemental oxygen | |
| <input type="checkbox"/> BVM | |
| <input type="checkbox"/> Suction | |
| <input type="checkbox"/> End-tidal CO ₂ monitoring | |

PRESENT IN ROOM (NAME AND ROLE)

PRE-SEDATION IV ACCESS

IVF _____ Site _____

Gauge _____ Rate _____

POST-SEDATION EVALUATION

- ☐ VS and SpO₂ stable and patient has returned to pre-sedation baseline.
- ☐ LOC at pre-sedation baseline.
- ☐ Airway protective reflexes intact or at pre-sedation baseline.
- ☐ Patient tolerates oral intake.
- ☐ Ambulation at baseline.

OUTCOMES AND MONITORING

Check all that apply:

- ☐ Apnea > 15 seconds.
- ☐ Intubation or positive pressure ventilation.
- ☐ Desaturation with SpO₂ <90% for >90 seconds.
- ☐ Vomiting.
- ☐ HR, CP, or RR change 30% from baseline.
- ☐ Emergency consultation with CRNA after start of procedure.
- ☐ No complications.

PROCEDURE SUMMARY

Date of procedure: _____
Procedure start time: _____
Procedure end time: _____
Time last sedating medication was given: _____
Deepest level of sedation achieved: _____
IVF received (type and total volume): _____

RESPIRATORY EFFORT QUALITY

N = normal L = labored
S = shallow R = regular
D = deep I = irregular

LOC SCALE

5 = awake and alert
4 = sleeping intermittently
3 = asleep but responds to voice
2 = responds to painful stimuli
1 = unresponsive

MODIFIED ALDRETE SCORE

Activity

Able to move four extremities voluntarily on command.	2
Able to move two extremities voluntarily on command.	1
Unable to move.	0

Respiration

Able to breathe deeply and cough freely.	2
Dyspnea or limited breathing.	1
Apnea.	0

Circulation

BP and HR ± 20% of pre-sedation level.	2
BP and HR ± 20-50% of pre-sedation level.	1
BP and HR ± 50% of pre-sedation level.	0

Consciousness

Fully awake and able to answer questions.	2
Arousable only to calling.	1
Unresponsive.	0

Oxygenation

SpO ₂ >90% on room air.	2
Requires supplemental oxygen to maintain SpO ₂ >90%.	1
SpO ₂ <90% despite supplemental oxygen.	0

SIGNATURES

Provider performing sedation: _____

Monitoring RN: _____

Provider performing procedure: _____

Place patient ID sticker here.

[illegible]

SIGNATURES

Monitoring RN(s): _____

Place patient ID sticker here.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline

Respiratory Distress & Bronchiolitis Management (<5 years)

Criteria for Levels of Care

Outpatient or Village

- WOB is mild or absent.
- No hypoxemia.
- Able to maintain hydration without IVF.
- Tolerating home therapy with reliable caregivers.
- No apnea.

Hypoxemia

- <90% while awake.
- <88% while asleep.
- Sustained for >10 minutes.

Inpatient at YKHC

- Requires supplemental oxygen to prevent hypoxemia or improve WOB. If requiring >2 L NC, reevaluate whether patient is appropriate to stay at YKHC.
- Requires IV or NG fluids.
- Question of apnea.
- Not tolerating home therapy or unreliable follow-up.
- Does not meet criteria for transfer to higher level of care.

Transfer to Higher Level of Care

- Requires >2-3 L oxygen for hypoxemia or to improve WOB.
- Witnessed apnea.
- Requires neb treatments more frequently than Q2-3h for >8 hours.
- Sustained tachycardia, tachypnea, or respiratory distress despite treatment.
- Significant pleural effusion.

SUPPORTIVE MEASURES

- Control fever, as it can be an independent cause of respiratory distress and tachycardia.
- Nasal suction with nasal bulb syringe and olive tip plus saline.
- Hydration by IV or enteral (including NG and G-tube).
- Gentle P&PD/CPT if helpful.
- Saline neb (either 0.9% or hypertonic 3%).
- Consider albuterol trial even if no wheezing heard, especially in Alaska Native patients as they have high rates of RAD.

Village Management

- Institute **SUPPORTIVE MEASURES**.
- Have low threshold to refer to Bethel or consult a pediatrician, especially if no improvement after 2-3 days.
- If Hx recurrent wheezing with viral illnesses, start budesonide 1 mg by nebulizer twice daily for 7 days.

Risk Factors for Apnea


- RSV
- Post-conceptual age <48 weeks
- Low birth weight
- Tachypnea or bradypnea
- Decreased oxygen saturation on room air

NOTE:

- If <3 months or history of prematurity, keep patient in Bethel and have low threshold for admission.
- If patient is <90 days and febrile, please see [fever guideline](#).

Pulse-Oximetry Monitoring

- Pulse-ox may be ordered Q4h (not continuously) if patient >6 months and stable.
- Being on oxygen does not mandate continuous pulse-oximetry if patient is stable.



Escalation of Care


Consult pediatrician.

- Nasal steroids (Pred-Forte (drops-ophth) 1 drop each nostril BID x5 days) and/or Neo-Syneprhine (1 spray each nostril BID x3 days).
- More frequent albuterol/hypertonic saline nebs. (May give Q2h nebs x3 on the floor, but if patient needs sustained Q2h treatment, initiate transfer to higher level of care.)
- Racemic epinephrine neb. (Use with caution if HR > 200.)

Steroids

- Recent national guidelines recommend that children ≤4 with recurrent wheezing with viral illnesses should be given a 7-10 day course of inhaled steroids like budesonide or fluticasone.
- National guidelines recommend against systemic steroids as the potential harm is generally greater than the potential benefit.
- If considering starting systemic steroids, please consult a pediatrician.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 3/11/24. Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact Amy_Carson-Strnad@ykhc.org.



Village Management

- Always check blood glucose.
- Patients <18 months with first-time febrile seizure or any patient with first-time non-febrile seizure should come to Bethel for evaluation.
- Patients ≥18 months meeting all low-risk febrile seizure criteria with identified low-risk source of infection (eg AOM) may stay in village until outpatient appointment available. Ensure close village follow-up.
- Strongly consider medevac if child is not back to baseline, has signs of meningitis, had an atypical seizure, or presented in status epilepticus.

Box 1: Detailed History

- When/where did it occur? Awake or asleep?
- What preceded the event (eg head trauma, crying, etc.)?
- How long did it last?
- Ask caregiver to recount, step-by-step, what happened.
- Type of movement and what part of body? Symmetric?
- Interventions?
- Incontinence?
- Behavior after event? How long till back to baseline?

HPI

- Intercurrent illness/fevers
- Medications
- Recent intake, including free water and diluted formula
- Ingestions
- Trauma – consider child abuse.

PMH

- Prior history of seizures
- History of breathholding

Family Hx: Seizures, febrile seizures, breathholding, etc.

Box 2: Low risk febrile seizure criteria

1. 6 months to 5 years of age.
2. Fever present.
3. Seizure generalized (nonfocal).
4. Seizure duration <15 minutes.
5. Child has normal neurologic examination.
6. Child has no history of previous neurologic or CNS abnormality.
7. Only one seizure in a 24 hour period.
8. Child has returned to baseline.
9. No meningial signs:
 - Irritability or inconsolability
 - Nuchal rigidity
 - Bulging fontanelle
 - Lethargy or somnolence
 - Focal neurologic findings
10. Child has NOT received antibiotics in the past 72 hours.

Box 3: Work-up

- Bedside glucose
- EKG for first event
- CBC, BMP, magnesium, phosphate
- Urine drug screen
- Perform LP if persistent altered mental status, meningitis suspected, or <18 months of age and delayed return to baseline.

Radiological studies:

Obtain head CT without contrast prior to LP if concerning neurologic status, persistently altered mental status, history of trauma, focal neurologic findings, or focal seizure.

Consider using the [Bacterial Meningitis Score for Children](#) to help rule-out meningitis.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 2/9/24. Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Amy_Carson-Strnad@ykhc.org.



Sepsis:
Suspected infection plus systemic inflammatory response.
Can use SIRS or qSOFA.
General signs:

- Temp > 100.4° or < 96.8° F
- HR > 100
- RR > 22
- Systolic BP < 100
- WBC > 12,000 or < 4,000

Severe Sepsis:
Sepsis plus evidence of end-organ damage. Can include:

- Hypotension (SBP < 90, MAP < 65, baseline drop in SBP > 40)
- Cool extremities, delayed cap refill
- Altered mental status (GCS < 15)
- Poor urine output
- New need for respiratory support (high flow oxygen, NIPPV)
- Lab indicators can include:
Lactate > 2
INR > 1.5, platelets < 100,000
Creat > 0.5 over baseline value
Bilirubin > 4

Septic Shock:
Severe sepsis persisting/worsening despite initial resuscitative measures.

In Bethel:

- Start pressors (see [medications](#)).
- Move toward central line placement, but ok to start first pressor peripherally.
- Consult ICU and move toward transfer.

In Village/SRC:

- Activate medevac if not done already.
- Consult ED physician for further management, including ongoing fluids, antibiotics, and pressors if available in SRC.


Intubation in Sepsis

- Higher risk for periintubation arrest due to hypotension, acidosis, etc.
- Strive for fluid resuscitation and/or pressors before intubation.
- Consider lower dose of induction agent (consult pharmacy or ICU).
- Vent settings: TV 6 mL/kg IBW, plateau pressures < 30.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 3/13/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact clinical_guidelines@ykhc.org.



IV Fluids in Sepsis

Historical consensus was every septic patient needed 30 mL/kg IVF as quickly as possible. There is not good evidence that this improves mortality. Likewise, fluid resuscitation guided by lactate alone is not associated with improved mortality. There is evidence of harm in over-fluid resuscitating patients, and in delay to initiating pressors if appropriate.

General Fluid Management Recommendations

- If hypovolemic, give fluids.
- If euvolemic, don't give excessive fluids.
- If progressive respiratory distress and pulmonary edema, stop fluids.
- Give smaller boluses 500-1000 mL and assess response.
- If CHF/renal failure/volume overload, fluids are not wrong but low threshold to consult ICU for assistance.

Medications Outside Bethel

Village formulary:

- Ceftriaxone 1-2 grams IM (for most cases)
- Metronidazole 500 mg PO (abdominal source, necrotizing SSTI, other need for anaerobic coverage)
- Azithromycin 500 mg PO (CAP)
- Clindamycin 900 mg PO (for anaerobic coverage, toxins in necrotizing infections)

SRC formulary:

- Ceftriaxone 1-2g IV/IM (for most cases)
- Levofloxacin 750mg IV (for pseudomonas coverage)
- Clindamycin 900 mg IV (for anaerobic coverage, toxins in necrotizing infections)
- Vancomycin 25 mg/kg or 2.5 g max IV (for MRSA)
- Pressors: epinephrine – consult pharmacist if considering.



Empiric Antibiotic Recommendations by Source of Infection

If possible, first dose of antibiotics should be administered as a 30 minute infusion to reduce time to therapeutic concentration.

Unknown Source

Vancomycin¹ Load 25 mg/kg using IBW.
Max dose 2500 mg.
OR
Linezolid² 600 mg IV Q12h.

AND

Cefepime 2 grams IV Q8h.

Community-Acquired Pneumonia

Ceftriaxone 1-2 grams IV Q24h.
OR
Ampicillin-sulbactam 3 grams IV Q6h.

AND

Azithromycin 500 mg IV Q24h.
OR
Doxycycline 100 mg IV Q12h.

If at risk for
aspiration,
consider
adding:

Metronidazole 500 mg IV Q8h
if not on Unasyn.

Hospital-Acquired Pneumonia or High Risk for Multi-Drug Resistant (MDR) Organisms

Vancomycin¹ Load 25 mg/kg using IBW.
Max dose 2500 mg.
OR
Linezolid² 600 mg IV Q12h.

AND

Cefepime 2 grams IV Q8h.

Meningitis

Dexamethasone 10 mg IV
prior to antibiotics.

AND

Vancomycin¹ Load 25 mg/kg using IBW.
Max dose 2500mg.

AND

Ceftriaxone
2 grams IV Q12h.

If >50
years,
ADD

Ampicillin
2 grams IV Q6h.

Urinary Tract Infection

Ceftriaxone
1-2 grams IV Q24h.

If urological interventions or MDR risk
factors, consider adding:
Vancomycin¹ Load 25 mg/kg using IBW.
Max dose 2500 mg.
OR
Cefepime 2 gram IV Q8h.

If at risk of ESBL, ADD:
Meropenem³
500 mg IV Q8h.

Intra-abdominal or Pelvic Infection

Piperacillin-tazobactam
4.5 grams IV Q6h.

OR

Cefepime 2 grams IV Q8h.
AND
Metronidazole 500 mg IV Q6h.

OR

Ciprofloxacin 400 mg IV Q12h.
AND
Metronidazole 500 mg IV Q8h.

Skin and Soft Tissue or Necrotizing Infections

IF PURULENT:
Vancomycin¹ Load 25 mg/kg using IBW.
Max dose 2500 mg.
OR
Linezolid² 600 mg IV Q12h.

IF NON-PURULENT:
Cefazolin 2 grams IV Q8h.
OR
Ceftriaxone 1-2 grams IV Q24h.
OR
Ampicillin-sulbactam 3 grams IV Q6h.

IF NECROTIZING:
Vancomycin¹ Load 25 mg/kg using IBW.
Max dose 2500mg.
AND
Piperacillin-tazobactam 4.5 grams IV Q6h.
AND
Clindamycin 900 mg IV Q8h.

Neutropenic Cancer Patients (ANC <500)

Cefepime 2 grams IV Q8h.
OR
Piperacillin-tazobactam
4.5 grams IV Q6-8h.

AND

Vancomycin¹ Load 25 mg/kg using IBW.
Max dose 2500 mg.
OR
Linezolid² 600 mg IV Q12h.

If concerned
for HSV or
VZV,
consider
adding:

Acyclovir
10 mg/kg Q8h.
Consult pharmacy for max dosing.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/13/23.

If comments about this guideline, please contact clinical_guidelines@ykhc.org

¹ Consult pharmacy for subsequent dose/schedule.

² Linezolid may be substituted for vancomycin in patients with relative contraindication to vancomycin for high risk for acute kidney injury. Pharmacy consult required.

³ Pharmacy consult required.



Vasopressors


Central venous access is preferred for administration of vasopressors, but these may be administered through peripheral IV if unable to obtain central access. If in an SRC, pressors may be available. Consult ED physician.

- | | |
|---|--|
| • Norepinephrine 2-80 mcg/min IV initial infusion rate. | First-line vasopressor of choice in sepsis. |
| • Vasopressin 0.03-0.04 units/min. | May be added to norepinephrine to increase MAP or decrease norepinephrine dose. DO NOT use as a single agent. |
| • Epinephrine 1-40 mcg/min initially, titrated to effect. | May be added or used in place of norepinephrine to maintain adequate BP. |
| • Dopamine 2-20 mcg/kg/min. | Second-line option in highly select patients as it causes more tachycardia. |
| • Phenylephrine 40-160 mcg/min IV initial infusion until stabilized.
Titrate to usual range of 20-400 mcg/min. | Can be used as salvage therapy for refractive hypotension associated with tachycardia. |
| • Dobutamine 2-20 mcg/kg/min IV infusion. | May be used for inotropic support in the presence of severe myocardial dysfunction or hypoperfusion with depressed cardiac output. |

Corticosteroids

Corticosteroids should NOT be administered for the treatment of sepsis in the absence of shock.

If considering use of corticosteroids for septic shock refractory to pressors after euvoemia and appropriate antibiotic therapy achieved, consult ICU. The exception is giving dexamethasone prior to first dose of antibiotics for meningitis.



Normal Vital Signs for Age

(Source: Harriet Lane Handbook)

Age	Heart Rate (beats/min)	Respiratory Rate (breaths/min)	Blood Pressure (mm Hg)	Mean Arterial BP (mm Hg)
0-3 months	100-150	35-55	65-85 / 45-55	50 th percentile 55 + (age x 1.5)
3-6 months	90-120	30-45	70-90 / 50-65	
6-12 months	80-120	25-40	80-100 / 55-65	
1-3 years	70-110	20-30	90-105 / 55-70	5 th percentile 40 + (age x 1.5)
3-6 years	65-110	20-25	95-110 / 60-75	
6-12 years	60-95	14-22	100-120 / 60-75	
>12 years	55-85	12-18	100-135 / 65-85	

Empiric Antibiotic Choice

≤28 days

Ampicillin 75 mg/kg AND gentamicin 5 mg/kg. If concern for meningitis, give cefepime 50 mg/kg IV.
If concerned about HSV or neurologic impairment, add acyclovir 20 mg/kg.

>28 days

Ceftriaxone 100 mg/kg (max 2000 mg) AND vancomycin 20 mg/kg (max 2000 mg)
If CVL in place, immunocompromised, or significant Hx antibiotics in past 30 days
Cefepime 50 mg/kg (max 2000 mg) AND vancomycin 20 mg/kg (max 2000 mg)

If allergic to PCN

Meropenem 15 mg/kg (max 500 mg) AND vancomycin 20 mg/kg (max 2000 mg)

If suspecting Staph or Strep


Consider adding clindamycin 13 mg/kg IV for anti-toxin effect.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 9/16/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact
Amy_Carson-Strnad@ykhc.org.





RIGHT

MOTOR

KEY MUSCLES

Elbow flexors C5

Wrist extensors C6

Elbow extensors C7

Finger flexors C8

Finger abductors (little finger) T1

SENSORY

KEY SENSORY POINTS

Light Touch (LT)

Pin Prick (PP)

UER

(Upper Extremity Right)

Comments: (Non-key Muscle? Reason for NT? Pain?)

LER

(Lower Extremity Right)

VAC Voluntary Anal Contraction

(Yes/No) ☐

MOTOR SUBSCORES

UER ☐ + UEL ☐ = UEMS TOTAL ☐ (50)

MAX (25)

SENSORY SUBSCORES

LTR ☐ + LTL ☐ = LEMS TOTAL ☐ (50)

MAX (25)

NEUROLOGICAL LEVELS

1. SENSORY ☐ R ☐ L ☐

2. MOTOR ☐ R ☐ L ☐

3. NEUROLOGICAL LEVEL OF INJURY ☐ (NLI)

4. COMPLETE OR INCOMPLETE? ☐

Incomplete = Any sensory or motor function in S4-5

5. ASIA IMPAIRMENT SCALE (AIS) ☐

Zone of Partial Preservation

Sensory ☐ R ☐ L

Motor ☐ R ☐ L

Zone of Partial Preservation

Sensory ☐ R ☐ L

Motor ☐ R ☐ L

LEFT

MOTOR

KEY MUSCLES

Elbow flexors C5

Wrist extensors C6

Elbow extensors C7

Finger flexors C8

Finger abductors (little finger) T1

SENSORY

KEY SENSORY POINTS

Light Touch (LT)

Pin Prick (PP)

UEL

(Upper Extremity Left)

Comments: (Non-key Muscle? Reason for NT? Pain?)

LEL

(Lower Extremity Left)

VAC Voluntary Anal Contraction

(Yes/No) ☐

MOTOR SUBSCORES

LER ☐ + UEL ☐ = LEMS TOTAL ☐ (50)

MAX (25)

SENSORY SUBSCORES

LTR ☐ + LTL ☐ = LEMS TOTAL ☐ (50)

MAX (25)

NEUROLOGICAL LEVELS

1. SENSORY ☐ R ☐ L ☐

2. MOTOR ☐ R ☐ L ☐

3. NEUROLOGICAL LEVEL OF INJURY ☐ (NLI)

4. COMPLETE OR INCOMPLETE? ☐

Incomplete = Any sensory or motor function in S4-5

5. ASIA IMPAIRMENT SCALE (AIS) ☐

Zone of Partial Preservation

Sensory ☐ R ☐ L

Motor ☐ R ☐ L

Zone of Partial Preservation

Sensory ☐ R ☐ L

Motor ☐ R ☐ L

Patient Name _____

Date/Time of Exam _____

Examiner Name _____

Signature _____

REV 11/15

If time allows, please print this, complete it, scan it into the patient's MultiMedia Manager, and send with patient at time of transfer.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/24/23.

If comments about this guideline, please contact Travis_Nelson@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline

Status Epilepticus Treatment (Adult)

Definitions

- This guideline is indicated for the emergent treatment of convulsive status epilepticus.
- For atypical seizure-like presentations without evidence of impending hemodynamic instability, consult Neurology on call at ANMC or PAMC.
- **Convulsive status epilepticus:**
 - Seizure that lasts >5 minutes or occurs multiple times without regaining consciousness.
 - Diffuse, often tonic-clonic motor activity AND loss of consciousness.

Adult patient with seizure.

- ABCs and neurologic exam.
- **Bedside glucose STAT.**
- Obtain PIV x 2, continuous SpO₂, & cardiac monitor.
- Ensure BVM and suction at bedside.
- If possible, obtain labs (see box).
- Get AMPLE history.

Village Management

[See Emergency RMT Seizure Scenario on the wiki.](#)

- ABCs. Prepare BVM and suction.
- Place patient on floor with space around.
- **Bedside glucose STAT.** If unable to get a glucose measurement, give glucose buccally.
- Follow flow for no IV in place.
- Discuss with E1/E2 and activate medevac.
- If seizure resolves, place patient in recovery position.

After five minutes of seizure activity, patient meets criteria for treatment of convulsive status epilepticus (see Definitions box).

IV in place

No IV in place

If IV access unsuccessful, begin treatment with "No IV" pathway while continuing to attempt access and/or placing IO.

Labwork

Labs: BMP, Mg, Phos, CBC, lactate, EtOH, UDS, U/A, hCG. If concern for infection, send blood cultures and pro-calcitonin. Consider CK to trend over time.

- Lorazepam 0.1 mg/kg IV @ 2 mg/min AND
- Levetiracetam 60 mg/kg IV (max 4500 mg). Give over 15 minutes.

Seizure continues 5 more minutes after lorazepam given.

- Lorazepam 0.1 mg/kg IV @ 2 mg/min.
- Prepare for intubation.

Seizure continues 5 more minutes.

Fosphenytoin 20 mg PE/kg IV (max 1500 mg). Give over 10 minutes. If seizure continues, give additional 10 mg PE/kg IV over 5-10 minutes.

- Contact ICU and activate medevac.
- **Intubate** patient.
 - Induction (choose ONE): Propofol 2 mg/kg OR midazolam 0.2 mg/kg.
 - Paralysis: Rocuronium 0.6 mg/kg (preferred over succinylcholine due to risk of rhabdomyolysis and hyperkalemia, but recommend this lower dose)
 - Consider sugammadex following intubation to avoid masking seizure activity. Discuss with intensivist.
- Be prepared to give vasopressors or push-dose epinephrine if needed.

Choose ONE:

- Propofol drip 20 mcg/kg/min, titrate to effect with goal 50-80 mcg/kg/min. Watch BP closely.
- Midazolam drip 0.1 mg/kg/hr gtt, titrate to effect

- Discuss further management with ICU.
- Prepare for medevac.
- Continue active management until patient leaves, including continuous VS, frequent labs, and monitoring of UOP.

Benzodiazepine (choose ONE):

- Midazolam 0.2 mg/kg IM (max dose 10 mg) x1.
- Diazepam 0.2 mg/kg (max 20 mg) PR x1.
- Diastat home dose x1.
- AND
- Levetiracetam 60 mg/kg (max 4500 mg) PO (if able) or PR. To give PR, give tablets as well as one packet of water-soluble lubricant.

Seizure continues 20 more minutes.

- Activate medevac if in village.
- Fosphenytoin 20 mg PE/kg IM (max 1500 mg).

Seizure continues 20 more minutes.

Repeat benzodiazepine dose.

Seizure continues 20 more minutes.

Phenobarbital 20 mg/kg IM (max 1000 mg).

Treatments for Provoked Seizures

- Hypoglycemia: Dextrose 50% IV. Give 25 grams IV push.
- Hyponatremia: Sodium chloride 3% 100 mL infusion over 10 minutes.
- Hypocalcemia: Calcium gluconate 1-2 gram IV push.
- Eclampsia: Magnesium sulfate 4-6 grams IV over 20 minutes followed by 1-2 gram/hour.
- Alcohol withdrawal: Phenobarbital 260 mg IV push followed by 130 mg Q30-60 minutes.

Notes

- If seizure occurs in outpatient clinic, place patient on floor with space around and call a Rapid Response.
- Avoid using lorazepam IM due to erratic absorption.
- Avoid mixing different benzodiazepines.
- Monitor CK and renal function. Patient may require aggressive IV fluid administration if risk for rhabdomyolysis.
- Obtain neuroimaging if any focal abnormalities on neuro exam.
- Perform LP if unable to exclude intracranial infection. (Perform CT prior to LP.)

Post Seizure Care

- Seizure recurrence typically occurs within 2-6 hours.
- If history of seizures, may discharge with responsible adult if patient is improving. If first-time seizure, monitor in ED or clinic until mentation is at baseline. No air travel until >6 hours from event.
- Consider admission for prolonged post-ictal state or if concern for persistent metabolic abnormalities.
- Place urgent referral to Neurology if first-time seizure without known cause. Consult Neurology if considering urgent neurologic evaluation or medication initiation or adjustment.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 5/28/24. Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact Megan_Young@ykhc.org.



Use the [Pediatric Critical Care Guide](#) and [ED Peds Critical Care PowerPlan](#) to check all medication dosing.

- ABCs. Ensure BVM at bedside and pediatric code cart within reach.
- **Bedside glucose STAT.**
- Obtain IV. Start supplemental oxygen.
- Consult pediatrics.
- Obtain brief history.
- Prepare first-line medication. If in the ED or NW, get the Peds Seizure Kit (see box).

Village Management

[See Emergency RMT Seizure Scenario on the wiki.](#)

- ABCs.
- **Bedside glucose STAT.**
- If unable to get a glucose measurement, give glucose buccally.
- Get BVM with appropriate sized mask to bedside.
- Follow flow to the left, using these drugs with dosing found on Pediatric Critical Care Guide:
 - Diastat home dose PR if available or midazolam 0.2 mg/kg intranasal (max dose 10 mg) or diazepam 0.5 mg/kg (max 10 mg) IV solution given RECTALLY.
 - Phenobarbital or fosphenytoin (kept refrigerated) IM. If giving either second-line drug, consult pediatrics and strongly consider activating a medevac.
- Consider placing IV and giving NS bolus 20 mL/kg.
- Low threshold to activate medevac for atypical or prolonged seizure.

Go to [Pediatric Post-Seizure Evaluation](#) guideline.

Peds Seizure Kit

- In the ED and Peds NW Pyxis.
- Type "seizure" and override.
- Includes:
 - Midazolam 10 mg/2 mL
 - Levetiracetam
 - Phenobarbital 130 mg/mL
 - Dosing cards from the pediatric critical care guide

Continuous Infusions

In all ages, in consultation with the PICU, consider preparing for intubation and continuous infusion after second-line drug has been given. Continue giving medications as detailed in the flow while infusion is being prepared.

If giving midazolam, make drip of 1 mg/mL and start at rate 0.1 mg/kg/hour.


Indications for Admission or Transfer

- Status epilepticus
- Cluster of seizures
- Increased intracranial pressure
- CNS infection
- Structural lesion
- Patient does not return to baseline mental status.
- New focal neurological deficit requires further work-up (often MRI) even if CT is normal.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 2/9/24. Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Amy_Carson-Strnad@ykhc.org.



In all ages, if hemodynamic instability or myocardial dysfunction, avoid phenobarbital and use alternate agents.



Goals

1. Evaluate carotid and vertebral arteries for injuries.
2. Evaluate bony/cartilaginous and neck soft tissue structures.
3. Evaluate brain for anoxic injury.

Note: Life-threatening injuries can be present up to one year after strangulation event.

Helpful Links

- S/Sx strangulation in [adults](#) and [children](#)
- Physiological consequences [timeline](#)

Patient presents with concern for strangulation

Are ANY of the following present?

- Airway: subcutaneous emphysema (can be a sign of tracheal or laryngeal rupture)
- Neurological: loss of consciousness, seizures, mental status changes, amnesia, cortical blindness, movement disorders, stroke-like symptoms
- HEENT:
 - Visual changes: spots, flashing lights, tunnel vision, etc.
 - Facial, intra-oral, or conjunctival petechial hemorrhage
 - Odynophagia
- Neck:
 - Ligature mark, neck contusion, soft tissue injury, swelling, carotid tenderness, etc.
 - Dysphonia/aphonia, hematoma, laryngeal fracture, recurrent laryngeal nerve injury
- Bladder or bowel incontinence
- Pulmonary: dyspnea, phrenic nerve injury

Yes to ANY

No to ALL

Rule Out Life-Threatening Injuries

- If GFR ≥ 30 : CT angio of carotid/vertebral arteries. This is the gold standard for evaluation of vessels and bony/cartilaginous structures but is not very sensitive for soft tissue trauma.
- If GFR < 30 : non-contrast CT of neck. This study is less sensitive than CT angio for vessel injury but gives good visualization of bony and cartilaginous structures.

Injury identified

- Consult trauma surgery and plan to transfer.
- Consider ENT consult for laryngeal trauma with dysphonia.

No injury identified

How recent was event?

≥ 48 hours ago

< 48 hours ago

Observe in ED or admit to inpatient until 48 hours post-event based on severity of symptoms.

Is there reliable home monitoring AND a safe place to go?

Yes

No

- Discharge home with detailed instructions to return to ED if any neurological signs/symptoms, dyspnea, dysphonia, odynophagia dysphagia, or voice changes occur or worsen.
- Give custom Strangulation Patient Education handout.

- Consider discharge to TWC.
- May call TWC Crisis Line (543-3456) for assistance with safe shelter.
- Also may call SART on call at 545-4238 for further assistance.

Tundra Women's Coalition (TWC)

- Crisis Line: 543-3456
- Main office: 543-3444
- On-call advocate: 545-4328

Services Provided by TWC

- Emergency shelter
- Hospital accompaniment
- Information about community resources
- Legal advocacy
- Violent crime compensation
- Funds for emergency air or cab transportation

If patient would like to report incident:

- If occurred in a village: Alaska State Troopers 543-2294
- If occurred in Bethel: Bethel Police Department 543-3781

Use the following autotexts in your documentation:

- ..hpiStrangulation
- ..physStrangulation

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 2/9/24. Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact Jennifer_Prince3@ykhc.org.



Box 1: If responding to scene

- Do not risk safety of medical staff under any circumstance.
- If scene is compromised by combative patient or unsafe bystanders, leave scene immediately and do not return until scene secured by law enforcement.
- If CPR in progress, stay on-scene; CPR is often interrupted or lowered in quality by transport.
- Otherwise, prioritize transport to clinic. Aggressive medical interventions in field delay definitive care.

Trauma patient outside Bethel

- Identify mechanism.
- Transfer to clinic with Spinal Motion Restriction (SMR) if indicated.
- See Box 1.

Box 2: Common conditions which warrant emergent transport

- Physiologic instability: MAP <70, RR >30, GCS <10 if not intoxicated.
- Anatomic injuries: *penetrating* wounds to head, neck, torso, eye.
- Crushed/degloved/mangled extremity.
- Non-digital amputation.
- Pelvic fracture.
- Open/depressed skull fracture.
- Paralysis.

Trauma Primary Survey: **ABCDE**

- **Airway:** Loss of airway, stridor, expanding neck/submental swelling, impending airway compromise
- **Breathing:** Hypoxia, marked tachypnea, flail chest, absent breath sounds
- **Circulation:** Absent pulses, pulsatile bleeding
- **Deficit:** Objective neurologic deficit
- **Exposure:** Undeveloped patient, eval for occult injuries

Box 3: Contents of Focused HPI

Age, sex, mechanism of injury (MOI)

Details by MOI:

1. Penetrating trauma:
 - Knife: Type, length, depth.
 - GSW: Caliber, distance from victim, entrance/exit.
2. Blunt trauma:
 - MVC: Vehicle type, speed, ±LOC, ±ambulatory afterwards, ±restraint, ±helmet.
 - Fall: Distance, ±LOC, ±ambulatory afterwards.
3. Environmental
 - Cold Exposure: Temperature, time of exposure.
 - Heat Exposure: Structure/materials involved.

Additional important information:

- Anticoagulants
- Pregnancy
- Presence of burns
- Ability to void since injury

Emergent findings in Primary Survey
AND/OR
Any condition in Box 2?

Yes

- Contact Emergency RMT/Wards Doctor STAT.
- Stabilize and evaluate. See Box 4.
- Proceed to [secondary survey](#) after patient is stabilized.

No

Proceed to focused HPI (Box 3) and
and [secondary survey](#).

Findings on secondary survey warrant
transfer to higher level of care.

No

- Discharge with thorough return precautions.
- Feel free to contact RMT provider if questions.

Yes

Patient is cognitively intact,
hemodynamically stable, and ambulatory.

No

- Likely to require medevac.
- Contact Emergency RMT/Wards Doctor.

Yes

- Likely candidate for commercial transfer.
- Contact RMT provider to notify.

**Contact

- To reach Wards Doctor, send message via Tiger Connect to "Yukon Wards Doctor (Emergency RMT)" or "Kusko Wards Doctor (Emergency RMT)."
- If this is not practical, call the ED at (907) 543-6395 and ask for the wards doctor to be paged.

Please use this guideline as well as ATLS principles in all trauma cases, including for delayed presentation to care. Although delayed presentations are often less emergent, these principles still apply, and this process should be followed.

If health aide present, consider asking them to look up and follow CHAM section on Major Trauma.

Abbreviations

MAP: mean arterial pressure
GCS: Glasgow coma scale
SMR: spinal motion restrictions
LOC: loss of consciousness
MOI: mechanism of injury

Box 4: Interventions

1. Stabilization
 - Two 18g (or largest bore available) PIV
 - Spinal motion restrictions (SMR) if indicated
 - Pressure dressing to briskly bleeding wounds
 - Pelvic wrap/binder if indicated
 - Splinting of fractures
 - Do not apply a tourniquet without input from RMT or ED provider.
2. Diagnostics
 - CXR, AP Pelvis
 - Glucose POC, CBC, CMP

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/24/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Clinical_Guidelines@ykhc.org.



Secondary Survey Checklist

Document in your note using autotext “..traumasurvey”

	Mental Status: GCS
	Scalp: <ul style="list-style-type: none"> • Lacerations / swelling • Evidence of skull fracture
	Eyes: <ul style="list-style-type: none"> • Visual Acuity • Pupil size/reactivity • Globe integrity • Extraocular muscle movement
	Ears: <ul style="list-style-type: none"> • Hemotympanum • TM rupture
	Face: <ul style="list-style-type: none"> • Nose: Epistaxis, septal hematoma, fracture • Mouth: Midline, symmetric jaw, able to open and close.
	Neck: <ul style="list-style-type: none"> • Swelling / soft tissue injury • TTP over cervical spine
	Chest: <ul style="list-style-type: none"> • Ecchymoses, swelling, flail chest • TTP, crepitus, displaced ribs • Bilateral lung sounds
	Abdomen: <ul style="list-style-type: none"> • TTP, distension, absent bowel sounds
	Pelvis/GU: <ul style="list-style-type: none"> • Stability to pressure at the anterior superior iliac spine • TTP of femoral head • Testicular swelling • Blood at urethral meatus
	Back: <ul style="list-style-type: none"> • TTP along T/L spine
	Long bones: <ul style="list-style-type: none"> • Deformity/TTP • Lacerations over fractures (should be treated as open fractures) • Limitations in active ROM
	Integument (all sites): <ul style="list-style-type: none"> • Cold, pale, cap refill >3 seconds • Lacerations: <i>If not over vascular area, explore with sterile glove</i> • Hematomas (watch for expansion) • Burns

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/24/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Clinical_Guidelines@ykhc.org.




Yukon-Kuskokwim HEALTH CORPORATION

Villages without Health Aides: Management of Emergency Patients

The top priority is to ensure the safety of all involved.

- This includes staff, bystanders, and former health aides.
- CHAP on call is often privy to information about safety and may overrule a plan in the interest of keeping everyone safe.
- Bringing personnel from another village may be an option, but safety must be carefully considered, as trails are often unsafe, especially in bad weather.
- In these situations, emotions often run high. Please be careful not to coerce or strongly urge personnel to do something if they feel unsafe.



Note: If unable to reach CHAP on call, consult the VHAC Excel spreadsheet and call the numbers at the top. The ER techs all have access to the VHAC.

Capabilities of Personnel

Personnel may need instructions but are permitted to do the following with phone support from provider.

- **Office Assistant:**
 - CAN check VS
 - CAN give supplemental oxygen
 - CAN give OTC meds or patient's own meds
 - CAN help set up a nebulizer if patient supplies meds
 - CAN set up Zoom
 - CANNOT give prescription medications
- **Lay Rescuers:**
 - CAN do all of the above except access Zoom
- **Former Health Aide:**
 - CAN perform all tasks that were part of previous level of training as a health aide
 - CAN access med room if key is available
 - CANNOT access controlled substances
 - CANNOT access computer system, including Zoom


This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/13/23.

If comments about this guideline, please contact Ellen_Hodges@ykhc.org.



****Note:** When an adult patient discloses a sexual assault, it is his or her choice to report to law enforcement. Reporting without a patient's consent may breach his or her confidentiality. A SART exam/kit can only be conducted by a trained SAFE/SANE. Law enforcement must be involved, unless the patient wishes to have an anonymous kit collection. (See box.)



If patient is <18 years, please see [Pediatric Sexual Abuse Procedure](#).

SART Team Members

- Law enforcement
- SANE/SAFE (Sexual Assault Nurse Examiner/Sexual Assault Forensic Examiner)
- TWC advocate

Contact Information

- **Tundra Women's Coalition:**
Business Line: (907) 543-3444
Crisis Line: (907) 543-3456
Toll Free: (800) 478-7799
- **Law Enforcement:**
Bethel Police Department: (907) 543-3781
Bethel Post of Alaska State Troopers: (907) 543-2294
Aniak Post of Alaska State Troopers: (907) 675-4459
Emmonak Post of Alaska State Troopers: (866) 949-1303
St. Mary's Post of Alaska State Troopers: (907) 438-2019
- **National Sexual Assault Helpline:**
(800) 656-4673
Available 24 hours a day, 7 days a week.
- **YKHC SAFE/SANE:**
Tiger Connect: SART Adult On Call
(907) 545-4273

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/13/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Jennifer_Prince3@ykhc.org.



Indicators of Abuse: History

- No/vague explanation of significant injury
- Important details of explanation change dramatically
- Explanation of injury is inconsistent with the child's physical and/or developmental capabilities
- Injury occurred as a result of inadequate supervision
- Delay in seeking medical care without reasonable explanation
- Children with injuries resulting from family/ domestic violence incident
- Previous history of inflicted injury
- Inappropriate caretaker behavior that places child at risk

Indicators of Abuse: Physical Exam

Bruising

- Bruising in infants < 6months of age or non-ambulatory infants
- Bruising in unusual locations in any age child: ear pinna, neck, under chin, torso, buttock
- Pattern Bruises: loop marks, hand print, subgaleal hematoma due to hair pulling

Bite Marks

- Semi-circular/oval pattern
- May have associated bruising

Burns

- Pattern contact burns
- Cigarette burns
- Stocking/glove pattern
- Mirror image burns on extremities
- Symmetrical burns on buttock
- Immersion burns

Facial Injury

- Unexplained torn frenulum in non-ambulatory child
- Unexplained oral injury
- Ear injury

Injuries Suggestive of Abuse

Skeletal

- Rib fractures
- Multiple fractures
- Long bone fractures in < 6 months
- Any fracture (including femur) in non-ambulatory child
- Scapular fracture
- Sternum fracture
- Fractures of hands and feet

Head

- Subdural hematoma with or without skull fracture
- Unexplained intracranial injury (Note: Infants with intracranial injuries frequently have no or non-specific symptoms)

Poisoning

- Any illegal drug exposure, prescribed controlled substance, ethanol, or marijuana

Suspicion, allegation, disclosure, or confession of child physical abuse.
Please see Indicators of Abuse AND Injuries Suggestive of Abuse.

Treat acute issues as appropriate. If patient is in village and stable please arrange to have patient sent to ED via next commercial flight. If unstable then activate medevac.

Mandatory reporters must report via phone to:
OCS AND law enforcement (AST if incident occurred in village or BPD if incident occurred in Bethel).

- Complete appropriate work-up (see table). Use Child Abuse Power Plan.
- Take photos of any injury visible on exam, especially bruising. Take photos at a distance AND close-up to establish relative size and landmarks. Include ruler to establish scale.

Send RAVEN communication to Child Abuse Pool detailing reports made to Law Enforcement and **OCS**. May contact **Child Abuse On-Call** via Tiger Connect if any questions or concerns.

If unable to reach a discharge plan with OCS that YOU think is safe, then consider admission for safety and send message to **Child Abuse On-Call** to help reach a safe discharge plan.

Contacts

- Child Abuse On-Call via Tiger Connect. May email ChildAbuse@ykhc.org with nonurgent questions.
- Office of Children's Services (**OCS**): (800) 478-4444 or reportchildabuse@alaska.gov (CC Child Abuse team).
- Alaska State Troopers (**AST**): (907) 543-2294
- Bethel Police Department (**BPD**): (907) 543-3781
- Alaska CARES: (907) 561-8301

Mandatory Reporters include:

- Medical providers, nurses, health aides, teachers, social workers, law enforcement officers, and mental health professionals.
- Report should be made by every mandated reporter who has a concern, even if you think a report has already been made. This helps keep reports up to date with new information.
- Always document date and time of call, name of OCS representative, and what was reported.
- To complete the State of Alaska training for mandatory reporters, click [here](#).


Note: Minor injuries (single bruise on forehead, occasional bruises on shins, minor oral trauma, etc.) in a child able to cruise or sit independently can be part of normal development. Always ask caregivers for story behind injuries. If history does not match injury or child's observed developmental level, strongly consider child abuse injury surveillance.

Child Abuse Injury Surveillance Table (Use Child Abuse Power Plan.)

	<6 months	6-24 months	2-5 years	>5 years
Full exam	Yes	Yes	Yes	Yes
Skeletal survey Including oblique rib films	Yes	Yes	If highly suspicious of severe abuse	If highly suspicious of severe abuse
Head CT Request 3D reconstruction and 3 mm slices	Yes	If neurological exam abnormal	If neurological exam abnormal	If neurological exam abnormal
Abdominal labs AST, ALT, lipase, bag or CC U/A	Yes	Yes	Yes	If abdominal trauma
Bone labs Calcium, magnesium, phosphorus, alkaline phosphatase, intact PTH, 25-OH	If fracture	If fracture	If fracture	If fracture
Coagulation studies PT/INR, PTT, factor VIII & IX activity levels, VWF activity & antigen, CBC with diff. Consider CK if significant bruising. If head trauma PT/INR, PTT, thrombin time, fibrinogen, D-dimer	If bruising	If concerning bruising	If concerning bruising	If concerning bruising
Head circumference	Yes	Yes	N/A	N/A
Urine drug screen ± expanded state screen (contact Child Abuse On Call if considering expanded screen)	Consider	Consider	Consider	No
Optometry consult (within 24 hours)	If head injury	If head injury	If head injury	N/A

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/14/23. Click [here](#) to see the supplemental resources for this guideline. If comments about this guideline, please contact Jennifer_Prince3@ykhc.org.



Contacts

- On-Call Peds SART: (907) 444- 8643 or TigerText On-Call Peds SART.
- Child Abuse On-Call via TigerText. May email ChildAbuse@ykhc.org with nonurgent questions.
- Office of Children's Services (OCS): (800) 478-4444 or reportchildabuse@alaska.gov.
- Alaska State Troopers (AST): (907) 543-2294
- Bethel Police Department (BPD): (907) 543-3781
- Child Advocacy Center (CAC): (907) 543-3144 or (907) 545-1178

Alaska Age of Consent

- The age of consent is 16, provided the older partner is not in a position of authority (example: teacher, coach, minister).
- Any two people who are over the age of 16 can consent to sex in Alaska, but if one of the partners is under 16, and there is at least a 3 year age difference between the partners, it is illegal for them to have sex and must be reported.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 8/2/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Jennifer_Prince3@ykhc.org.



Source: ADA guidelines for treatment [here](#) and the abbreviated version [here](#).

Screen all overweight or obese adults with one or more other [risk factors](#) and all adults >35 years for type 2 diabetes mellitus.

Comorbidities and ASCVD Risk

Comorbidities must be evaluated at every visit. Document in chart and address Assessment and Plan where appropriate.

- ASCVD/CHF
- Hypertension
- Hyperlipidemia
- CKD
- Obesity
- Sleep apnea
- Tobacco and alcohol use
- NAFLD
- Hemoglobinopathies (including anemia)
- Major depressive disorder/general anxiety disorder
- Diabetes distress

Diagnostic Criteria

Unequivocal symptoms of hyperglycemia (thirst, polyuria, weight loss, and blurry vision) and either any one of the following OR any two of the following. (Take confirmatory test as close as possible to initial lab value to avoid treatment delays.)

- FPG* ≥ 126 mg/dl
- 2 hour PG ≥ 200 mg/dl during OGTT
- Hgb A1c ≥ 6.5
- RPG ≥ 200 mg/dl and symptoms of hyperglycemia or hyperglycemic crisis

Order CBC and iron profile if needed, as anemia can affect the accuracy of Hgb A1c.

Note: Fasting is defined as no caloric intake for at least 8 hours.

See diagnostic criteria.

Confirm diagnosis and add to problem list in RAVEN.

- Refer all new diagnoses of diabetes to the Diabetes Department.
- In RAVEN, type "Refer to Diabetes Internal" and select "DSMES (Diabetes Self Management Education and Support)," "MNT (Medical Nutrition Therapy)," and provider.
- Refer to Wellness Center for exercise education.

Schedule follow up appointment for 1-2 weeks and coordinate with diabetes department if possible.

At initial and annual diabetes visits:

- Review and complete health maintenance:
 - Foot exam
 - Labs
 - Immunizations
 - Mental health screening
- Encourage lifestyle changes (see box).
- Set [A1c target](#) based on age and risk factors or complication risk.
- Encourage purposeful blood glucose monitoring.
- Discuss family planning/sexual health.
- Refer to optometry, dental, audiology if needed, physical therapy if needed.

Remember: language matters.
See this [ADA resource](#).

Lifestyle Changes

- Advise 7-10% weight loss.
- Advise minimum 150 minutes of exercise per week.
- Advise traditional native diet with minimal carbs.
- Encourage [PLATE](#) method.
- Advise ≥ 7 -8 hours of sleep per night.
- Encourage DSMES participation.
- Limit alcohol consumption: one drink per day for females and two drinks per day for males.

For Optometry Referrals

- Either provider or patient must call Optometry at x6336 to schedule appointment.
- Provider must state in note that patient is to be referred to Optometry for a diabetic eye exam. This is necessary for travel to be arranged.

Abbreviations/Acronyms

ADA = American Diabetes Association
 ASCVD = Arteriosclerotic cardiovascular disease
 BH = Behavioral Health
 CGM = Continuous glucose monitoring
 CKD = Chronic kidney disease
 CMP = Complete Metabolic Profile
 DM = Diabetes mellitus
 DSMES = Diabetes self management, education, and support
 FPG = Fasting Plasma Glucose
 Hgb A1c or A1c for short = Hemoglobin A1c or glycosylated hemoglobin
 HTN = Hypertension
 MNT = Medical nutrition therapy
 OGTT = Oral Glucose Tolerance Test
 OSA = Obstructive sleep apnea
 PG = Plasma Glucose
 RPG = Random Plasma Glucose
 SMART = Specific, Measurable, Achievable, Realistic, Time-limited

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
 Approved by Clinical Guideline Committee 8/23/23.
 If comments about this guideline, please contact
Elizabeth_Tressler@ykhc.org.



Primary Treatment Goal of DM 2 is ASCVD risk reduction.

ABC's of DM 2 Care

- A1c: [individualized goal](#)
- BP $\leq 130/80$ (see antihypertensive box)
- Cholesterol (see lipids box)

Antihypertensives (in order of preference)

1. ACE/ARB
2. CCB/diuretic
3. Mineralocorticoid

Avoid beta-blockers unless necessary.

Medication selection is based on comorbidities and patient centered goals.

Always begin with lifestyle interventions. These are essential as medication response is often dependent on lifestyle measures.

Metformin: generally first-line unless true allergy, CKD, CHF, or ASCVD. Remember to use extended release and titrate.

Indicators of high-risk for ASCVD or established ASCVD, CKD, or HF?

Yes

No

Consider using a SGLT2i or GLP-1 RA independent of baseline A1c or A1c target. SGLT2i for CKD or HF and GLP-1 RA if ASCVD predominates.

Using shared decision making with patient, choose from any of the four classes: GLP-1 RA, SGLT2i, DPP-4i, TZD. Use GLP-1 RA or SGLT2i if weight loss/maintenance a goal.

Follow-up in 1-3 months.

If not achieving targets, continue to add classes of medications with the following suggestions:

- > Minimize hypoglycemia <
DPP-4i, GLP-1 RA, SGLT2i
For SU or basal insulin, consider agents with lower risk of hypoglycemia
- > Minimize weight gain/promote weight loss <
GLP-1 RA OR SGLT2i
- > Consider cost and access <
Certain insulins available at lower generic cost, SU

Shared decision making includes an educated and informed patient and their family/caregiver, patient preference, motivational interviewing, goal setting, ensuring access to DSMES, and empowering the patient.

Indications/Qualifications for CGM

- A1c ≥ 9 and/or prescribed insulin
- All CGM Rx for GDM patients must be prescribed by Compton
- All CGM Rx for non-pregnant patients must be prescribed by Nelson FNP

Follow up visits 1 month after adding new meds, OR every 3 months if stable, until lifestyle and A1c goals achieved.

If not achieving A1c goals, consider using CGM, revise SMART goals, utilize DSMES, DM support group, screen for Diabetes Distress or other psychosocial issues.

- If not achieving A1c goals and on four classes of medication including basal insulin, consider referral to ANMC Diabetes program and/or multidisciplinary discussion with diabetes team.
- Add prandial insulin as needed and ensure insulin teaching, self-management goals, and that patient is performing appropriate monitoring
- Continue to utilize a patient centered approach with shared decision making. Revisit lifestyle behaviors, patient specific motivators, psychosocial factors, and address medical comorbidities.
- To avoid therapeutic inertia, reassess and modify treatment regularly (3-6 months)

Diabetes Distress refers to negative psychological reactions to the emotional burden and patient worries specific to their experience of managing a complicated and demanding chronic disease. See ADA [position statement](#).

Abbreviations

- DPP-4i = dipeptidyl peptidase 4 inhibitor or gliptins. YKHC formulary saxagliptin (Onglyza).
- GLP-1 RA = glucagon-like peptide-1 receptor agonist. YKHC formulary liraglutide (Victoza).
- SGLT2i = sodium-glucose co-transporter-2 inhibitor. YKHC formulary empagliflozin (Jardiance).
- SU = sulfonylureas. YKHC formulary glipizide.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 8/23/23.
If comments about this guideline, please contact
Elizabeth_Tressler@ykhc.org.



Our pediatric endocrinologist, Dr. Rachel Lescher, has created the following protocols to aide us in managing patients with endocrinologic disorders. Please follow these recommendations.

As always, contact the pediatric hospitalist on call with any questions via the Tiger Connect role, "Peds Wards on Duty." We have access to the endocrinologist call/coverage schedule and can help direct consults as needed.

Endocrine Emergencies

Protocols for managing the following:

- Severe hypoglycemia
- Adrenal insufficiency/crisis (including patients with CAH)
- Hypercalcemia
- Hypocalcemia
- Thyrotoxic crisis (thyroid storm)

Diabetic Ketoacidosis

- Definitions and formulae
- Management
- Monitoring parameters
- Discussion and management of cerebral injury
- Prevention
- Sick day plans

Routine Follow-up of Endocrine Disorders

Protocols for managing the following:

- Congenital adrenal hyperplasia
- Congenital hypothyroidism/Hashimoto thyroiditis/goiter
- Hypopituitarism/septo-optic dysplasia/optic nerve hypoplasia
- Short stature work-up
- Growth hormone injections
- Insulin resistance/obesity
- Diabetes mellitus (type 1 and type 2)

These files were written by pediatric endocrinologist Rachel Lescher.

This is for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 9/25/23.

If comments about this protocol, please contact Leslie_Herrmann@ykhc.org.



Congenital Sucrase-Isomaltase Deficiency (CSID)

- This condition leads to an inability to digest sucrose (table sugar).
- Signs/symptoms:
 - Watery diarrhea after food containing sucrose
 - Abdominal pain/distension
 - Malnutrition, poor growth, FTT
- The condition is seen in Alaska Native people but is often under-diagnosed because patients unknowingly manage it with a traditional diet.

If you are considering this diagnosis, please consult a pediatrician. There are many more resources in the Pediatrics Folder on the vault, including sucrose content of medications and formulas.

CSID suspected

- Consult a YKHC pediatrician.
- Consult the [ANMC CSID Guideline](#) for details on testing.

- After CSID has been confirmed, treat with sacrosidase enzyme replacement (Sucraid). Sucraid is not covered by Medicaid, so there are many necessary steps.
- To obtain Sucraid:
 1. Go to: [Sucraid.com](#) → [How to Order](#).
 2. Click Physician Prescription Form.
 3. Fill out the information in the form with CSID as the diagnosis with 11 refills. (Must fill out this form annually.)
 4. Fax this form to the number at the top.
 5. Instruct family to fill out HIPAA form, found [here](#). This is the form to get the Sucraid for free via the financial assistance program.
 6. Fax this form to the number at the top.
 7. Get a reliable phone number for the family and tell them they must answer their phone when the company calls. They will need to give more information over the phone.
 8. Call the company to confirm everything has been arranged: 1-833-800-0122.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 6/1/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



Background Information

- 75% of the ANAI population is colonized with *H. Pylori* (range 61-84%).
- Screening or testing for *H. Pylori* for routine evaluation of dyspepsia or other GI symptoms is not clinically useful or supported by evidence for high prevalence populations.
- For routine clinical practice, there is **insufficient evidence-based data** to support community-wide eradication as a mechanism for gastric cancer prevention.
- Current literature **DO NOT** support a test and treat method with noninvasive tests.

Pediatrics

- Goal is to determine underlying cause of symptoms, not solely the presence of *H. pylori* infection.
- Diagnostic testing is NOT recommended with functional abdominal pain.
- Consult pediatrics if considering this diagnosis.
- Urgent referral for endoscopy for: occult blood in stool, weight loss, vomiting/hematemesis, melena, or clinical suspicion for gastric or duodenal peptic ulcer disease.
- Consider referral for: iron deficiency that is unexplained and refractory to treatment, malnutrition without other cause, chronic immune thrombocytopenia.

Pregnancy and Lactation

- Delay treatment until after pregnancy.
- DO NOT use in Pregnancy: bismuth and tetracycline.
- DO NOT use in lactation: bismuth, metronidazole, levofloxacin.

H. Pylori identified by histology and/or CLO test from EGD
AND

Endoscopy reveals the following:

- Duodenal ulcers
- Gastric ulcer
- MALT lymphoma
- Gastric intestinal metaplasia

Treat for *H. Pylori* with antibiotics

Adult Dosing (14 day course)

Preferred Treatment:

Metronidazole 500 mg PO QID
Amoxicillin 1000 mg PO BID
Omeprazole 20 mg PO BID
Bismuth subsalicylate 524 mg PO QID

PCN allergic (anaphylactic):

Metronidazole 500 mg PO QID
Doxycycline 100 mg PO BID
Omeprazole 20 mg PO BID
Bismuth subsalicylate 524 mg PO QID

Recurrence/Failure:

Metronidazole 500 mg PO QID
Doxycycline 100 mg PO BID
Omeprazole 20 mg PO BID
Bismuth subsalicylate 524 mg PO QID
OR
Amoxicillin 1000 mg PO BID
Levofloxacin 500 mg PO daily (FDA Black Box)
Omeprazole 20 mg PO BID

Pediatric Dosing (14 day course)

Metronidazole 10 mg/kg PO BID
Amoxicillin 25 mg/kg PO BID
Omeprazole 1 mg/kg PO BID
Bismuth subsalicylate
<10 years: 262 mg PO QID
>10 years: 524 mg PO QID

See [ANMC Helicobacter pylori Pediatric Treatment Guideline](#).

Other causes of dyspepsia that antibiotics will NOT help,
EVEN IF *H. Pylori* is detected:

- GERD
- Irritable Bowel Syndrome
- Mild/moderate gastritis
- Excessive/chronic NSAID use
- Heavy alcohol use
- Poor gastric mobility

Symptomatic Relief Medications

Adults:

Famotidine 10 mg PO BID
Omeprazole 20 mg PO BID

Children:

Famotidine 0.5 mg/kg PO BID

After Treatment


- Goal is eradication. Test of cure with Urea Breath Test (UBT) must be performed ≥ 8 weeks after end of treatment course. Must be off PPI ≥ 2 weeks prior to UBT. UBT can be performed on children as young as 3 years.
- For children, if UBT positive after treatment, consult Infectious Disease.
- For adults, if UBT positive, use one of the "Recurrence/Failure" regimens. If unsuccessful, consult Infectious Disease.
- The stool antigen test available at YKHC is not recommended for test of cure.
- 10-35% of individuals will fail treatment.
- Serologic testing is not recommended due to prolonged antibody persistence beyond date of cure and false positive results.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 4/28/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Clinical_Guidelines@ykhc.org.




Endoscopy Referrals

- Adults: Criteria for YKHC found [here](#). Use orders "Refer to Adult Surgery Internal – YK EGD" or "Refer to Adult Surgery External (only EGD/CS)."
- Pediatrics: "Refer to Peds Gastroenterology External" or "Refer to Peds Surgery External."



Note: this guideline does NOT apply to patients with acute blood loss requiring transfusion.

Adult male with Hgb < 13 g/dL or
Adult non-pregnant female with Hgb < 12 g/dL



Oral Iron Repletion

- Ferrous sulfate 325mg daily or every other day is generally acceptable
- No data to support vitamin C co-administration
- If GI side effects, consider ferrous gluconate 324mg daily or liquid ferrous sulfate

Candidates for IV Iron (See [Iron Infusion Guideline](#).)

- Elderly patients
- Hx gastric surgery, PUD, H pylori
- Hospitalized with acute bleeding needing transfusion (can give in same hospitalization)
- Heart failure with reduced ejection fraction
- IBD, celiac disease, other malabsorption syndromes
- Inability to tolerate oral iron
- Lack of response to oral iron

Findings diagnostic of iron deficiency:

- Ferritin < 30ng/ml or < 100ng/ml in patient with chronic inflammation
- Transferrin saturation (Tsat) < 20%

Findings supporting iron deficiency:

- RBC count low
- Corrected retic count low
- TIBC increased

References: UpToDate, Dynamed

B12 Reference Values

> 300pg/ml: Normal

200-300pg/ml: Borderline, check MMA/homocysteine

< 200pg/ml: Low

NB: High B12 in pt not on supplementation should prompt hematology referral due to risk of MDS

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 8/2/24.

If comments about this guideline, please contact clinical_guidelines@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline Iron Infusion for Chronic Iron-Deficiency Anemia (Adult & Pediatrics)

Iron-Deficiency Anemia Work-Up

- Evaluate for blood loss.
- Evaluate for dietary deficiencies.
- Labwork classically shows:
 - ↓ Hgb
 - MCV < 80
 - Ferritin < 30
 - ↑ TIBC
- Consider checking a lead level in children <6 years. See [lead screening guideline](#).

Diagnosis of iron-deficiency has been established.

Patient meets criteria for iron infusion, and patient or parent has agreed to infusion.

Provider places order "Refer to Infusion – Internal." Send message to "Infusion Center" role on Tiger Connect. Include patient's phone number.

Provider places future orders using "AMB IV Iron" or "PED Pediatric Iron Infusion" Power Plans.

- Provider updates Problem List with Iron-Deficiency Anemia.
- In the comments, provider states the plan (iron infusion with date ordered) and includes goal hemoglobin after infusions.

Infusion clinic nurse schedules patient for infusion. Case Managers write Letter of Medical Necessity. Village clinic arranges travel.

- Infusion(s) given per orders.
- All patients should have a follow-up hemoglobin level checked one month after infusion.
- If not at goal hemoglobin, patient should return to Bethel outpatient clinic for further evaluation.

See [Anemia in Pregnancy](#) guideline for indications in pregnancy.

Indications for Iron Infusion

If patient is hemodynamically unstable due to anemia, consider transfusion regardless of hemoglobin level. Ensure iron studies have been sent prior to transfusion.

- Hemoglobin between 5 and 7 in a hemodynamically stable, asymptomatic patient:
 - Patients <18 years: iron infusion likely indicated. Consult pediatric hematologist.
 - Patients ≥18 years: consider iron infusion alone vs transfusion followed by iron infusion based on clinical judgment.
- Hemoglobin between 7 and 8 with failure of oral iron therapy. Failure is defined as:
 - Minimal improvement in hemoglobin level despite at least two months of compliance with oral iron (in children 6 mg/kg/day; in adults ferrous sulfate 325 mg PO daily with ascorbic acid 500 mg PO daily)
 - Intractable GI side effects
 - Non-compliance after at least three attempts at oral iron therapy.
- Other patients may receive iron infusion if recommended by a hematologist.

Note: Patients <2 should have a hematology consult prior to beginning an infusion. The Infusion Center does not generally treat children <2, so they are generally admitted to Inpatient Pediatrics for iron infusions.

Iron Replacement Dose Calculation

$$\text{Total Iron Replacement Dose (in mg)} = 0.6 \times \text{weight} \times \left[100 - \left(\frac{\text{actual hemoglobin}}{\text{desired hemoglobin}} \right) \times 100 \right]$$

For pediatric patients:

- Using iron sucrose, this dose should be given in aliquots of 5-7 mg/kg until the full replacement dose has been given. Max dose is 100 mg for initial dose and 300 mg for repeat doses.
- Per Pediatric Hematology, may give children two iron sucrose doses 24 hours apart and then repeat two doses 24 hours apart Q1-2weeks until full replacement dose has been given. Giving more frequent dosing or more than two daily doses in a row results in decreased absorption and increased side effects in children.

For adult patients:

- Dose is typically iron sucrose 300 mg IV daily x3 doses.

Side Effects/Reactions

Efficacy and safety have been evaluated in adults and children older than two years. Consult pediatric hematologist for children younger than two years.

Specific reactions (rare):

- Hypersensitivity, including anaphylaxis and angioedema. Stop infusion immediately and treat as anaphylaxis.
- Hypotension (related to high total doses or rapid infusions). Stop infusion and treat with IVF, as appropriate.
- Infection: avoid administering if active systemic infection.
- For IV infiltrates, place cold pack.

Resources

- Consult Peds Wards On Duty by Tiger Connect.
- A pediatric hematologist can be reached for further questions at Alaska Pediatric Oncology at (907) 929-3773.
- ANMC Adult Hematology Oncology can be reached at (907) 729-1180.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 8/23/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



Background

- Only 4-9% of those...labeled [penicillin-allergic] are currently allergic. It is important to identify those who are not allergic, because children mislabeled as penicillin-allergic have more medical visits, receive more antibiotic prescriptions, and have longer hospitalizations with more antibiotic-related complications.¹
- Up to 10% of children develop rashes while receiving antibiotics. Most are diagnosed...as allergic to the implicated antibiotic, and most continue to avoid the suspect antibiotic in favor of alternatives, which may be less effective, more toxic, and more expensive.²
- Do not label a patient as allergic to penicillin/amoxicillin unless he or she has true hives, anaphylaxis, or a life-threatening reaction. Please include photos of rashes in RAVEN.
- Children labelled as allergic to penicillin/amoxicillin often carry that label for the rest of their lives.
- Please consult a pediatrician with any questions.

Anaphylaxis

- Acute onset – several minutes to hours from exposure.
- Generalized hives, pruritis or flushing, swelling of lips/tongue/uvula, and at least one of the following:
 - Dyspnea, bronchospasm, stridor
 - Hypotension
 - Evidence of hypoperfusion of end-organs
 - Persistent crampy abdominal pain, and/or vomiting or diarrhea

Hives vs Viral Rash

- True hives are raised, itchy, larger than dime-sized, come and go, move around the body, and change shape and size. True hives are uncomfortable. Ask if the rash bothered the child.
- Keep in mind that many parents refer to any rash as "hives." Get a description every time.
- A viral exanthem is typically diffuse, fine, pinpoint red dots and can be dense, coalesced, larger raised lesions. The rash typically covers the face and chest but can cover the whole body. The rash typically worsens and takes days to clear.

NOTE: If amoxicillin is needed to treat a life threatening infection, consult Allergy & Immunology to discuss possible desensitization. Alaska Asthma, Allergy, & Immunology can be reached at (907) 562-6228.

References

1. Kelso JM. "Provocation challenges to evaluate amoxicillin allergy in children." JAMA Pediatrics 2016;170(6):e160282.
2. Mill C, et al. "Assessing the diagnostic properties of a graded oral provocation challenge for the diagnosis of immediate and nonimmediate reactions to amoxicillin in children." JAMA Pediatrics. 2016;170(6):e160033.

Patient labeled with a penicillin/amoxicillin allergy.

Review history. (See box.)

Was the reaction anaphylaxis (see box) or other life-threatening reaction (eg Stevens-Johnson syndrome, etc.)?

Yes

- Do not give drug or perform trial.
- Update chart, including the Problem List and a comment on the allergy.
- Refer to Allergy & Immunology at 5.

No

What was the reaction?

Rash

Vomiting and/or diarrhea without any other S/Sx anaphylaxis

Other

Get more history. Consider pediatric consult.

True hives, skin blistering/peeling, or mucous membrane involvement

Viral-appearing rash or other type of rash

- Do not give drug or perform trial.
- Update chart, including the Problem List and a comment on the allergy.
- Refer to Allergy & Immunology at 5.

- Not a true allergy.**
- Educate and perform Amoxicillin Trial (see box).
 - If patient/family refuses trial, update Problem List.
 - Offer future trial or refer to Allergy & Immunology at age 5 for amoxicillin allergy testing.

- Not a true allergy.**
- Educate and perform Amoxicillin Trial (see box).
 - If patient/family refuses trial, update Problem List.
 - Offer future trial or refer to Allergy & Immunology at age 5 for amoxicillin allergy testing.

This guideline is designed for patients that are extremely unlikely to be allergic to amoxicillin. There is no upper or lower age limit. Consult pediatrics with any questions.

Amoxicillin Trial Procedure²

Use AMB Amoxicillin Trial Power Plan.

1. Obtain VS. Perform physical exam, including lung exam. Have appropriate dose of EpiPen or epinephrine. Epinephrine (1 mg/mL): 0.01 mg/kg (or 0.01 mL/kg) IM Q5-15 minutes. Per AAP recommendations:
 - 7.5-25 kg: use EpiPen Jr (0.15 mg)
 - ≥ 25 kg: use EpiPen (0.3 mg)
2. Calculate weight-based dose of amoxicillin. Give patient 10% of that dose.
3. Place patient in nearby room and instruct caregiver to notify staff of any changes in status.
4. If no reaction by 20 minutes, give patient remaining 90% of weight-based dose of amoxicillin.
5. Observe another 60 minutes. If no reaction, check VS and physical exam. If all stable, discharge home with regular course of drug.
6. Give patient and family amoxicillin trial education sheet.
7. Update allergy alert in RAVEN. Click the allergy in the banner. Right click over the drug name and choose "cancel." On the "reason" drop-down menu, choose "OK on Retrial."


Notes:

- If patient is on a beta-blocker, stop this for 24 hours prior to procedure, if possible. Beta-blockers can interfere with treatment for anaphylaxis, if it occurs.
- Ensure that patients with asthma have optimal control prior to this procedure.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner. Approved by Clinical Guideline Committee 8/23/23. Click [here](#) to see the supplemental resources for this guideline. If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



Suspected Botulism



Note: Botulism toxin only causes flaccid paralysis. Patients are awake, alert, and aware. Procedures should be explained and appropriate pain control and sedation for intubated patients should be provided.

Infant Botulism:

This is rare, with only 5 reported cases in AK in the past 65 years. If suspected, see Epi Procedure Manual, Botulism at State website.

Resources

- AK State Office of Epidemiology [Website](#):
-907-269-8000 (M-F, 8-5) and 800-478-0084 (after hours)
- State Lab [Website](#):
-1-855-222-9918
- Division of Public Health Healthcare Provider [Checklist](#)

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guidelines Committee 3/13/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact clinical_guidelines@ykhc.org.



Definitions

• Bronchiectasis is a lung condition with chronic wet cough and lung infections and is diagnosed by CT with contrast.

Use ICD10 code J47 – “Bronchiectasis.”

• Bronchiectasis risk is defined as ≥ 3 episodes of wet cough >4 weeks in the past 2 years, often in a setting of persistent infiltrates and recurrent pneumonia.

Use ICD10 code J41.1 – “Chronic purulent bronchitis.”

• All patients with either diagnosis should be made CPP and referred to pediatric pulmonology.

Stable Chronic Management

Comorbidities

• Aspiration: Trial thickener if <3 years, feed with swaddling in side-lying position at 45 degrees with slow-flow nipple, consider speech therapy.

• TB: Place PPD, send sputum/gastric aspirates if indicated (see [Pediatric TB Evaluation & Treatment guideline](#)).

• Asthma: Bronchodilators, inhaled steroids.

• Immunodeficiencies: Consider referral to Alaska Asthma, Allergy, & Immunology for work-up.

• CF: Confirm screen negative on newborn screen.

Maintenance Management

• Follow-up with pulmonology clinic Q3-6mo and pediatrician or health aide Q2-3mo to check symptoms and medications. At every visit:

- Patient and caregiver should verbalize diagnosis.
- Review plan for exacerbations.
- Check that Problem List is up-to-date with plan.

• Annual PFTs if >5 years.

• Annual sputum culture if chronic productive cough.

• Annual flu and COVID vaccines.

• Pneumococcal vaccines: PCV-13 series followed by one dose of PPSV-23 (Pneumovax) at ≥ 2 years.

• Treat dental caries.

• Optimize environmental health with woodstove safety, vents, irritant reduction, smoking cessation, etc.

• Airway clearance: P&PD/chest PT, consider acapella.

• Consider allergy testing.

Transition of Care

• Review diagnosis and management with patient and caregiver at each visit. Patient and caregiver should verbalize diagnosis, treatment, and exacerbation plan.

• At age 17, a pediatrician should review chart and refer patient to pediatric pulmonology for chest CT, treatment plan, and handoff visit.

• By age 18, a pediatrician should schedule a transition of care appointment with family medicine, write an Alert Note that includes a summary of medical history and current treatment plan, and refer to adult pulmonologist.

Exacerbation Management

Consider if:

- Persistent infiltrate >6 weeks
- or
- Chronic wet cough ≥ 4 weeks
- or
- Fever, increased wet cough, dyspnea, etc.

- Treat with Augmentin 45 mg/kg/dose BID or cefdinir 14 mg/kg/dose daily for at least 2 weeks.
- Consider probiotics.


• If able, do sputum culture (via RT in Bethel). If patient cannot produce sputum, use method described in [Induced Sputum Collection Checklist](#).

• Ask screening questions for dysphagia and have low threshold to thicken feeds.

• Chest physiotherapy TID.

• Recheck after two weeks.

• Consider systemic steroids if significant bronchospasm.





Molnupiravir

- Mechanism: The oral prodrug of a ribonucleoside with activity against RNA viruses.
- Regimen: 800 mg PO twice daily for five days. Initiate within five days of symptom onset.
- Main concerns: Risk of fetal toxicity.

Patients may request molnupiravir directly from the pharmacy.

See [Policy & Procedure](#) for details.

Criteria:

- Age ≥18 years.
- Mild to moderate disease in the outpatient setting
- High risk of progressing to severe illness.
- Alternative antiviral therapies not accessible or clinically appropriate.

No contraindications, warnings, or precautions. (See box.)

Counsel patient and document per requirements in box.

- Prescribe molnupiravir as soon as possible after positive COVID test and within five days of symptom onset.
- Patient should take molnupiravir 800 mg (four 200 mg capsules) PO twice daily for five days.

Adverse Reactions

In the clinical studies quoted in the EUA, the following adverse events were reported: diarrhea, nausea, and dizziness.

Contraindications, Warnings, and Precautions

- Molnupiravir is NOT authorized for use in patients who are hospitalized, requiring supplemental oxygen, or requiring more than their baseline supplemental oxygen flow rates due to COVID.
- Pregnancy: Due to risk of fetal toxicity, molnupiravir is NOT recommended for use during pregnancy.
- Breastfeeding: Not recommended to breastfeed during treatment period and for four days after the last dose. Instruct patients to pump and discard milk.
- Patients with childbearing potential:
 - Females: Instruct patients to use effective contraception during the treatment period and for four days after the last dose.
 - Males: Instruct patients with partners of childbearing potential to use effective contraception during the treatment period and for three months after the last dose.
- <18 years: Due to risk of bone and cartilage growth disruption, molnupiravir is NOT recommended for patients younger than 18 years old.

Documentation Requirements for Molnupiravir

Communicate and document the following in the medical record:

- [Fact Sheet for Patients and Parents/Caregivers](#) given to patient/caregiver.
- Inform patient/caregiver of alternatives to receiving molnupiravir. See [clinicaltrials.gov](#) for emerging data.
- Inform patient/caregiver that molnupiravir is an unapproved drug that is authorized for use under Emergency Use Authorization.

Reporting of Adverse Events

The prescribing health care provider is responsible for mandatory reporting of all medication errors and adverse events potentially related to molnupiravir. Reports must be made within seven days of the event.

Serious adverse events include: death; life-threatening adverse event; inpatient hospitalization or prolongation of existing hospitalization; persistent or significant incapacity or substantial disruption of the ability to conduct normal life function; congenital anomaly/birth defect; or medical or surgical intervention to prevent death, a life-threatening event, hospitalization, disability, or congenital anomaly.

Submit report to FDA MedWatch by completing the online form [here](#). The report should include "use of molnupiravir under Emergency Use Authorization (EUA)" in the "Describe Event" section.

See the [FDA MedWatch program](#) for more information.

Resource: Fact Sheet for Health Care Providers Emergency Use Authorization (EUA) of Molnupiravir.
Updated July 2023. Click [here](#) for source.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 9/25/23.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



Ritonavir-Boosted Nirmatrelvir (brand name Paxlovid)

- Mechanism: Nirmatrelvir is a protease inhibitor; ritonavir is a cytochrome P450 3A4 inhibitor that increases nirmatrelvir concentrations.
- Regimen: Paxlovid is packaged with nirmatrelvir 150 mg x2 and ritonavir 100 mg. Take all three pills (nirmatrelvir 300 mg and ritonavir 100 mg) PO twice daily for five days. Initiate within five days of symptom onset.
- Main concerns: Significant drug-drug interactions.

Criteria:

- Age ≥ 12 years and weight ≥ 40 kg
- Mild to moderate disease in the outpatient setting
- High risk of progressing to severe illness.
- No contraindications (see box).



- Providers should document using “..Paxlovid” autotext and comment on need for labs. May place this in an Alert Note and complete full documentation later.
- Consult a pharmacist. **A pharmacist must be involved with all Paxlovid prescriptions.** See [Policy & Procedure](#) for details.
- Patients may request Paxlovid directly from the pharmacy. See [Policy & Procedure](#) for details.

Note

- Ritonavir can have significant [drug-drug interactions](#). These interactions are increased with renal or hepatic insufficiency.
- Pharmacist involvement is essential in making adjustments to chronic medications and creating a patient-specific, tailored plan.

Indications for Labwork (CMP)

- Age ≥ 65 years.
- Hypertension, diabetes, or CVD
- Other chronic viral illness (HIV, Hepatitis C)
- Malignancy, autoimmune diseases, nephrolithiasis, or recurrent UTIs
- Chronic use of nephrotoxic medications
- Family history or past history of CKD
- Clinical judgment.

(May defer If checked in the last 12 months and no suspicion for worsening renal or hepatic impairment in that time.)

Adverse Reactions

In the clinical studies quoted in the EUA, the following adverse events were reported: dysgeusia, diarrhea, hypertension, and myalgia.

Contraindications

- Paxlovid is NOT authorized for use in patients who are hospitalized, requiring supplemental oxygen, or requiring more than their baseline supplemental oxygen flow rates due to COVID.
- Do not give to any patient with known hypersensitivity to any ingredient of Paxlovid.
- Review patient's medications (including herbal supplements) for drug-drug interactions, summarized at the [NIH COVID Treatment Guidelines](#).

Special Populations

- Pregnancy & Breastfeeding: There are no available data in these populations to use to make a recommendation.
- Renal Impairment:
 - Moderate (eGFR ≥ 30 to <60 mL/min): change dose to nirmatrelvir 150 mg (one tab) and ritonavir 100 mg (one tab)
 - Severe (eGFR <30 mL/min): not recommended
- Hepatic Impairment: not recommended if [Child-Pugh Score](#) Class C.



Signs of Impending Airway Compromise

- Drooling
- Lethargy
- Tripod position
- Marked retractions
- Tachycardia
- Cyanosis or pallor
- Rapid progression of symptoms

NOTE: Use extra caution in children with airway anomalies or ANY history of prior intubation.

Important Supportive Measures

1. Keep child upright or in position of comfort.
2. Turn lights down and minimize unpleasant interventions.
3. May take child outside for cool air.
4. Minimize invasive measures – keep child CALM!
5. **DO NOT** give albuterol; this can worsen croup.


In Village

If no racemic epinephrine available, mix 0.5 mL/kg of 1 mg/mL (1:1000) epinephrine (max dose 5 mL) with 1 bullet of NS and give via nebulizer.

DDx Stridor

- Croup (most common in ages 6 months to 3 years)
- Foreign body
- Tracheomalacia
- Angioedema
- Tracheitis
- Epiglottitis
- Abscess

Note: if prolonged symptoms (>3-5 days without any improvement), consider diagnosis other than croup.



The **Croup Severity Score** may be helpful in clinical decision making.

Low-Risk Criteria

- No stridor at rest
- Normal pulse-oximetry
- No increased WOB
- Good air exchange
- Normal color
- Normal mental status
- Tolerating PO
- Caregivers understand to return to clinic for recurrent stridor and/or increased WOB.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 11/21/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline

Fever $\geq 100.4^{\circ}\text{F}$ in Infants 0-90 Days

Infant 0-90 days with recorded or reported rectal temperature $\geq 100.4^{\circ}\text{F}$.

Sick-appearing

Well-appearing

Village Management

- If well-appearing and meets criteria for evaluation in Bethel, send on next commercial flight. If travel to Bethel will be delayed, infant must have recheck with health aide within 12 hours and be followed at least daily until seen in Bethel.
- If infant is not well-appearing, consult peds to discuss treatment options.
- If giving ceftriaxone IM in the village, DO NOT say "ceftriaxone per CHAM." Give the health aide the exact dose per box below.

- Evaluate patient in Bethel ED.
- Perform immediate work-up:
 - CBC with manual differential
 - Procalcitonin, CRP
 - Blood culture
 - U/A, urine culture
 - CXR
 - LP if stable
 - RSV/flu/COVID swabs

Admit or medevac, as appropriate.
Give antibiotics. (See box.)

0-28 days

29-60 days

61-90 days

- Evaluate patient in Bethel ED.
- Perform work-up:
 - CBC with manual differential
 - Procalcitonin, CRP
 - Blood culture
 - U/A, urine culture
 - Consider CXR, RSV/flu/COVID swabs

No temp > 101.3
AND
Fever resolves by 48 hours
AND
Meets all Low Risk History & Exam Criteria?

No Yes

Low Risk Criteria

History & Exam

- Well-appearing (soothable, feeding well, etc.)
- No significant prior hospitalization.
- Gestational age > 37 weeks.
- Neonatal course not complicated by: chromosomal anomaly, surgery (including G-tube), or infection.
- No evidence of invasive bacterial infection like cellulitis or osteomyelitis.

Lab

- Procalcitonin < 0.5
- CRP < 2
- Absolute neutrophil count (ANC) 1000-4000

CSF

- Do Multiplex PCR if any suspicion for meningitis.
- See Harriet Lane (not the results in RAVEN) for normal results by day of life.
- If concerned for bacterial meningitis, consult pediatrics and strongly consider medevac.
- Do not use correction formulas for traumatic LPs.

Meets all Low Risk Criteria (see box)?

No

Yes

- Perform LP.
- Give antibiotics. (See box.)
- Consider admission.

Evidence of UTI, AOM, or PNA?

Yes

No

- Consider LP based on clinical status, age, etc. LP not mandatory in well-appearing infants.
- Admission or daily follow-up until blood culture negative at least 36 hours.
- If UTI, treat empirically with cephalexin or ceftriaxone, pending speciation of culture.
- If AOM or PNA, use clinical judgment to decide on antibiotic treatment (oral or parenteral).

NOTE: If 22-28 days old and well-appearing with low-risk lab criteria, recent studies allow deferral of LP if admitted \pm antibiotics. Discuss with pediatrician and family if considering this option.

- Evaluate patient in Bethel.
- Perform work-up:
 - U/A, urine culture
 - Consider bloodwork and LP based on history, exam, and risk factors.

- LP and antibiotics unnecessary unless status changes.
- Daily follow-up until blood culture negative at least 36 hours.

- Daily rechecks in village, outpatient clinic, or ED until improving.
- Low threshold to bring to Bethel ED if any concerns.
- Don't give antibiotics without an in-person provider evaluation OR consulting a pediatrician.

Acetaminophen

- Acetaminophen should NOT be given prior to vaccines, as there is some evidence it blunts the immune response.
- Acetaminophen should NOT be given around-the-clock in this age group.
- Acetaminophen MAY be given after a fever has been documented and the infant evaluated by a health aide or provider **EXCEPT in babies 61-90 days old who are being managed in the village as this may blunt the fever curve. If a child in the village is already scheduled to come to Bethel for further evaluation, acetaminophen may be given.**

Antibiotic Treatment

- 0-7 days: Consult a pediatrician, pharmacist, or Neofax.
- 8-28 days:
 - If well-appearing and low suspicion for meningitis: **ampicillin** 50 mg/kg IV Q8h AND **gentamicin** 5 mg/kg IV Q24h.
 - If well-appearing and any suspicion for meningitis: **ampicillin** 75 mg/kg IV Q6h AND **cefepime** 50 mg/kg IV Q12h.
 - If ill-appearing and/or positive CSF Gram stain: please consult a pediatrician and/or a pharmacist.
- 29-90 days:
 - If low suspicion for meningitis: **ceftriaxone** 50 mg/kg IV/IM Q24h
 - If concern for meningitis: **ceftriaxone** 100 mg/kg IV once then 50 mg/kg IV Q12h AND **vancomycin** 20 mg/kg IV Q8h.
 - Continue IV/IM antibiotics until cultures are negative at least 36 hours and patient is clinically stable or until sensitivities are available to direct therapy.
 - Dose #2 of ceftriaxone may be given 12-24 hours after dose #1.

Special Circumstances

1. If fever within 48 hours of immunizations, well-appearing, and meets all history & exam low-risk criteria: no work-up necessary but must follow-up in village or Bethel within 12-24 hours. If fevers are rising or infant is not well-appearing, perform evaluation as above.
2. Pre-treatment with antibiotics but otherwise meeting low-risk criteria: infant must be observed a full 48 hours off antibiotics.
3. Unsuccessful LP: treat if appropriate and consider a repeat LP in 12-24 hours and determine treatment course based on cell counts. If repeat LP not performed or unsuccessful, either treat for 10-14 days with meningitic dosing of IV antibiotics or stop antibiotics at 48 hours and observe infant for an additional 48 hours off antibiotics. Consider admission.
4. If known HSV exposure, seizures, severe illness, hypothermia, lethargy, HSM, vesicular rash, conjunctivitis, interstitial pneumonitis, hepatitis, or thrombocytopenia: **acyclovir** 20 mg/kg IV Q8h with IVF, perform HSV work-up (CSF Multiplex PCR; CMP; blood HSV PCR; nasopharyngeal, conjunctival, and anal swabs and vesicle fluid for HSV PCR), and consult pediatrics.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 9/16/24. Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline Fever in Underimmunized Children (3-36 Months)

Please note: There is no expert consensus on this topic. These recommendations are adapted from [this](#) UpToDate article. As always, feel free to consult a pediatric hospitalist via Tiger Connect role "Peds Wards on Duty."

Note: Children with fevers lower than 102.2 may still require evaluation/work-up, especially if medical complexity, ill appearance, or any other concerns. For fevers lasting five days or more, consider Kawasaki disease or [MIS-C](#). Consult pediatric hospitalist.

Consult pediatric hospitalist.

Yes

Child 3-36 months old with
<5 days of fever $\geq 102.2^{\circ}\text{F}$.

Is child toxic-appearing or in
respiratory distress?

No

Is child fully immunized for
PCV and HiB? (See box.)

Yes

Work-up as appropriate.

No

Is there an identifiable source listed in box?

Yes

Treat as appropriate.
Encourage caregiver update vaccines as soon as possible.
If able, offer vaccines or outpatient appointment.

No

Perform work-up:
• CBC with differential
• Procalcitonin
• Blood culture
• Urinalysis and urine culture
• CXR

Note: The presence of AOM or viral URI does not decrease the risk of occult bacteremia. Thus, an underimmunized child with viral URI should still undergo this work-up; a work-up should be considered if the source is deemed to be AOM.

Is ANC $>10,000$ OR
is procalcitonin >0.5 ?

Yes

• Consider LP. May consult pediatric hospitalist.
• Give ceftriaxone 50 mg/kg IV or IM.

No

• Treat UTI if present. (See [guideline](#).)
• If child is well-appearing, is well-hydrated and tolerating PO intake, and has reliable caregivers, discharge with daily follow-up until blood culture is negative at 36 hours or sooner for any worsening.

Haemophilus influenzae Type A (HiA)

- There is a [high prevalence of HiA in rural Alaska](#). Our region has had several pediatric deaths due to HiA in the past five years.
- There are few evidence-based recommendations, as this is rare elsewhere.
- Children with HiA can have meningitis, septic arthritis, severe cellulitis, necrotizing fasciitis, osteomyelitis, pneumonia, and isolated bacteremia.
- In most of the local cases, the child had a fever >104 at some point in the course.
- Children with history of HiA are at risk for recurrence, especially in the first year after initial presentation. These children should have blood cultures drawn with any fever (>100.4) until at least 12 months after the initial infection.
- YKHC pediatricians recommend that any child with a fever $\geq 104^{\circ}\text{F}$ be considered at risk for HiA. Blood culture is not mandatory but should be considered. Good follow-up must be assured.

"Fully Immunized"

If children are under-immunized for pneumococcus (PCV) or *Haemophilus influenzae* type B (HiB), they are at a higher risk for occult bacteremia. Check vaccination status in the Immunization History tab in RAVEN and on VacTrAK for all febrile children.

• **PCV:** YKHC uses the Prevnar 20 vaccine. The recommended schedule is 2 months, 4 months, 6 months, and 12-15 months. **Children are considered "fully immunized" if they have received three doses by one year and four doses thereafter.** If series was started late or interrupted, consult [CDC vaccine schedule](#) to determine if child is "fully vaccinated."

• **HiB:** YKHC uses the PedVaxHib vaccine. The recommended schedule is 2 months, 4 months, and 12-14 months. **Children are considered "fully immunized" if they have received two doses by one year or three doses thereafter.** Exceptions:
- If first dose was given at 12-14 months, a second dose will be given >8 weeks for a total of two doses.
- If first dose was given at >15 months, no further doses are required.

NOTE: Some experts consider all children <6 months under-immunized and recommend a cautious approach. If choosing not to do a work-up, child must be followed daily until improvement is documented.

Identifiable Sources

Bacterial:

- Cellulitis and/or abscess
- Bone or joint infection
- Acute suppurative otitis media (see note to left)

Viral:

- Croup
- Influenza
- Herpetic stomatitis
- Hand-foot-mouth disease
- COVID-19
- Kawasaki disease




Testing

For thorough information about testing for influenza, please see [this page from the CDC](#).

Treatment

- For guidance on influenza treatment, please see [this page from the CDC](#).
- This includes a list of high-risk conditions that warrant treatment.
- Please note: Oseltamivir is a limited resource. Thus, the YKHC Antimicrobial Stewardship Program recommends that usage be limited to patients with additional risk factors for complications beyond Alaska Native or American Indigenous ethnicity.



Further Work-up

Perform careful exam for lymphadenopathy of other locations.
For any child with nontender lymphadenopathy or lack of improvement after specified period, consider, as appropriate:

- PPD/TB work-up
- CBC with diff
- CRP
- LFTs
- Blood culture
- HIV testing
- Testing for specific diseases (low threshold to consult ID if considering): syphilis, toxoplasmosis, *Bartonella*, *Brucella*, tularemia, HHV-6
- EBV, CMV titers
- LDH, uric acid
- CXR
- Hematology/oncology consult
- Infectious disease consult

Most Common Causes

- **Reactive lymphadenopathy** due to local infection, including dental (may take 4-6 weeks to resolve).
 - **Unilateral:** *Staph aureus*, Group A Strep, Group B Strep, anaerobes, TB, nontuberculous mycobacterium (NTM)
 - **Bilateral:** respiratory viruses (enterovirus, adenovirus, influenza, etc.), Group A Strep, HSV (primary), EBV, CMV, *Mycoplasma*, *Arcanobacterium*, TB
- Less Common Causes to Consider**
- Kawasaki disease; periodic fever with aphthous stomatitis, pharyngitis, and adenitis (PFAPA); leukemia; lymphoma; HIV; tularemia; brucellosis (if recent contact with caribou); *Bartonella*

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 8/2/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



The following is adapted from the "[ANMC Pediatrics Statement on Dexamethasone and Hearing Screening in Meningitis](#)," dated 2/4/20.

Haemophilus influenzae type A

In recent years, *Haemophilus influenzae* type A (HiA) meningitis has been more common than other causes of bacterial meningitis in children admitted to ANMC. Many of these children have been transferred from YKHC. See this [State Epidemiology Bulletin](#) for information about Alaska cases in 2014-2018, including the outbreak in 2018.

The pattern of disease in HiA is similar to that seen in *Haemophilus influenzae* type B (HiB) meningitis. In HiB meningitis, dexamethasone has been shown to decrease the incidence of severe hearing loss. In Alaska, there have been multiple cases of sensorineural hearing loss associated with HiA meningitis. It is suspected that dexamethasone may confer similar benefits in HiA meningitis. As a result, our local experts (including infectious disease and endocrinology experts) recommend giving dexamethasone with all cases of suspected bacterial meningitis.

Dexamethasone

- **Indications:** A child >6 weeks old with clinical meningitis or visibly purulent spinal fluid.
- **Timing:** First dose should be given 10-20 minutes prior to or concurrent with the first dose of antibiotics; if given after antibiotics have been given, there is no evidence that dexamethasone will improve outcomes.
- **Dose:** Dexamethasone 0.15 mg/kg/dose IV.
- **Course:** If dexamethasone is initiated and HiA/HiB is confirmed, continue dexamethasone 0.15 mg/kg/dose IV Q6h for 2-4 days. If CSF culture/PCR show a different pathogen or are negative, stop the dexamethasone.

Hearing Screening

- All children with bacterial meningitis should be referred to audiology.
- Hearing evaluation should be scheduled one month after hospital discharge.



Tecovirimat (also called TPOXX)

- Tecovirimat is an inhibitor of the orthopoxvirus VP37 envelope wrapping protein, which prevents the formation of egress-competent enveloped virions necessary for cell-to-cell and long-range dissemination of virus.
- Tecovirimat is approved by the FDA to treat smallpox under the [Animal Rule Regulations](#). It has not been studied in humans, as smallpox has been eradicated globally, and exposing people to smallpox virus for the purpose of a clinical trial is not ethical.
- Tecovirimat has also not been studied in the treatment of mpox. However, the FDA and CDC have an expanded access Investigational New Drug (IND) protocol (also known as “compassionate use”) that allows tecovirimat to be used to treat mpox under strict requirements. The drug is only available from the Strategic National Stockpile.
- YKHC’s Institutional Review Board has approved the use of tecovirimat as long as the [CDC/FDA protocols](#) are followed.

CDC Expanded Access IND Protocol

Patient with lesions suspicious for mpox with an indication to consider tecovirimat (see box).

Swab lesions and send for [testing](#).
Note: decision to treat is clinical, as results take a long time to return.

Counsel patient/family on tecovirimat, including experimental nature and possible risks.
May use this [informed consent](#) form to guide counselling.

If patient/family agrees to tecovirimat, contact COVID-19 Response Team by Tiger Connect.
Note: only providers on the IND registry may prescribe tecovirimat; thus, the COVID team has agreed to manage this drug.

- COVID team will discuss further with patient, prescribe drug, complete all required IND forms, complete consent form, and document in RAVEN.
- Standard course of therapy is 14 days. See [protocol](#) for dosing and mixing instructions.
- COVID team will do periodic telehealth visits with patient to monitor progress and document in RAVEN.

Please see [these resources](#) for pain management and other supportive measures.

Indications to Consider Tecovirimat

- **Severe disease:** hemorrhagic disease; large number of confluent lesions; sepsis; encephalitis; ocular or periorbital infections; or other conditions requiring hospitalization.
- **Involvement of anatomic areas which might result in scarring or strictures:** lesions directly involving the pharynx causing dysphagia, inability to control secretions, or need for parenteral feeding; penile foreskin, vulva, vagina, urethra, or rectum with the potential for causing strictures or requiring catheterization; anal lesions interfering with bowel movements (for example, severe pain); and severe infections (including secondary bacterial skin infections), especially those that require surgical intervention such as debridement.
- **Severe immunocompromise:** advanced or poorly controlled human immunodeficiency virus (HIV), leukemia, lymphoma, generalized malignancy, solid organ transplantation, therapy with alkylating agents, antimetabolites, radiation, tumor necrosis factor inhibitors, or high-dose corticosteroids, being a recipient of a hematopoietic stem cell transplant <24 months post-transplant or ≥24 months but with graft-versus-host disease or disease relapse, or having autoimmune disease with immunodeficiency as a clinical component.
- **Pediatric populations:** particularly patients younger than 8 years of age.
- **Pregnant or breastfeeding people**
- **Concurrent conditions affecting skin integrity:** atopic dermatitis, eczema, burns, impetigo, varicella zoster virus infection, herpes simplex virus infection, severe acne, severe diaper dermatitis with extensive areas of denuded skin, psoriasis, or Darier disease (keratosis follicularis).

Contraindications & Risks

- Patient or legally authorized representative unwilling to sign an informed consent and refuse tecovirimat treatment
- Known allergy to tecovirimat and/or inactive ingredients in tecovirimat
- For IV tecovirimat only: patients with severe renal impairment (CrCl <30 mL/min)*. Oral tecovirimat is an option for patients with severe renal impairment.
- Co-administration with repaglinide may cause hypoglycemia. Monitor blood glucose and monitor for hypoglycemic symptoms during co-administration.

Adverse Reactions

In a Phase 3 clinical trial, the most common reported events were headache; nausea; vomiting; abdominal pain; and infusion site pain, swelling, erythema, and extravasation. Other events were reported in <2% of patients.

Reporting of Adverse Events

The prescribing health care provider is responsible for mandatory reporting of all medication errors and adverse events potentially related to tecovirimat. Reports must be made within seven days of the event.

Serious adverse events include: death; life-threatening adverse event; inpatient hospitalization or prolongation of existing hospitalization; persistent or significant incapacity or substantial disruption of the ability to conduct normal life function; congenital anomaly/birth defect; or medical or surgical intervention to prevent death, a life-threatening event, hospitalization, disability, or congenital anomaly.


Submit report to FDA MedWatch by completing the online form [here](#). The report should include “use of tecovirimat under Emergency Use Authorization (EUA)” in the “Describe Event” section.

See the [FDA MedWatch program](#) for more information.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline Multisystem Inflammatory Syndrome in Children (MIS-C)



Case Definition for Multisystem Inflammatory Syndrome in Children (MIS-C) According to the CDC & CSTE

An individual <21 years presenting with:

1. Measured or subjective fever $\geq 100.4^{\circ}\text{F}$.
2. Laboratory evidence of inflammation with CRP ≥ 3.0 mg/dL.
3. Evidence of clinically severe illness requiring hospitalization with new onset manifestations in at least two of the following categories:
 - **Cardiac:** elevated troponin (or specific echo findings)
 - **Mucocutaneous:** rash, inflammation of the oral mucosa (eg, mucosal erythema or swelling, drying or fissuring of the lips, strawberry tongue), conjunctivitis or conjunctival injection, or extremity findings (eg, erythema or edema of the hands or feet)
 - **Shock**
 - **Gastrointestinal:** abdominal pain, vomiting, or diarrhea
 - **Hematologic:** platelet count $<150,000$ cells/ μL or absolute lymphocyte count (ALC) <1000 cells/ μL
4. Absence of a more likely alternative diagnosis.
5. Evidence of or high suspicion for current or recent (within the last four weeks) COVID-19 infection.

Differential Diagnosis

- Kawasaki disease and other vasculitides
- Severe acute COVID-19
- Sepsis
- Toxic shock syndrome
- Appendicitis
- Viral infection like EBC, CMV, adenovirus, and enterovirus (although unlikely to cause severe multisystem disease if immunocompetent)
- Rare syndromes including HLH/MAS and SLE

NOTE: MIS-C is a reportable disease. Please ask the accepting facility who should make the report. The form can be found [here](#).

Note: Our understanding of MIS-C is evolving. At the time of publication, the standard of care for MIS-C treatment includes IVIg, so YKHC pediatricians recommend transfer to a higher level of care for all patients who meet criteria for MIS-C. However, if diagnosis is uncertain, admission with observation at YKHC may be appropriate. Have a low threshold to consult infectious disease and pediatric cardiology if considering this.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 9/16/24.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



Box 1: AOM Decision-Making Principles

- If observation is warranted, do not prescribe antibiotics.
- Always treat pain.
- If patient has not received amoxicillin within 30 days, start with amoxicillin to treat new infection.
- Do not treat fluid that develops after AOM if child is asymptomatic – observe up to 3 months.
- Do not use antibiotic prophylaxis.
- Do not send ear drainage for culture.

Box 2: Eligibility for Observation for 48-72 hours

- 6-24 month old with mild, uncertain, or unilateral AOM.
- >24 month old with mild/moderate (non-bulging) AOM.
- Caregiver comfortable withholding antibiotics.
- Follow-up assured.
- Antibiotics can be started promptly if symptoms persist or worsen.
- No fever >102°F and only mild otalgia.
- No otorrhea (unless tympanostomy tubes present).

Box 3: AOM Treatment

Antibiotic duration, by age:

- < 2 years: 10 day course of oral antibiotic
- 2-5 years: 7 day course of oral antibiotic
- ≥ 5 years: 5 day course of oral antibiotic
- Note: in patients with TM perforation or history of recurrent/complicated/chronic infections, treat for 10 days.

Antibiotic choice:

- 1st line: amoxicillin 45 mg/kg/dose PO BID
- 2nd line: Augmentin 45 mg/kg/dose PO BID
- 3rd line: cefdinir 14 mg/kg/dose PO QD
- OR ceftriaxone 50 mg/kg IV/IM QD for 1-3 days

Otitis-conjunctivitis syndrome

Augmentin 45 mg/kg/dose PO BID

Try to avoid using cephalosporins. They are less effective at treating the most common organisms that cause OM.

Do not use azithromycin, erythromycin, cephalexin (Keflex), or Septra for AOM.

For PCN allergy: Please refer the patient for an allergy trial if not already done.

cefdinir 14 mg/kg/dose PO QD

OR

ceftriaxone 50 mg/kg IV/IM QD for 1-3 days

AOM with Otorrhea

- If patient has ruptured TM and no tubes, treat with oral antibiotics for ten day course with dosing as above and otic antibiotics. Oral antibiotics may improve TM healing.
- If patient has tympanostomy tubes that are confirmed to be still in place, may treat with otic antibiotics only.

Otic Antibiotics

Wick ears prior to giving drops. After instilling drops, child should lie with affected side up for several minutes.

- Ciprofloxacin 4 drops BID for 7 days
- Ciprofloxacin + dexamethasone 4 drops BID for 7 days


AOM ≥3 months

Acute onset of:

- Fever and ear pain
 - Bulging TM and decreased mobility
- See Box 1.

Always address pain:

- If >3 months old, use acetaminophen.
- If >6 months old, use acetaminophen and/or ibuprofen.



AOM <3 Months Old

If otorrhea, bulging TM, or other suspicion of AOM <3 months old, patient must be seen by provider within 24 hours.

- ≤28 days old: patient must be seen in the ER for full lab work-up including LP and treatment with IV antibiotics.
- 29-60 days old with or without fever, patient must be seen in the ER for evaluation. Even if no fever, follow recommendations on [fever <90 days clinical guideline](#).
- 61-90 days old:
 - If febrile, follow [fever <90 days clinical guideline](#).
 - If afebrile and sick-appearing, perform work-up as clinically appropriate. May consult peds as needed.
 - If afebrile and well-appearing, lab work-up not necessary. May treat with antibiotics as appropriate.

AOM via RMT

If considering antibiotics for AOM, always request that health aide sends photos of the tympanic membrane.

Consider **Otitis Media with Effusion (OME)** if no acute symptoms but decreased TM mobility. Non-infected fluid may persist for 3 months after AOM.

Tympanostomy Tubes

- **Indications:** OME for at least three months or recurrent episodes of AOM with at least three episodes in the past six months or at least four episodes in the past year (with at least one in the past six months).
- **Process:** Place order for "Refer to Audiology Internal." Audiology at YKHC will evaluate the child and refer to ENT if indicated.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/13/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Amy_Carson-Strnad@ykhc.org.



Symptoms of Peritonsillar Abscess/Cellulitis

- Progressively increasing throat pain and swelling
- Muffled speech / change in voice
- Neck pain, typically unilateral
- Fevers, chills, myalgias
- Dysphagia, odynophagia

Labs

- CBC, BMP, CRP
- If needle aspiration, culture aspirate
- If SIRS or qSOFA ≥ 2 , add lactate, procalcitonin, blood cultures

- Prepare for **intubation**. Anticipate difficult airway. Consider calling CRNA.
- Place IV; get labs. (See box.)
- Give antibiotics. (See box.)
- Transfer to higher level care.

Impending
airway
compromise?

Yes

No

Signs/Symptoms of Impending Airway Compromise

- Drooling
- Patient in "sniffing position" (leaning forward)
- Anxious appearance with suprasternal retractions with or without stridor

Indications for CT Soft Tissue Neck with IV Contrast as Part of Initial Workup

- Toxic appearance
- Submental tenderness to palpation
- Neck stiffness, swelling, or pain with extension

Village Management

- Amoxicillin/clavulanic acid (preferred)
- If unable to swallow, IM penicillin OR ceftriaxone + clindamycin
- Ketorolac/acetaminophen
- Consider dexamethasone 10 mg.

Commercial flight to Bethel ER; discuss with ED MD if concern for airway compromise.

Consider CT. (See box.)

Examination with ANY?

- Trismus
- Uvular deviation
- Peritonsillar swelling

No

Examination with
peritonsillar erythema?

No

Acute pharyngitis

Yes

Attempt **needle
aspiration**

No pus

Peritonsillar cellulitis

Pus

Peritonsillar
abscess

- Place IV; get labs. (See box.)
- Give antibiotics. (See box.)
- IV fluids.
- Consider dexamethasone 10 mg.
- Analgesia (non-opioid first).
- Monitor in ED minimum 3 hours.

Improving

Discharge on PO
antibiotics with
recheck in 24 hours.

Not improving

Obtain CT neck with contrast.

Deep tissue abscess

Consult ENT.

No deep tissue abscess

Admit to inpatient on IV antibiotics.

Microbiology & Antibiotics

Continuum from pharyngitis > cellulitis/phlegmon > abscess. Often polymicrobial, typically GAS, *Strep viridans*, *Staph aureus*, fusobacterium, bacteriodes. MRSA coverage not indicated unless patient does not respond to initial antibiotic selection.

IV

Ampicillin/sulbactam 3 grams Q6h (preferred)
OR
Piperacillin/tazobactam 3.375 grams Q6h
OR
Ceftriaxone 1 gram Q12h + metronidazole 500 mg Q6h
OR
Clindamycin 600 mg Q6-8h (if penicillin allergy)

PO

Amoxicillin/clavulanate 875 mg BID (preferred)
OR
Cefpodoxime 300 mg Q12h + metronidazole 500 mg Q6h
OR
Clindamycin 300 mg Q6h (if penicillin allergy)

Treatment duration 14 days.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved 8/2/24. Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact Travis_Nelson@ykhc.org.



Maintain High Index of Suspicion

1. Classic findings include inspiratory whoop and staccato cough.

2. Infants often do not have the "whoop."

This [video](#) from the Mayo Clinic is a great example of a classic infant presentation.

3. Pertussis is predominantly a clinical diagnosis: If you are very suspicious (especially in babies), treat empirically while awaiting test results. Do not report to state until confirmed.

Prevention

1. Update DTaP and Tdap for anyone eligible. [Here](#) are the CDC vaccine schedules, including catch-up.

2. Pregnant patients are due for Tdap after 28 weeks. This gives protection to the baby.

The [Alaska Department of Health Pertussis page](#) includes detailed information.

Isolation

• Recommend patients isolate and wear a mask around others for all respiratory symptoms.

• If test is positive, recommend isolation until five days of treatment have been completed.

Tests Available Through YKHC

(Use a regular viral respiratory swab with a red top to collect both tests.)

• "Respiratory Panel 2 In-House" is a large panel of respiratory tests, including pertussis. Very expensive but turnaround time is several hours. Given expense, should only be used on young or highest risk patients.

• "B. pertussis and B. parapertussis LC" is a send out test to LabCorp. Cheaper option but longer turnaround time (at least 7 days).

Medication Regimens

(Same regimen for both treatment and prophylaxis.)

• **< 6 months:** azithromycin 10 mg/kg PO daily x5 days.

• **≥ 6 months:** azithromycin 10 mg/kg PO x1 then 5 mg/kg PO daily x4 days.

• **Adults and patients ≥50 kg:** 500 mg PO x1 then 250 mg PO daily x4 days.

If true macrolide allergy:

• **≥ 2 months:** Septra 4 mg/kg TMP PO twice daily x14 days.

• **Adults and patients ≥40 kg:** Septra DS (160 mg TMP) PO twice daily x14 days.

Pertussis is suspected.

Droplet precautions for all suspected cases.

Who to test:

- **< 6 months:** suspicious symptoms OR exposure with ANY symptoms: Test with "Respiratory Panel 2 In-House."
- **≥ 6 months:** with suspicious symptoms OR exposure with ANY symptoms: Test with LabCorp test.
- Do not test completely asymptomatic people.
- Any household contact of a known case may be treated without a test.

Who to treat:

- Patients <12 months within 6 weeks of cough onset. If high level of suspicion for patients at high risk, treat empirically while awaiting test result.
- Patients >12 months within 3 weeks of cough onset. If high level of suspicion for patients at high risk, treat empirically while awaiting test result.
- Pregnant patients (especially if near term) within 6 weeks of cough onset.
- Any household contact of a known case may be treated without a test.
- [This document](#) from State Epi provides more guidance about who to treat.

Factors to consider about hospitalization:

- Infants <4 months:
 - Check CBC with diff.
 - Low threshold to hospitalize these infants until they have begun to show some improvement.
 - Risk factors for significant morbidity (including "rapid, unpredictable deterioration"): apnea, true cyanosis, and WBC >30K. If any of these are present, consider transfer to a facility with a PICU.
- Older patients: Consider hospitalization and/or empiric treatment for patients with history of prematurity, chronic lung disease, neuromuscular disorders, etc. Feel free to consult Peds Wards on Duty with any questions.

Pertussis is confirmed.

- Provider must report to State Epi within two business days. May call (907) 269-8000 or fax [this form](#) to (907) 561-4239.
- Provider will send message to Population Health team via Tiger Connect. Include patient name, DOB, MRN, and any other information (if any contacts are known, etc.).

- Population Health will work with Public Health Nurses to identify contacts needing treatment.
- These names will be sent to Pertussis Response Providers, who will prescribe medications as needed.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 10/23/24.

Click [here](#) to see the supplemental resources for this guideline.


If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



For thorough information about the diagnosis and treatment of Streptococcal pharyngitis, please see [this page](#) from the CDC.

Other Considerations:

- Consider testing for oral GC/CT in at-risk populations.
- Testing for Group A streptococcal (GAS) pharyngitis is NOT recommended for acute pharyngitis with clinical features that strongly suggest viral etiology (e.g. cough, rhinorrhea, etc).
- Routine use of back-up cultures for those with a negative rapid test is not needed for adults; there is a low incidence of GAS in adults and risk of subsequent acute rheumatic fever is exceptionally low.
- It is NOT recommended to test for GAS in patients under the age of 3; the risk of rheumatic fever in this age group is exceptionally low.
- Patients are contagious for up to 24 hours after starting antibiotic treatment.
- Treatment for asymptomatic GAS carriers is not recommended, nor is testing or empiric treatment of household contacts.
- Refer to [Peritonsillar Abscess guideline](#) if appropriate





Transfer to Higher Level of Care

Criteria

- Requires >2-3 L supplemental oxygen to prevent hypoxia or improve WOB.
- Witnessed apnea.
- Requires neb treatments more frequently than Q2-3h for >8 hours.
- Sustained tachycardia, tachypnea, or respiratory distress despite treatment.
- Significant pleural effusion.

Antibiotics per box.

Supportive care per [Respiratory Distress Guideline](#).

Consider [high flow nasal cannula](#).

Inpatient Treatment at YKHC

Criteria

- Requires supplemental oxygen to prevent hypoxia or improve WOB. If requiring >2 L NC, reevaluate whether patient is appropriate to stay at YKHC.
- Requires IV or NG fluids.
- Question of apnea.
- Not tolerating home therapy or unreliable follow-up.
- Does not meet criteria for transfer to higher level of care.

Antibiotics per box.

Supportive care per [Respiratory Distress Guideline](#).

Outpatient Treatment

Criteria

- WOB is mild or absent.
- No hypoxia.
- Able to maintain hydration without IVF.
- Tolerating home therapy with reliable caregivers.
- No apnea.

Antibiotics per box.

Supportive care per [Respiratory Distress Guideline](#).

After 48-72 hours

Patient is improving and now meets outpatient treatment criteria.

No Yes

- Consult pediatrics.
- Consider repeating CXR and labs.
- Consider IVF.
- Consider broadening antibiotics.

Discharge with plan for total 7-10 days of antibiotics.

- Transfer to higher level of care.
- Supportive care per [Respiratory Distress Guideline](#).
- Consider [high flow nasal cannula](#).

Improving?

Yes

Continue care until meets criteria for outpatient management.

Antibiotics

Outpatient – total course of treatment 5 days

- 1st line: amoxicillin 45 mg/kg/dose PO BID
- 2nd line: Augmentin 45 mg/kg/dose PO BID
- 3rd line: cefdinir 7 mg/kg/dose PO BID

Inpatient – total course of treatment 7-10 days

- 1st line: ampicillin 50 mg/kg/dose IV Q6h
- 2nd line: Unasyn 50 mg/kg/dose IV Q6h
- 3rd line: ceftriaxone 50 mg/kg/dose IV Q24h

If not fully immunized, suspicion for *H influenzae*, or complicated pneumonia (pleural effusion, multilobar involvement, concern for bacteremia, etc.): Ceftriaxone until improving.

For *H influenzae* type A: At least one dose of ceftriaxone or four days of rifampin is necessary for decolonization. Remainder of course may be completed with a penicillin, if sensitive.

For PCN allergy: If reaction was non-anaphylactic, may trial amoxicillin with monitoring. If reaction was anaphylaxis, treat with a cephalosporin. If any questions, please obtain a pediatrics consult.

Azithromycin: Do not prescribe azithromycin unless there is evidence of an atypical pathogen and child is >5 years. Must be prescribed in addition to primary treatment above.

RUL infiltrate: Consider starting with Augmentin/Unasyn to cover for oral anaerobes. Consider thickener.

For Chronic Cough: See [Bronchiectasis/Chronic Cough guideline](#).

Follow-up for All Patients

- Within 48-72 hours.
- Do not repeat CXR unless recurrent infiltrate in same lobe; in that case, repeat CXR in 4-6 weeks.

REMEMBER:


- If patient is <90 days and febrile, please see [fever guidelines](#).
- Pneumonia is a clinical diagnosis and does not require X-ray findings.
- Place PPD if older than 6 months and no PPD in past 6 months.
- Any child <5 years with suspected pneumonia should be evaluated in Bethel or an SRC.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/11/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact
Amy_Carson-Strnad@ykhc.org.





Please see the [ANMC Pediatric Acute Hematogenous Septic Arthritis/Osteomyelitis Guideline](#).

- Please note: this guideline was designed at ANMC, where recommended labs, MRI, and operative management are immediately available and antibiotics can be started after these interventions.
- When evaluating a patient at YKHC with possible septic arthritis or osteomyelitis, strongly consider empiric antibiotics if there is going to be a delay of >6 hours to perform the recommended work-up (joint aspiration, surgical drainage, etc.), as noted in ANMC's guideline.
- Always discuss antibiotics with ANMC consultants and advocate for empiric usage if appropriate. Keep in mind possible delays, including weather, transport difficulties, and other emergencies. If deferring antibiotics, ensure that patient is closely monitored for development of worsening infection.
- Always feel free to consult YKHC pediatric hospitalist with any questions.



Universal Screening Recommendations

- All sexually active patients starting at age 14: annual screening for GC/CT, HIV, and syphilis.
- Any time GC and CT are tested for, HIV and syphilis screening should also be performed if not done in the last 12 months.
- Regardless of sexual activity, all teenagers should be screened for HIV by the age of 18. Additionally, all teenagers should undergo yearly GC/CT screening with, at minimum, a dirty urine.

Symptoms of Genital Infection

- Sores (genital, oral, or anal)
- Discharge or burning
- Dysuria
- Groin pain
- Pelvic pain
- Sore throat
- Rectal itching
- Discomfort or pain with bowel movement
- Vaginal itching or odor
- Testicular pain, swelling, or twisted feeling (can be off and on)
- Pain with ejaculation or sex


Age of Consent in the State of Alaska


Two people who are both 16 or older can legally agree to have sex with each other. When a person involved in sex is under the age of 16, Alaska law looks at the difference in ages to decide whether consent can be legally given.

- No person over 16 can legally have sex with someone who is 13 or younger.
- No person under 16 can legally have sex with someone who is 4 or more years older.
- No person under 16 can legally have sex with a person in a position of authority over them (including a teacher, coach, or minister).

A positive STI test in a patient who fits the above scenarios should be reported to OCS, law enforcement (BPD if in Bethel or AST if in a village), and the Child Abuse Pool in RAVEN.

Please note: There is no lower age limit for STI testing. Any patient may be tested, regardless of age, without special consent.





Expedited Partner Therapy (EPT)



EPT is a method of treating partners by asking the patient to take the doses to the partner. This is the standard of care for chlamydia at YKHC.

Process:

1. Treat the patient with azithromycin 1000 mg PO UNLESS there is concern for anal chlamydia, in which case the treatment of course is doxycycline 100 mg PO BID x7 days.
2. For azithromycin, give the patient pre-packaged doses for each sexual contact in the last three months. Give a handout explaining the process. This can be found under Patient Education → All → "EPT Partner Chlamydia (Custom)."
3. Encourage all contacts to be seen for full STI testing.

Retesting for All STIs

- If positive for anal and/or oral infection: Retest in one month at affected site (anal and/or oral) plus urine/vaginal.
- If pregnancy: Retest in one month.
- For all other cases: Retest in three months.





Mycoplasma Genitalium

- This organism has been recently identified in our population.
- Prevalence is estimated at 1% in the general population but is much higher in individuals at high risk of STIs.
- Mgen is a known cause of nongonococcal urethritis in males, cervicitis in females, and possibly pelvic inflammatory disease (PID) in females. It is uncertain whether it causes proctitis in men who have sex with men (MSM). There is no clear evidence that M. genitalium is associated with any human diseases outside the anogenital tract. (Source: UpToDate)
- Testing should be considered in patients who have persistent symptoms despite completing an empiric course of treatment.

Female patient with either:

- Persistent or recurring cervicitis
- PID

OR

Male patient with either:

- Non-gonococcal urethritis
- Persistent urethritis

Ensure full STI testing has been performed (HIV, syphilis, GC/CT/TV).
Consider retesting for GC/CT.
Treat per [guideline](#).

Test for *Mycoplasma genitalium* (Mgen) using these orders:

- Mycoplasma genitalium, NAA swab
- Mycoplasma genitalium, NAA urine

Collection tubes are:

Aptima: yellow for urine, orange for vaginal swab, blue/purple for endocervical swab.

Tests are run at the Alaska state public health lab and will take up to a week to result.

If NAA positive for Mgen:

Doxycycline 100 mg oral BID for 7 days
FOLLOWED BY
Moxifloxacin 400 mg oral daily for 7 days.

No test of cure or test for reinfection is necessary.
No contact tracing is necessary.
This is NOT reportable to the state.



If considering the diagnosis of bacterial sinusitis in a child younger than 4, please consult a pediatrician.

Fever and rhinorrhea in child >4 years old

90-98% of pediatric sinus infections are caused by viruses. Consider bacterial sinusitis only in the following scenarios.

Persistent Illness
Nasal discharge and daytime cough for >10 days
with no improvement

Observe for 3 days. Follow-up by phone or by appointment.

If no improvement

Worsening Course
One week of worsening nasal discharge, daytime cough, and fever after initial improvement

Severe Onset
Fever >102°F and purulent nasal discharge for >3 consecutive days

Treatment

1st line High-dose Augmentin 45 mg/kg/dose PO BID for 10-14 days
2nd line Cefdinir 14 mg/kg/dose PO daily for 10-14 days
Try to avoid using cephalosporins. They are less effective at treating the most common organisms that cause sinusitis.
For PCN allergy: Please obtain a pediatrics consult. Do not prescribe azithromycin or Septra. The most common pathogens in pediatric sinusitis have high resistance rates to these antibiotics. Avoid fluoroquinolones.

Follow-up by phone or by appointment at 3 days. If no improvement, consider broadening to next line of treatment.

Follow-up 10-14 days after starting treatment. If still symptomatic, consider ENT consult.

Differential Diagnosis

- foreign body
- seasonal/environmental allergies
- recurrent/back-to-back viral rhinitis or nasopharyngitis
- GERD

Sinus Development in Children



- Maxillary: present at birth, fully developed at 12 years
- Ethmoid: present at birth, fully developed at 12 years
- Frontal: present at 3 years, fully developed at 18-20 years
- Sphenoid: present at 8 years, fully developed at 12-15 years

Source: https://www.google.com/search?biw=1536&bih=740&btm=isch&a=1&ei=THW3rCHRJA4hKaICA&q=sinus+development&oeq=sinus+development&gs_l=psy-ab.3..0i67k1j0i5j0i5i30k113j0i8i30k13424034657.0352992.2.0.0.0.200.398.0tj1.2.0...1c.164psy-ab..0.2.395..030k1.0.rSFh1aHlq74#imgres=ID5BxCfQakWM:&sp=1518811500752

Imaging

Do not routinely obtain imaging studies in suspected sinusitis unless there is concern for a complication like orbital or CNS involvement.

Do not treat sinusitis, in the absence of symptoms, if it is an incidental finding on an imaging study.

Adjuvant Therapies


- Saline nasal spray
- Steam
- Oral hydration
- Tylenol and ibuprofen
- Do not routinely give decongestants and antihistamines (especially Benadryl). They have been proven ineffective in children and are unsafe under 6 years old.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 1/19/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact
Leslie_Herrmann@ykhc.org.




This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/24/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact clinical_guidelines@ykhc.org.






If you diagnose active TB, please contact a TB Officer.

DO NOT PUT A PATIENT WHO MAY HAVE ACTIVE TB ON A PLANE UNLESS ACUTELY ILL: This could expose the other passengers. Perform evaluation in village as able. Consult TB Officer.

TB Symptoms and Risk Factors (clinical judgment required)

- Hemoptysis
- Cough > 3 weeks
- Fever
- Night sweats
- Weight loss
- Persistent pneumonia
- Atypical CXR
- Household contact of active TB
- Prior active or latent TB infection
- Foreign born from endemic area
- Immunosuppression (HIV, diabetes mellitus, prednisone >15 mg/day for > 1 month, or TNF-alpha blocker)
- Born before 1960 and long-term resident of western Alaska or other endemic area



Abbreviations

AFB: acid-fast bacilli
 DOT: directly observed therapy
 EMB: ethambutol
 IGRA: interferon gamma release assay, e.g. QuantiFERON Gold
 INH: isoniazid
 LTBI: latent TB infection
 MOX: moxifloxacin
 MTB-RIF: mycobacterium tuberculosis nucleic acid amplification test that also tests for rifamycin resistance
 PZA: pyrazinamide
 RIF: rifampin(a rifamycin)
 RPT: rifapentine (another rifamycin)
 TST: tuberculosis skin test

TB Discharge and Follow up

- For hospitalized patients, use the [Heartland Criteria](#) to determine suitability for discharge
- For patients listed as having completed TB treatment, an appropriate initial follow up is a symptom screen and chest x-ray. However, there can be much nuance to follow up so consult TB officer if questions.

Contact Information

How to Consult a TB Officer: Send a message via Tiger Connect to "TB Officers" Team.

- Public Health Nursing (PHN):
 Phone: 907-543-2110
 Fax: 907-543-0435
 All directly-observed therapy (DOT) will be arranged by PHN.
- Curry Center TB Warm Line: (877) 390-6682
- Dr. Jacob Gray, ANMC Infectious Disease (Tiger Text)
- State Epidemiology: (907) 269-8000

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
 Approved 8/2/24. Click [here](#) to see the supplemental resources for this guideline.
 If comments about this guideline, please contact Robert_Tyree@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline

Tuberculosis, Latent (≥14 years)

Symptoms

- Cough for more than three weeks
- Weight loss
- Fever
- Night sweats
- Hemoptysis

Do not perform TB skin test or QuantiFERON Gold on anyone with a prior positive. QuantiFERON Golds can be ordered Monday through Thursday only, and they cannot be done in villages.

What is a positive TB skin test?

- At least 10 mm of induration OR >5 mm of induration for patients who are high risk for TB. (See box.)
- Must be read 48-72 hours after placement to be a true negative.
- If positive, the induration can remain up to seven days and can be read until then.

High Risk for Tuberculosis

1. Immunosuppressed, HIV positive, prednisone >15 mg/day for >1 month, TNF-α blocker.
2. Suspicious chest X-ray.
3. Household contact with active TB.

LTBI Treatments: Choose one option.
DOT is optional for all three treatment options.

1. 3HP: INH 15 mg/kg PO weekly, rounding to nearest 50 mg (max dose 900 mg) x 12 weeks AND Rifapentine PO weekly x12 weeks.
Rifapentine Dosing:
 - 32.1-49.9 kg: 750 mg
 - >50 kg: 900 mg (max dose)
2. Rifampin 10 mg/kg PO daily (max dose 600 mg) x4 months.
3. INH 5 mg/kg PO daily (max dose 300 mg) x9 months.
 - If on INH, give pyridoxine (vitamin B6) 50 mg PO daily to prevent neuropathy.
 - If patient is pregnant or HIV infected, the preferred treatment is INH for 9 months. In HIV infection, avoid rifampin and rifapentine.

Abbreviations

3HP: three month regimen of INH and rifapentine
AFB: acid-fast bacilli
DOT: directly-observed therapy
INH: isoniazid
LTBI: latent tuberculosis infection
MTB-RIF: mycobacterium tuberculosis nucleic acid amplification test that also tests for rifampin resistance
PHN: Public Health Nursing
TNF-α: tumor necrosis factor alpha

Patient ≥14 years with:
New positive TB skin test
OR
New positive QuantiFERON-Gold.

At least one symptom

No

Thirty minute appointment in Bethel for:
• Physical exam
• Chest X-ray
• Labs: LFTs, HIV, and hCG if female

Abnormal chest X-ray?

No

LTBI

- Call PHN with plan of care.
- Begin treatment per box, using LTBI Power Plan.
- Send LTBI prescriptions to the YKHC pharmacy and securely email notification to LTBI_Case_Managers@ykhc.org.

Yes

Patient in village?

No

Yes

If patient has symptoms concerning for TB, see [Active TB Guideline](#). Do not send patient to Bethel unless patient is medically unstable.

DO NOT PUT A PATIENT WHO MAY HAVE ACTIVE TB ON A PLANE UNLESS ACUTELY ILL; this could expose the other passengers. Perform evaluation in village, as able.

ACTIVE TB IS SUSPECTED

- Patient must wear surgical mask AND stay in a negative pressure room, if available, until MTB-RIF result is negative.
- Collect sputum samples using the "AMB NEW +PPD/ LTBI" Power Plan. This generates orders for all three sputum samples, including the 2nd and 3rd day samples.
- First, if 18 or older, collect one sputum sample (3 mL in a urine cup) for MTB-RIF and send to lab.
- Next collect three sputum samples for AFB smear and TB culture (5 mL in a conical tube) at least eight hours apart.
- Obtain labwork: LFTs, HIV, and hCG if female.
- Chest X-ray if available.

MTB-RIF positive?

Yes

Active TB.
Contact TB officer and see [Active TB Guideline](#).

No

- Consult TB officer regarding whether to treat for active TB
- Consider treating for [CAP](#).

Sputum positive? No response to CAP tx?

No

LTBI.

Continue full course of treatment per LTBI case managers.

Yes

Contact Information

How to Consult a TB Officer: Send a message via Tiger Connect to "TB Officers" Team.

- Public Health Nursing (PHN):
Phone: 907-543-2110
Fax: 907-543-0435
All directly-observed therapy (DOT) will be arranged by LTBI Case Managers.
- Curry Center TB Warm Line: (877) 390-6682
- Dr. Jacob Gray, ANMC Infectious Disease (Tiger Text)
- State Epidemiology: (907) 269-8000


This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved 8/2/24. Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact Robert_Tyree@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline

Tuberculosis Evaluation & Treatment (<14 years)



This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved 8/2/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Mien_Chyi@ykhc.org.



This protocol has been designed to maximize efficacy, use the least invasive measures that are still effective, and minimize hospital length of stay. **Please follow these steps to optimize sample quality.**

- ☐ 1. **Premedicate** with albuterol 2.5 mg/3mL (0.083%) solution – 3 mL via nebulizer to induce bronchodilation, facilitate delivery of hypertonic saline, and help prevent bronchospasm during delivery of hypertonic saline. May substitute MDI with mask and spacer. **DO NOT COMBINE with hypertonic saline.**
- ☐ 2. Administer 5 mL of 3% hypertonic saline solution via nebulizer **over a period of at least 10 minutes.** Prolonged administration has been shown to yield better samples.
- ☐ 3. If patient has copious nasal secretions, consider nasal suction with olive tip.
- ☐ 4. Obtain sample using mucus specimen trap with suction catheter appropriate for patient size. Measure from tip of nose to the tragus to cricoid cartilage for depth of catheter insertion and obtain sample via suction of the nasopharynx. The goal is to induce a gag and then a cough. *Sample is expected to be blood-tinged.*

(Note: This process may induce a vagal response. The patient should be sitting up with feet supported or lying down, NOT standing. If vasovagal syncope does occur, immediately place the patient supine with the legs elevated.)

- ☐ 5. Place specimen in appropriate collection container for desired test.
 - a. For rule-out pulmonary tuberculosis:
 - i. Collect three induced sputum samples **at least 8 hours apart** – one must be first morning sample (fasting goal 6-8 hours). Send for Acid Fast Bacilli Smear and Culture. Sample must be in an AFB container (conical with orange top), with a minimum volume of 2 mL (although 5 mL is preferable); sterile water may not be added to dilute sample.
 - ii. Two samples should also be sent for Xpert MTB-RIF. This test requires 3-5 mL of mucous in a sterile specimen cup. **DO NOT DILUTE**, or "saline wash" nares during suction for this specimen.
 - iii. AFB and Xpert may be obtained at the same time; if quantity not sufficient for both tests, prioritize the AFB.
 - b. Standard sputum cultures do not have a minimum volume and can be placed in a sterile specimen cup.
- ☐ 6. Label with full name of collector and date and time of the collection. This should be written **below the barcode**, NOT beside it. **If not labelled correctly, state lab will reject specimen.**
- ☐ 7. Collect specimen in RAVEN. Confirm the correct accession number and deselect any additional (future) accession numbers. **Ensure the collector ID, date, and time entered into RAVEN are an exact match to the written label.**

Contraindications to collecting an induced sputum: oxygen saturation of <92% despite supplemental oxygen therapy, inability to protect the airway, severe bronchospasm, or designation as inappropriate by the clinician for another reason (eg., midface trauma). After exclusion or resolution of these conditions, sputum induction can be considered.

Special considerations:

This procedure can also be used for patients who are able to follow instructions but do not have a productive cough. In these cases, suction may or may not be necessary.

While there are no contraindications due to age, for infants younger than 6 months, the sensitivity of induced sputum samples is lower than that of gastric aspirates. Thus, three first morning gastric aspirates collected 24 hours apart or a single first morning gastric aspirate followed by 2-3 induced sputum samples eight hours apart may be preferable. Please consult a pediatric TB officer to discuss this plan.

NOTE: Gastric aspirate samples cannot be sent for sputum culture or Xpert MTB-RIF.


Young infants with CPT1A-AV may need dextrose-containing mIVF while NPO. Very young infants may not tolerate fasting intervals of 6-8 hours; consider allowing breastmilk up to 4 hours pre-procedure and/or clear liquids up to 2 hours pre-procedure.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline

UTI (Adult)



This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 12/12/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Kaia_Pearson@ykhc.org.



Risk Factors for UTI

- Constipation
- Poor hygiene
- Uncircumcised boy

High Concern for UTI

- Fever >102
- No source
- Fever >48 hours
- History of UTI

May use [UTI Risk Calculator](#) to help risk-stratify.

Signs and Symptoms of UTI

- Fever
- Dysuria
- Hematuria
- Vomiting
- Abdominal pain
- New daytime or nighttime wetting
- Increased frequency of voiding

Differential Dx for Dysuria

- Vulvovaginitis
- Candida infection
- Bowel-bladder dysfunction
- Poor hygiene
- Sexual abuse (consider collecting dirty urine for GC/CT; see [Suspected Pediatric Sexual Abuse Procedure Guideline](#) for more information)
- Age-appropriate self-exploration
- UTI

Resistance

- Empiric drug choice is based on local resistance patterns (see [YKHC Antibigram](#)) and consultation with ID specialist.
- If urine culture grows an Extended-Spectrum Beta-Lactamase (ESBL) producing organism, please obtain an infectious disease consult during business hours and add ESBL to Problem List.

Indications for VCUG

- Recurrent febrile UTI.
- Major anomaly on ultrasound. Consult pediatric urologist and consider obtaining VCUG in Anchorage.

Note: study available in Bethel 1-2 times per year when radiologist in-house.

Child less than 5 years of age with concern for UTI

Obtain catheterized or true clean catch urine sample for urinalysis (UA) AND culture.

UA positive for one or more:

- Leukocyte esterase
- Nitrites
- >5 WBC/HPF on microscopy

Ensure urine culture is sent.

Patient appearing toxic or not tolerating PO?

Yes

Empiric antibiotic treatment:
ceftriaxone
50 mg/kg/day
IV/IM

No

Consider empiric treatment with cephalexin 20 mg/kg/dose Q8h x5-10 days. Max dose 1000 mg. Note: consider 10 day course if <6 mo. May defer empiric treatment and await culture sensitivities.

Urine Culture Results

Diagnose UTI if:

- Single species ≥50,000 CFU/mL in a catheterized sample
- Single species ≥100,000 CFU/mL in a clean catch sample.
- Single species ≥10,000 CFU/mL in a catheterized sample with positive UA.
- ≥50,000 CFU/mL in a clean catch sample with positive UA.

Yes

Begin empiric treatment and narrow coverage per sensitivities.

If first UTI and ≤24 months, perform renal ultrasound at first opportunity to evaluate anatomy.

Renal abnormality identified.

- If mild pelviectasis identified, repeat ultrasound in 3 months.
- If any other anomaly identified, consult pediatrics.

Symptomatic Care

- If dysuria, irritation, etc. recommend A+D ointment and instruct family to do soaks/baths with warm water and no soap.
- May consider baking soda ¼ cup per tub.

Village Management

- DO NOT treat any child empirically in the village without sending clean catch urine sample for culture.
- Consider symptomatic care (see box) for possible vulvovaginitis.
- If patient has dysuria, increased frequency, new-onset enuresis, and/or abnormal clean catch urinalysis, consider further evaluation in Bethel.

DO NOT...

- Treat any child for UTI empirically without culture pending.
- Routinely collect urine via bag.
- Routinely perform a test of cure.
- Routinely start UTI prophylaxis.
- Perform suprapubic taps.
- Routinely obtain bloodwork for uncomplicated UTI.
- Add UTI to Problem List until confirmed by culture.

No UTI.
Stop antibiotics.

NOTE: For infants <90 days with fever, see the [Fever in Infants 0-90 Days Guideline](#).

If diagnosis of UTI is suspected after appropriate work-up, follow this guideline for full management recommendations.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 1/19/24.
Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact Clinical_Guidelines@ykhc.org.



True Varicella infection is RARE in our region:

- DO NOT diagnose Varicella** without confirmatory lab testing.
- Per the CDC:
 - Two doses of VZV vaccine are 88-98% effective at preventing all VZV infections.
 - One dose of VZV vaccine is 80-85% effective at preventing all VZV infections.
- All confirmed Varicella must be confirmed to State Epidemiology with this form: <http://dhss.alaska.gov/dph/Epi/Documents/pubs/conditions/frmlnfect.pdf>

Differential Diagnosis

- Hand-foot-mouth disease
- Scabies
- Stomatitis
- Eczema herpeticum
- Diffuse impetigo

Documentation for Suspected Varicella Infection

- Date of symptom onset
- Date of suspected diagnosis
- Date of rash onset
- Location of rash, including where first noted
- Number of lesions
- Photos of lesions
- Evolution of rash (including appearance of new groups of lesions)
- Appearance of lesions (are there lesions in all stages of development at once?)


High Risk Exposures

- Inquire if any pregnant women or immunocompromised people have been exposed.
- For pregnant women: find out if she has a history of varicella or has received the vaccine. If not, then consult HROB to consider further treatment.
- For immunocompromised patients: refer to a provider for evaluation.

Note: Unconfirmed cases must still receive the vaccine. There is no increased harm in vaccination if the patient had varicella in the past.

Typical Presentation for Chickenpox/Varicella


- 10-21 days after exposure, fever appears, followed by rash that often starts on the head and then moves down.
- Rash appears in successive crops over several days.
- Rash begins as macular and then progresses to vesicular, then pustular, then crusted.
- There are lesions in different stages of development on different parts of the body.**
- New vesicles stop forming within four days.



This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 1/19/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Lestlie_Herrmann@ykhc.org.



Checklists for Specific Types of Admissions

- **TB evaluation** (see [guideline](#)):
 - Work-up: CXR, LFTs, HIV test
 - Contact TB officers via Tiger Connect and ensure they are aware of admission and have no further recommendations.
 - Discuss expected course with caregivers, including induced sputum collection and expected stay of ~24 hours.
- **Iron Infusion** (see [guideline](#)):
 - If first infusion, ensure work-up (CBC, iron panel, lead level) is complete and within past month.
 - Discuss IV access with admitting provider and accepting nursing team. Where IV access is obtained should be a team decision.
 - Discuss expected course with caregivers.
- **Skin Care** (severe eczema exacerbation, etc.):
 - If patient with history of chronic problems, work-up and IV are often not required. Discuss with accepting physician.
 - Discuss expected course with caregivers, including to expect to stay at least 3-5 days.
- **Failure to Thrive** (see [guideline](#)):
 - Discuss appropriateness of work-up and IV access with accepting physician. These decisions are tailored to each patient's unique presentation and needs.
 - Discuss expected course with caregivers, including to expect to stay at least 3-5 days.
- **Constipation/Clean-out**:
 - Discuss appropriateness of work-up and IV access with accepting physician. These decisions are tailored to each patient's unique presentation and needs.
 - Discuss expected course with caregivers, including to expect to stay at least two days.
- **Respiratory** (see [pneumonia guideline](#), see [respiratory distress guideline](#)):
 - Discuss appropriateness of work-up and IV access with accepting physician. These decisions are tailored to each patient's unique presentation and needs.
- **Skin and Soft Tissue Infections** (see [guideline](#)):
 - Discuss IV access with admitting provider and accepting nursing team. Where IV access is obtained should be a team decision.
 - Discuss expected course with caregivers.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 9/16/24.
Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact
Leslie_Herrmann@ykhc.org.



IF ANY CONCERN FOR APNEA, please consult a pediatrician immediately to determine need for further evaluation, transfer, medevac, etc.

Recommendations on Management of Caffeine After NICU Discharge

- Recommended dose of caffeine is 12 mg/kg PO daily.
- Patient should be seen in Bethel by a pediatric provider within one week of returning to the region.
- Dose should be weight-adjusted every 1-2 weeks. This can occur in outpatient clinic with a pediatric provider or a pediatric consult, in an SRC with a pediatric consult, or in a village by RMT to Chronic Peds.
- Stop the caffeine when the baby is 42 weeks corrected gestational age.
- Discontinuation of caffeine may be delayed for another week so as not to coincide with immunizations, recent URI, or planned anesthesia (as all of these events can cause re-emergence of intermittent hypoxia with periodic breathing).



When a Baby is Discharged from the NICU on Caffeine

- Update the Problem List with the plan, including the target dose, how often to weight-adjust, and the expected end date (when 42 weeks corrected gestational age will be).
- Write a prescription for the caffeine. Include the target dose. Under "eRx Note to Pharmacy," state "do not fill until family calls for refills."
- Assess caffeine dose at every encounter.

Rationale

- In the past, premature infants were given caffeine until about 34 weeks post-menstrual age. Some needed caffeine past this point and went home on caffeine and an apnea monitor.
- Recent studies have shown that many preterm infants who have been taken off caffeine will go on to have intermittent hypoxia and subclinical apnea and bradycardia events after discharge from the hospital.
- Evidence is also building that prolonged use of caffeine results in better neurodevelopmental outcomes.
- As of January 2019, caffeine has been continued in preterm infants after discharge from the PAMC NICU.
- The PAMC NICU stopped the routine use of apnea monitors for babies discharged on caffeine due to sub-optimal monitor technology and frequent frustration among parents and providers. They prefer to emphasize the importance of giving caffeine rather than use of apnea monitors.

Source

Adapted from letter from Alaska Neonatology Associates, Inc., Pediatrix Medical Group, an affiliate of MEDNAX.
1/10/2019
Providence Alaska Medical Center (PAMC)
Neonatal Intensive Care Unit (NICU)

This resource is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 1/11/23.
If comments about this resource, please contact Leslie_Herrmann@ykhc.org.



Criteria

- Weight for age <5th percentile on WHO Growth Chart (0-24 months). (Note: The growth chart in RAVEN defaults to the CDC. Select the WHO growth chart by clicking on "CDCWHO.")
- Weight for length <5th percentile on WHO Growth Chart (0-24 months).
- Decrease across two major percentile lines over a 3-6 month period.

High-Risk Criteria (Consider admission.)

- <1 month: not regaining birth weight by 21 days of life, continued weight loss after 7 days of life.
- 1 month-12 months: no weight gain, weight loss, weight for length z-score of -2 or lower. (Note: to see z-score in RAVEN, select the weight for length growth chart and click "table.")
- Medical instability
- Moderate to severe malnutrition with concern for refeeding syndrome
- Moderate to severe dehydration
- Failed outpatient managements including multiple missed appointments
- Suspected abuse/neglect

Frequency of Weight Checks

- <1 month: Q1-3 days
- 1-6 months: Q1-2 weeks
- 6-12 months: Q2-4 weeks
- 12-24 months: Q2-4 weeks

Patient at risk for malnutrition or failure to thrive (FTT)

Initial Evaluation

- Full history (see box on [next page](#)). Use autotext "...pedFTThpi."
- Full physical exam (see box on [next page](#)). Use autotext "...pedFTTpe."

Please have a low threshold to consult a pediatrician for any child you are concerned about.

Does patient have any High-Risk Criteria? (See box.)

Yes

No

- Consider differential diagnosis.
- Perform lab work-up (see box).
- Consider admission.

- Give feeding recommendations per box on [next page](#).
- Schedule weight checks at frequency in box.
- Consider fortification of feeds.

- Daily naked weights.
- Strict I/O, calorie counts.
- If breastfeeding, do weights pre- and post-feeds. (Naked except for diaper that should not be changed until after weight.)
- Implement initial feeding recommendations per box on [next page](#).

At weight check, is patient gaining adequate weight?

Yes

Continue scheduled weight checks until adequate weight gain at three consecutive visits.

Is patient gaining adequate weight after 2-3 days?

No

Yes

- Implement secondary feeding recommendations on [next page](#).
- Consider transfer to higher level of care if patient does not gain weight after these measures.

- Discharge with:
- Detailed feeding plan that includes timing, volume, calorie density, supplements, etc.
 - Scheduled follow-up per box. If returning to village, consider a weight check in 24-48 hours in outpatient clinic before returning to village.

Differential Diagnosis: General Categories and Symptoms

Inadequate Intake

- Long intervals between feeds (Sleep >3 hours if <2 months old)
- Falling asleep during feeds
- Limited number and volume of feeding per day
- Improper mixing of formula
- Lactation problems: poor supply, difficulty with latching
- Limited urine diapers (<1 wet diaper per 8 hours)
- Food insecurity/inability
- Excessive vomiting/spitting up/reflux
- Increased hunger cues/caregiver isn't recognizing cues
- Symptoms of maternal depression
- Birth weight not regained in 2 weeks
- Oral Motor Dysfunction

Malabsorption

- High volume, extremely loose stools
- Clay-colored stools
- Greasy or significantly foul-smelling stools
- Chronic diarrhea
- Abdominal distention, gassiness with diarrhea
- Blood in stools

Increased Metabolic Demand

- Cardiac: heart murmur, tachypnea, sweating or cyanosis with feeds, feeding fatigue
- Respiratory: noisy breathing, tachypnea, difficulty breathing with feeds, nasal obstruction
- Neurologic: increased or decreased tone, abnormal movements
- Metabolic/genetic: abnormal newborn screen, dysmorphic features
- Renal: urologic abnormalities, renal tubular acidosis
- Endocrinology: tachycardia, diaphoresis

Lab Workup, By Age

Use Power Plan "PED Pediatric Failure to Thrive" to place orders in RAVEN.

- <1 month:
 - Verify Newborn Screen, CMP, CBC, urinalysis.
 - Consider metabolic evaluation.
- 1-24 months:
 - CBC, CMP, urinalysis, TSH, HIV, PPD (if <6 months but only actionable if positive) or Quantiferon (if >6 months), tTG IgA and total IgA if > 6 months and gluten exposure.
 - Consider sending stool for occult blood, metabolic evaluation.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 8/2/24.

Click [here](#) for supplemental resources for this guideline.

If comments about this guideline, please contact
Jennifer_Prince3@ykhc.org.

History

Use autotext “..pedFTThpi” to document in RAVEN.

General

- Recurrent fevers or infections
- Detailed birth history

Cardiovascular

- Sweating and/or fatigue with feeds

GI

- Constipation
- Vomiting

Neurologic

- Depressed mental status, inconsolability, sleepiness
- Developmental delay
- Abnormal movements

Feeding

- Breastfeeding
 - Frequency, length, number per day, longest interval between feedings, night vs day?
 - One or both breasts, softer after feeding, ± nipple shield, any pain or difficulty with latch?
 - If pumping, how much is produced?
 - Can you see or hear the baby swallow?
 - Any supplementation (expressed breast milk or formula)?
 - Does baby fall asleep at breast?
- Formula
 - Frequency, length, amount per feed and per day, longest interval between feeds, night vs day?
 - Type of formula and recipe
 - Type and size of bottle and nipple
 - Any supplementation (either addition to the bottle or solids)?
- Swallow problems
 - Coughing during feeding
 - Wet or gurgly sounds during or immediately after feeding
 - Frequent upper respiratory tract infections, fevers, or pneumonia
- Reflux
 - Coughing, choking, gagging, or any respiratory symptoms with feeds
 - Spitting up/vomiting
 - Arching, fussiness, or discomfort with feeds

Social

- Who feeds the baby? Who lives at home? Is there a feeding schedule?
- If bottle fed, are there concerns about obtaining enough formula?

Elimination

- Number of wet and stool diapers per 24 hours
- Stool appearance (consistency, color, any orange/red crystal/powder, any blood or mucus)

Please see [ANMC's Preterm Nutrition Resource](#) for more information, including recipes for mixing high caloric density formula.

Initial Feeding RecommendationsBreastmilk/Formula

- Minimum Intake Recommendations:
 - Term Infant: 108 kcal/kg/day = 162 mL/kg/day of 20 kcal/oz formula/breast milk
 - Preterm Infant: 110-130 kcal/kg/day = 177 mL/kg/day of 22 kcal/oz preterm formula
- Feeding Frequency:
 - <3 months: Q3h or ≥8 feeds/day. No more than 3 hours between feeds.
 - ≥3 months: Q3h during day with ≥6 feeds/day
- Wake the baby to feed if necessary.

For Solids

- Infant must be taking at least 24 oz/day of formula or breastmilk.
- Limit any other fluids like water or juice.
- By 12 months, goal 4-6 servings of >4 tablespoons per day.

Secondary Feeding Recommendations

- If patient is able to tolerate goal feed volume, increase volume by 10% to max 180 mL/kg/day OR increase caloric density by 2 kcal/ounce to max 24 kcal/ounce.
- Allow at least 24 hours to assess tolerance to any changes.
- If patient is taking solids and >9 months, consider increasing calories in solids.
- If patient is not able to consistently and safely take enough by mouth to gain weight, consider NG feeds.

Physical

Use autotext “..pedFTTpe” to document in RAVEN.

General

- Cachexia, decreased subcutaneous stores, decreased muscle bulk
- Relative macrocephaly
- Lack of caregiver bonding or responsiveness to patient
- Dysmorphic features or syndromic appearance

HEENT

- Scleral icterus
- Nasal congestion or obstruction
- Cleft lip or palate
- Macroglossia or ankyloglossia
- Micrognathia

Respiratory

- Stridor
- Difficulty breathing, tachypnea
- Abnormal breath sounds including wheezing, crackles, etc.

Cardiovascular

- Murmurs
- Diminished or absent peripheral pulses

GI

- Hepatosplenomegaly
- Abdominal distension
- Palpable stools

Skin

- Jaundice
- Rashes or skin breakdown (including in diaper area)
- Severe atopic dermatitis

Neurologic

- Depressed mental status, inconsolability, sleepiness
- Developmental delay
- Abnormal movements

Caloric Needs by Age

If preterm, use corrected age.

- <37 weeks: 110 -130 kcal/kg/day
- 37 weeks-6 months: 108 kcal/kg/day
- 7-12 months: 98 kcal/kg/day
- 12-24 months: 75-95 kcal/kg/day

Average Daily Weight Gain by Age


Age (corrected)	Median (grams/day)	
	Girls	Boys
2-4 weeks	29	34
4 weeks-2 months	34	40
2-3 months	24	27
3-4 months	20	21
4-5 months	16	17
5-6 months	13	14
6-8 months	11	11
8-10 months	9	9
10-12 months	8	8
12-15 months	4-9.5	4.5-10
15-18 months	4-9.5	4-9
18-21 months	4-9.5	4-9
21-24 months	3.5-9	3.5-9

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 8/2/24.

Click [here](#) for supplemental resources for this guideline.

If comments about this guideline, please contact
Jennifer_Prince3@ykhc.org.



Barlow and Ortolani Tests

- The Barlow test is for laxity of the hip joint. It should be performed gently with no posterior force. If positive, you will feel laxity or the hip will sublux or dislocate.
- The Ortolani test is the maneuver to reduce a dislocated hip. If positive, you will feel a clunk.
- Per the AAP, "One can think of the Barlow and Ortolani tests as a continuous smooth gentle maneuver starting with the hip flexed and adducted, with gentle anterior pressure on the trochanter while the hip is abducted to feel whether the hip is locating into the socket, followed by gently adducting the hip and relieving the anterior pressure on the trochanter while sensing whether the hip slips out the back. The examiner should not attempt to forcefully dislocate the femoral head."
- See [this video](#) for AAP guidance on these exam maneuvers.

Swaddling

Swaddling techniques that restrict hip and leg movement have been associated with developmental dysplasia of the hip. Parents should be advised to avoid tight swaddling of the lower extremities

Orthopedics Consults & Referrals

1. Consultation:

- Beneficiary patients: contact ANMC orthopedic surgeon on call at (907) 563-2662 (*97) or send message through Tiger Connect.
- Non-beneficiary patients: contact Dan Brown, MD at Anchorage Fracture & Orthopedics at (907) 563-3145.

2. Referral:

- Place an order for "Refer to Orthopedics External" with brief history. Note the orthopedist who was consulted. Indicate where the referral should be sent.
- Send a RAVEN Communication to Chronic Peds Case Manager Pool about the referral and level of importance.

Imaging

Patient must have either ultrasound or X-ray, as below.

1. Hip ultrasound: 6 weeks to 4 months of age.

- Performed at ANMC for beneficiaries and Imaging Associates for non-beneficiaries.
- Beneficiary patients: Place order for "Refer to Pediatric Clinic External (MRI / EEG / VFSS / Hip US)" with brief history. Request follow-up appointment with SCF Team B
- Non-beneficiary patients: Place order for "Refer to Peds Other External," facility "Other – Imaging Associates" with brief history. Request follow-up appointment with pediatric provider in Bethel
- Send a RAVEN Communication to Chronic Peds Case Manager Pool about the referral and level of importance.

2. X-ray, AP pelvis: over 4 months of age. (Note: in premature infants, ossification of femoral heads is delayed. May use corrected gestational age of 4 months or later.)


- Performed at YKHC.
- Place an order for "XR Pelvis (Pelvis AP only)" and put in comments "AP view with hips in neutral position to rule-out developmental dysplasia of the hip."
- Send a RAVEN Communication to Chronic Peds Case Manager Pool stating the order was placed and requesting an appointment for this with a pediatric provider in Bethel.

This protocol is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee. 7/23/25.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this protocol, please contact Zoe_Storck@ykhc.org.



How to Use [PediTool Calculator](#) (Note: Website may not work in Firefox.)

1. Enter infant's gestational age, age in hours, and total bilirubin level.
2. Note if infant has any neurotoxicity risk factors. (See box.)
3. Click "submit."
4. The next page will plot the level on a graph. You will see if the infant meets criteria for phototherapy and/or exchange transfusion.
5. If not starting phototherapy, scroll down to the table labelled, "Post-birth hospitalization discharge follow-up for infants who have NOT received phototherapy." Follow these recommendations for repeat levels.

Phototherapy

1. Order using one of the following:
 - PED Pediatric Admission Power Plan → PED Phototherapy sub-phase
 - OB/Newborn orders folder → OB Newborn Phototherapy Power Plan
2. **Check hemoglobin and hematocrit on all patients receiving phototherapy. May obtain via heel stick when checking bilirubin.**
3. Check serum total bilirubin level Q12h (or more frequently if neurotoxicity risk factors or concern for ongoing hemolysis). If level is trending up, consult pediatrician and consider broadening differential and work-up. Note: Transcutaneous bilirubin is not reliable until 24 hours after phototherapy has been stopped.
4. Encourage frequent feeding, but try to limit time out of phototherapy to no more than 20 minutes Q3h. Use bili blanket for feeds.
5. IV fluids are unnecessary unless infant has signs of dehydration.
6. Keep infant supine with eye protection and diaper (for gonad protection) while under phototherapy.
7. Stop phototherapy when serum total bilirubin level is ≥ 2 mg/dL below the phototherapy initiation level, using the hour of life at which phototherapy was initiated.
8. Obtain rebound bilirubin level 6-12 hours after stopping phototherapy if patient required phototherapy in first 48 hours of life, <38 weeks, concern for hemolysis, or DAT positive.
9. For all babies who met the phototherapy threshold, check a serum bilirubin level the day after stopping phototherapy.

Concerning Scenarios

- If jaundice is visible before 24 hours of life, check level immediately.
- If infant is not well-appearing, consider sepsis, consult pediatric hospitalist, and follow recommendations on [fever <90 days guideline](#).
- If total bilirubin level meets PediTool's Escalation of Care criteria, the NICU should be consulted and Labs for Expanded Work-up collected.
- If direct bilirubin > 1 mg/dL, do not start phototherapy. Direct hyperbilirubinemia requires a work-up and often gastroenterology consultation. Obtain pediatrics consult.

Neurotoxicity Risk Factors

Use this list to answer question on [PediTool calculator](#).

- Isoimmune hemolytic disease (positive DAT, etc.), G6PD deficiency, or other hemolytic condition
- Sepsis
- Clinical instability in past 24 hours
- Low albumin (if known)

Direct Antibody Test (DAT)

- Order a DAT if:
 - Mother is type O or Rh negative and did not receive Rhlg during pregnancy.
 - The infant's total bilirubin level has a high rate of rise (0.3 mg/dL/hour in first 24 hours or 0.2 mg/dL/hour after first 24 hours).
- If infant has a positive DAT, check transcutaneous bilirubin immediately and then retest Q4h x2 then Q12h x3.

Labs for Expanded Work-up
Consider in infants with jaundice at <24 hours of life, rising levels despite phototherapy, or recurrent jaundice.

- Blood type, DAT (Direct Antibody Test, or Coombs)
- CBC with manual differential and reticulocyte count
- CMP
- Thyroid studies (if prolonged or recurrent)
- GGT
- G6PD

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/23/25

Click [here](#) to see the supplemental resources for this guideline.


If comments about this guideline, please contact Mien_Chyi@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline

Care of Late Preterm & Low Birth Weight Newborns



- Encourage mother to express breastmilk.
- If infant is stable, encourage bonding and breastfeeding while awaiting medevac.

NOTE: If infant of any GA is unstable at any time, please contact the pediatric hospitalist (Tiger Connect Peds Wards on Duty) and prepare for transfer.

Parent Education

- Educate parents regarding vulnerability of late preterm neonate and late preterm protocol.
- Attach completed Late Preterm Crib Card to crib.
- Ensure parents have received the Late Preterm Handout and use as a resource.
- Emphasize need for follow-up with outpatient appointment prior to return to village.
- Ensure and encourage proper pediatric follow-up.
- Education regarding feeding plan and follow-up resources.

Infant Stability

- Temperature ≥ 97.7 (axillary) for 6 hours in open crib.
- Cardiovascular and respiratory stability as determined by the medical team.
- Able to tolerate oral feeds without color change or increased WOB: breastfeeding or tolerating 5-10 ml EBM or formula at a minimum of every 3 hours.

Strikes

- Any temperature < 97.7
- Any weight < 2200 grams
- Any blood glucose level below target for age

****NOTE:** Term babies with BW < 2200 grams do not need to be automatically transferred if stable. For these infants, this guideline should be applied, with the BW counting as one strike. There should be a huddle at 24 hours of life or sooner if infant receives two more strikes.

- Admit patient to OB using the Late Preterm Power Plan.
- Infant is observed in the mother's room or in the Newborn Treatment Room for at least four hours to ensure stability.
- VS Q4h, including temperature, throughout entire stay.
- Weigh baby Qshift.
- Blood glucose screening per [protocol](#) for full first 24 hours of life.
- Establish feeding plan with parents (see box).
- Ensure parents are educated (see box).
- Follow Late Preterm Goals of Care worksheet (to be placed on baby's hard chart).
- On day of birth, schedule outpatient appointment for DOL 4-5 to ensure appointment availability.

Huddle at 24 hours of Life

(to include bedside nurse, charge nurse, family medicine hospitalist, and pediatric hospitalist if needed)

- Points to discuss: how the baby is feeding, %weight loss, can we safely manage the baby's needs, unit acuity/staffing ratios, does the baby need to be transferred at this time, time for next huddle (if needed).
- If infant receives three "strikes" on the Late Preterm Goals of Care worksheet, there must be a huddle to discuss if the infant should be transferred. (See Strike box.)

Goals for Discharge

- All late preterm babies are admitted for at least 72 hours.
- Weight loss $< 8\%$ below BW.
- Temperature $\geq 97.7^\circ\text{F}$ x24 hours in an open crib.
- Well-established feeding plan.
- Follow-up appointment scheduled in outpatient clinic in Bethel in 24-48 hours. If weekend, may have this follow-up on OB by pediatric hospitalist.
- Must have warm handoff with message sent to provider seeing patient for follow-up that includes minimal requirements to be met for discharge back to village.
- Follow-up weekly in village or outpatient clinic until corrected GA of 40 weeks.
- Prescribe Poly-Vi-Sol WITH Iron at discharge.

Definitions

- GA: gestational age at birth
- Late preterm: GA 34 weeks 0 days to 36 weeks 6 days
- Early term: GA 37 weeks 0 days to 38 weeks 6 days
- Term: GA 39 weeks 0 days to 40 weeks 6 days
- Low birth weight is any baby born < 2500 grams

Characteristics of Late Preterm Infants

- Low birth weight
- Low body fat
- Poor thermoregulation
- Low glycogen stores
- Low tone
- Poor state regulation
- Immature immune system
- Immature suck and swallow
- Delay in bilirubin metabolism

Late Preterm Infants Are at Risk For:

- Hypothermia
- Hypoglycemia
- Sepsis
- Poor feeding and infrequent feeds can lead to inadequate maternal milk supply/breast feeding failure
- Poor suck and swallow may lead to inadequate milk intake
- Excessive weight loss, failure to thrive
- Hyperbilirubinemia with late rise (expect peak on DOL 5)
- Increased readmission rate (5-13 times that of term infants)
- Respiratory instability in upright car safety seats or other upright infant devices
- Hospital readmission

Feeding Plan

Infants meeting any of the following criteria should be assessed for the need for supplementation:

- Birth weight < 2500 grams
- Poor reserve (evidenced by temperature instability or hypoglycemia)
- Poor feeding (LATCH < 7 or < 10 minutes at breast)
- Weight loss $> 3\%$ per day or $> 8\%$ total
- Minimum volumes for both bottlefed and breastfed babies:
 - 0-24 hours: 5-10 mL
 - 25-48 hours: 10-20 mL
 - 49-96 hours: 20-30 mL
- If bottlefeeding, advance feeds as tolerated.

If Breastfeeding

- Lactation evaluation within 24 hours of birth.
- **LATCH score** documented at least Qshift.
- Infant should be put to breast at least Q3h.
- Use Supplemental Nursing System (SNS) **to ensure measurable amounts each feed with the above minimum volumes.**

Supplementation

- Supplementation should be given by SNS (preferred), cup, or finger feeds rather than nipple and bottle. May receive formula if milk volume not meeting fluid needs.
- Mother to pump every 3 hours after nursing unless infant nursing vigorously.
- Bedside nurse and medical team should re-evaluate feeding plan daily.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/14/23.

Click [here](#) to see the supplemental resources for this guideline.


If comments about this guideline, please contact Amy Carson-Strnad at amy_carson-strnad@ykhc.org.



mPEWS (modified Pediatric Early Warning System)

- YKHC uses the mPEWS to monitor admitted pediatric patients. Scoring is required prior to and throughout admissions to screen patients for acuity and help determine appropriate disposition.
- This is a scoring system that can help identify patients at risk for deterioration. YKHC uses it as a communication tool to highlight these patients.
- The score is calculated using the Ad Hoc form called "mPEWS," found in three places: (1) in ER encounters, under "mPEWS;" (2) in inpatient encounters, under "Assessments" → "mPEWS;" and in other encounters in "Asmt/Tx/Monitoring" folder.

- The ED will calculate a score prior to admission to the inpatient unit using the most recent vital signs and several aspects of PMH. If PMH unclear, the provider of record will clarify.
- Upon admission, the Inpatient Unit will calculate a score using the first vital signs taken.





Indications for Use

- Neonate with respiratory distress.
- Head circumference ≤ 39 cm and age ≤ 2 months.
- Has stabilized on CPAP via NeoPuff.
- Must have a respiratory rate (NO apnea).
- Anticipated prolonged need for CPAP at YKHC due to weather, NNP team unavailable for transport, etc.
- Note: If newborn <26 weeks, discuss with NICU prior to use.

Note: If using on a non-newborn, must request NNP/ neonatal transport team to ensure level of support can be maintained during transport.

Mask/Bonnet Set-Up

1. Measure the head circumference with the tape provided to determine the correct bonnet size.
2. Place the bonnet on the baby's head. Make sure to cover the bottom of the ears and the back of neck.
3. Mount the mask on the tubing.
4. Attach the fixation pillow to the hat, secure the tubes in the grooves, and secure with Velcro.
5. Use the fixation straps to hold the mask in place.
6. To hold optimal pressure, try to keep the infant's mouth closed. May offer a pacifier with Sweet-Ease.



Ventilator Settings

1. While in standby, select the neonatal option. Then select "Modes."



2. Enter the patient's weight.

3. Select nCPAP mode and "Confirm."



4. Adjust PEEP and FiO_2 and then select "Confirm."

5. Select "Start Ventilation."

See the YKHC [Neonatal Resuscitation Summary](#) for weight-based drug doses and equipment.

This resource is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 2/9/24.

If comments about this resource, please contact Zoe_Storck@ykhc.org.




Tubing Set-Up



Hamilton ventilator

Inspiratory limb (white)
Expiratory limb (blue)
Pressure sensor (blue, small)

Humidifier (sterile saline bag attached)



Troubleshooting Tips

Ventilator Settings:

- Start by matching the PEEP level infant stabilized on with the NeoPuff, generally 5-8 cm H₂O.
- Adjust settings as needed. If FiO₂ <40%, consider weaning PEEP. If requiring >8 cm H₂O, discuss with neonatologist.

Tubing Set-Up:

- Try to maintain tubing proximal to infant parallel with the infant's face and midline.
- Ensure there is no tension on the tubing.

Bonnet:

- Choose the smallest bonnet you can fit on the baby. A snug fit will hold everything in place.
- Ensure velcro is not rubbing against the baby's skin.

Mask:

- Avoid overtightening but ensure good seal.
- If requiring nasal CPAP >6 hours, watch closely for skin breakdown and consider alternating mask with prong interface.





Signs of Neonatal Sepsis

- Temp ≥ 100.4 or ≤ 97.5
- Irritability
- Poor Feeding
- Hypoglycemia
- Hypothermia
- Tachypnea
- Tachycardia
- Grunting
- Hypoxia
- Lethargy
- "Not acting right"

If any of these signs are present, consider obtaining a pediatrics consult.

Is infant ≥ 35 weeks?

No

This guideline does not apply to infants <35 weeks. Consult pediatrics. See [this link](#) for resources.

Yes

Is one or more of the following present?

- Any maternal fever during intrapartum period
- Rupture of membranes ≥ 18 hours prior to delivery
- GBS unknown or GBS positive by culture or PCR

No

- Routine care.
- Consult pediatric hospitalist with any questions or concerns.

Yes

- Go to the [Kaiser Neonatal Early-Onset Sepsis Calculator](#).
- Enter infant and maternal information.
- For local incidence of newborn sepsis, choose CDC national incidence.

Note: For the purposes of this calculator, only penicillin, ampicillin, and cefazolin are considered GBS-specific antibiotics. If the mother received clindamycin or vancomycin alone, enter "no antibiotics."

- Follow recommendations for blood culture, antibiotics, and vital sign frequency based on early-onset sepsis risk for infant's clinical status.
- "Well-appearing," "equivocal," and "clinical illness" are defined [here](#).
- Note: If the calculator recommends observation, this should be for at least 48 hours with vital signs Q4h.
- If maternal GBS positive and inadequately treated, observe infant for at least 48 hours regardless of calculator recommendations. If GBS positive and adequately treated, may discharge when clinically appropriate.

If any clinical concerns about infant, consult pediatric hospitalist

If giving antibiotics:

- Send blood culture, CBC with differential, and CRP.
- Order ampicillin and gentamicin, using [Neofax](#) or the [Neonatal Resuscitation Summary](#) for dosing.
- Consult pediatric hospitalist and prepare to transfer infant to NICU.

Note: CBC and CRP are not routinely recommended anymore. However, they can be useful to trend if starting antibiotics. Thus, we recommend they be obtained if starting antibiotics.

References

- *Pediatrics* 2019: [Management of Infants at Risk for Group B Streptococcal Disease](#)
- *Pediatrics* 2018: [Management of Neonates Born at \$\geq 35 0/7\$ Weeks' Gestation with Suspected or Proven Early-Onset Bacterial Sepsis](#)

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner. Click [here](#) to see the supplemental resources for this guideline.

Approved by Clinical Guideline Committee 7/23/25.

If comments about this guideline, please contact Amy_Carson-Strnad@ykhc.org.



Target Glucose Levels for Age	
Birth to 4 hours of life	≥ 35 mg/dL
>4 – 24 hours of life	≥ 45 mg/dL
>24 – 48 hours of life	≥ 50 mg/dL
>48 hours of life	≥ 60 mg/dL

Symptoms of Hypoglycemia

- Irritability
- Tremors/jitteriness
- Exaggerated Moro reflex
- High-pitched cry
- Seizures
- Lethargy
- Floppiness
- Cyanosis
- Apnea
- Poor Feeding


0-4 HOURS OF AGE

At Risk Infants (See Indications Box)

Begin feeding within one hour of birth. First glucose should be obtained 30 minutes after completion of first feed.

Indications for Screening of At Risk Infants


- **SGA** (<10%ile BW) or IUGR
- **LGA** (>90%ile BW)
- Infant of Diabetic Mother
- Maternal exposure to labetalol
- Late preterm (34 0/7 – 36 6/7) or post-term (>42 weeks)
- Perinatal stress (C-section for fetal distress, maternal preeclampsia or severe HTN, etc.)
- Physician discretion



GLUCOSE 40% GEL DOSING

Birth Weight	Dose
≤ 2 kg	1 mL
>2-2.5 kg	1.25 mL
>2.5-3 kg	1.5 mL
>3-3.5 kg	1.75 mL
>3.5-4 kg	2 mL
>4-4.5 kg	2.25 mL
>4.5-5 kg	2.5 mL

> 4 - 48 HOURS OF AGE



If infant has severe symptoms or BG is <25 after first Dextrose Gel dose THE ABOVE PROTOCOL NO LONGER APPLIES.

- Give Glucose Gel dose.
- Start IV.
- Give D10 2 mL/kg bolus at 1 mL/minute.
- Start D10 infusion at 80 mL/kg/day.
- Goal is to keep baby's serum glucose at 60.
- Check glucose 30 minutes after each bolus or rate change and Q1-2h until stable.
- If glucose remains low, give another D10 2 mL/kg bolus and increase hourly rate by 1 mL/hour.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 3/11/24.
Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact
Amy_Carson-Strnad@ykhc.org.



PAMC Transfer center - 907-212-7363
NICU (907) 212-3614 – Ask for attending neonatologist on call.
Neonatologist direct line (for emergencies) (907) 212-2068.

GESTATIONAL AGE (weeks)	24	26	28	30	32	34	36	38	40
ESTIMATED WEIGHT (grams)	700	900	1100	1350	1650	2100	2600	3000	3500
EQUIPMENT/SUPPLIES: NG/OG Tube - 5 French + UVC <32 weeks - 3.5 French + UVC ≥32 weeks - 5 French									
Laryngoscope Blade	00	00	00	0	0	0	0	0-1	0-1
ETT Size	2.5	2.5	2.5-3.0	3.0	3.0	3.0-3.5	3.5	3.5-4.0	3.5-4.0
ETT Depth lip to tip. <i>Place at T2 above the carina.</i>	6.5-7 cm	6.5-7 cm	7 cm	7-7.5 cm	7.5 cm	8 cm	8.5 cm	9 cm	9.5 cm
Laryngeal mask airway (LMA)	none	none	none	none	Consult NICU.	1	1	1	1
Needle decompression. <i>See kit and protocol in neonatal code cart.</i>	18 gauge	18 gauge	18 gauge	18 gauge	18 gauge	16 gauge	16 gauge	16 gauge	16 gauge
UVC insertion. <i>Place just above diaphragm. Add umbilical stump length.</i> May insert UVC 2-4 cm for emergency access.	6.5 cm	6.9 cm	7.2 cm	7.5 cm	8 cm	8.7 cm	9.4 cm	10 cm	10.8 cm

VITAL SIGNS: Heart Rate 120-160 + Respiratory Rate 30-60 + Mean Blood Pressure = Gestational age in weeks									
INITIAL VENTILATOR SETTINGS									
Positive Inspiratory Pressure (PIP) cm H ₂ O	16-22	16-22	16-22	16-22	18-24	18-24	18-24	20-28	20-28
Positive End Expiratory Pressure (PEEP) cm H ₂ O	4-6	4-6	4-6	4-6	4-6	5-6	5-6	5-6	5-6
Inspiratory Time (seconds)	0.3-0.35	0.3-0.35	0.3-0.35	0.3-0.35	0.3-0.35	0.3-0.35	0.35-0.4	0.35-0.4	0.35-0.4
Respiratory Rate (breaths per minute)	30-45	30-45	30-45	30-45	20-40	20-40	20-40	20-40	20-40
Saturation Goal after 10 Minutes	88-95%	88-95%	88-95%	88-95%	88-95%	88-95%	88-95%	95-98%	95-98%

MEDICATIONS									
Epinephrine IV/IO 0.1 mg/mL 0.2 mL/kg. <i>May repeat every 3 minutes for asystole.</i>	0.14 mL	0.18 mL	0.22 mL	0.27 mL	0.33 mL	0.4 mL	0.5 mL	0.6 mL	0.7 mL
Epinephrine ET ONLY 0.1 mg/mL 1 mL/kg. <i>May repeat every 3 minutes for asystole.</i>	0.7 mL	0.9 mL	1.1 mL	1.3 mL	1.6 mL	2.1 mL	2.6 mL	3 mL	3.5 mL
Curosurf (poractant alfa 80 mg/mL) 2.5 mL/kg as a single dose. Give Curosurf <26 weeks OR 26-29 weeks requiring ≥40% FiO ₂ OR >29 weeks with CXR-proven RDS. See guideline for more details.	1.8 mL	2.2 mL	2.8 mL	3.4 mL	4 mL	5.2 mL	6.6 mL	7.6 mL	8.8 mL
FOR HYPOGLYCEMIA: Give D10 bolus 2 mL/kg IV/IO at 1 mL/min. Increase D10 maintenance fluid rate by 1 mL/hour for <2 kg or 2 mL/hour for ≥2 kg.	1.4 mL	1.8 mL	2.2 mL	2.7 mL	3.3 mL	4.2 mL	5.2 mL	6 mL	7 mL
Ampicillin (Dilute to 100 mg/mL) 50 mg/kg IV/IM	35 mg (0.35 mL)	45 mg (0.45 mL)	55 mg (0.55 mL)	68 mg (0.68 mL)	83 mg (0.83 mL)	105 mg (1.05 mL)	130 mg (1.3 mL)	150 mg (1.5 mL)	175 mg (1.75 mL)
Gentamicin (2 mg/mL) 5 mg/kg IV as one-time dose. May give IM. DO NOT USE IN VILLAGE.	3.5 mg (1.75 mL)	4.5 mg (2.25 mL)	5.5 mg (2.75 mL)	6.8 mg (3.4 mL)	8.2 mg (4.1 mL)	10.4 mg (5.2 mL)	13 mg (6.5 mL)	15 mg (7.5 mL)	17.6 mg (8.8 mL)
Volume Expanders: NS or albumin 10 mL/kg IV/IO. Give over 15-30 minutes; give faster if unstable; give slower for extreme premies.	7 mL	9 mL	11 mL	13.5 mL	16.5 mL	21 mL	26 mL	30 mL	35 mL
D10 Maintenance Fluids: <750 grams give 90-100 mL/kg/24 hours + ≥750 grams give 80 mL/kg/24 hours. Goal blood glucose is 50-110 mg/dL.	3 mL/hour	3 mL/hour	3.7 mL/hour	4.5 mL/hour	5.5 mL/hour	7 mL/hour	8.7 mL/hour	10 mL/hour	12 mL/hour
Phenobarbital (130 mg/mL) 10 mg/kg IV/IO/IM/PR. May give additional 10 mg/kg dose.	7 mg (0.05 mL)	9 mg (0.07 mL)	11 mg (0.08 mL)	13.5 mg (0.1 mL)	16.5 mg (0.13 mL)	21 mg (0.16 mL)	26 mg (0.2 mL)	30 mg (0.23 mL)	35 mg (0.27 mL)



PAMC Transfer center - 907-212-7363
NICU (907) 212-3614 – Ask for attending neonatologist on call.
Neonatologist direct line (for emergencies) (907) 212-2068.

Epinephrine 0.1 mg/mL

- This is the pre-filled syringe concentration.
- Draw up doses by inserting needle through the thick rubber stopper.
- Flush with 3 mL of NS regardless of weight or gestational age.

Ampicillin 100 mg/mL

Products needed:

- Ampicillin 500 mg vial
- Sterile water for injection, 10 mL vial

How to mix:

1. Reconstitute 500 mg vial with 4.8 mL sterile water for injection. This will result in a 100 mg/mL final concentration.
2. The Neonatal Resuscitation Summary (page 1) lists the total dose and volume – draw up dose from vial.
3. Dose must be used within 1 hour of reconstitution.

Administration:

- Doses less than 500 mg can be injected via slow IV push over 3 to 5 minutes.
- Not compatible with D10W.
- Administer before gentamicin – do not administer at the same time.

Gentamicin 2 mg/mL

Product needed:

- Gentamicin 100 mg/50 mL pre-mixed bag.

DO NOT ADMINISTER THE BAG – the dose will be administered via syringe pump.

The Neonatal Resuscitation Summary (page 1) lists the total dose and volume – draw up this volume from the bag and **immediately dispose of the bag.**

Administration:

- Administer after ampicillin – do not administer at the same time.
- Administer via **syringe pump** over 30 minutes.
- Compatible with D10W.

Rapid Sequence Intubation Medications: Consult NICU prior to use. Do not use in routine resuscitation. Consider for surfactant administration and in difficult airway.

GESTATIONAL AGE (weeks)	24	26	28	30	32	34	36	38	40
ESTIMATED WEIGHT (grams)	700	900	1100	1350	1650	2100	2600	3000	3500
Atropine (0.1 mg/mL) – 0.02 mg/kg	0.01 mg (0.1 mL)	0.02 mg (0.2 mL)	0.02 mg (0.2 mL)	0.03 mg (0.3 mL)	0.03 mg (0.3 mL)	0.04 mg (0.4 mL)	0.05 mg (0.5 mL)	0.06 mg (0.6 mL)	0.07 mg (0.7 mL)
Fentanyl (**10 mcg/mL**) – 1 mcg/kg (May repeat dose once.) Push slowly over 3-5 minutes. Have dose of rocuronium drawn up in case of chest wall rigidity.	0.7 mcg (0.07 mL)	0.9 mcg (0.09 mL)	1.1 mcg (0.11 mL)	1.4 mcg (0.14 mL)	1.7 mcg (0.17 mL)	2.1 mcg (0.21 mL)	2.6 mcg (0.26 mL)	3 mcg (0.3 mL)	3.5 mcg (0.35 mL)
Rocuronium (10 mg/mL) – 0.6 mg/kg Do not routinely use. Reserve for difficult airways.	0.4 mg (0.04 mL)	0.5 mg (0.05 mL)	0.7 mg (0.07 mL)	0.8 mg (0.08 mL)	1 mg (0.1 mL)	1.3 mg (0.13 mL)	1.6 mg (0.16 mL)	1.8 mg (0.18 mL)	2.1 mg (0.21 mL)
Naloxone (0.4 mg/mL) – 0.1 mg/kg	0.07 mg (0.18 mL)	0.09 mg (0.23 mL)	0.11 mg (0.28 mL)	0.14 mg (0.35 mL)	0.16 mg (0.4 mL)	0.2 mg (0.5 mL)	0.26 mg (0.65 mL)	0.3 mg (0.75 mL)	0.35 mg (0.9 mL)

Fentanyl 10 mcg/mL

Products needed:

- Fentanyl 50 mcg/mL, 2 mL vial
- Preservative-free normal saline

How to mix:

1. Draw up 1 mL of fentanyl 50 mcg/mL.
2. Add to 4 mL of normal saline.

Administration:

- Inject via slow IV push over 3 to 5 minutes.
- If chest wall rigidity develops, give dose of rocuronium or naloxone.

RSI Drug Notes


• Drug Locations:

- Atropine and naloxone located in neonatal code cart.
- Fentanyl 50 mcg/mL and rocuronium located in OB Pyxis.


- May flush with 0.5-1 mL of NS if needed.





Attach the oxygen tubing to a 15 L flow meter.
Set blender to 21% and consider increasing depending on clinical status.
Set the flow meter to **10 L**.



Occlude both the mask and the hole.
Set the PIP: Turn the knob labeled Peak Inspiratory Pressure until the arrow on the dial points to **20**.





Occlude only the mask.
Set the PEEP: Turn the PEEP knob until the arrow on the dial points to **5**.




Troubleshooting: If you cannot achieve the desired pressures, try changing the liters on the flow meter or turning the Max Pressure Relief knob located under the flap.





- Turn Gas Supply switch on. Down is ON.
- Set blender to 21% and consider increasing depending on clinical status.
- Set the top flow meter to **10 L**.
- The bottom flow meter is for use with nasal cannula.



- Occlude both the mask and the hole.
- Set the PIP: Turn the knob labeled Airway Pressure Relief until the arrow on the dial points to **20**.



- Occlude only the mask.
- Set the PEEP: Turn the PEEP knob until the arrow on the dial points to **5**.




Troubleshooting: If you cannot achieve the desired pressures, try changing the liters on the flow meter.

This protocol is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 9/16/24.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.





- Turn on top left switch. Down is ON.
- Set blender to 21% and consider increasing depending on clinical status.
- Set the right flow meter to **10 L**.
- The left flow meter is for use with nasal cannula.

- Occlude both the mask and the hole.
- Set the PIP: Turn the knob labeled PIP until the arrow on the dial points to **20**.



- Occlude only the mask.
- Set the PEEP: Turn the PEEP knob until the arrow on the dial points to **5**.




Troubleshooting: If you cannot achieve the desired pressures, try changing the liters on the flow meter.

This protocol is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 9/16/24.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



There are two ON switches – one in the back (top photo) and one on front below the screen (bottom photo).

The respiratory equipment may be used without turning the warmer on.

(Please follow the shut-down procedure located on the warmer when finished.)


- When turned on, the warmer will automatically heat to 100% for three minutes and then will decrease to 30%.
- To change this, tap “Pre.” A percentage will appear. Turn the knob to the right and below the screen to adjust the percentage. Tap the knob to confirm the setting.



Skin Mode (sometimes called servo mode)

This mode is like a thermostat. The warmer heats and cools in response to the baby's temperature and auto-adjusts to maintain the temperature set.

- To use this mode, plug a temperature probe in.
- Tap “Skin Mode” on the screen.
- Affix the temperature probe to the baby's skin overlying the liver (RUQ) with a metallic sticker on top.
- To adjust the desired temperature, tap the number above “Skin temp.” Turn the knob to change the number. Tap the knob to confirm the setting.



Celsius	Fahrenheit
36°	96.8°
36.5°	97.7°
37°	98.6°
37.5°	99.5°
38°	100.4°

This protocol is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 9/16/24.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



Surfactant Administration Protocol

Indications for Curosurf®

- GA < 26 weeks.
- GA 26-29 weeks with supplemental oxygen requirement $\geq 40\%$.
- GA > 29 weeks with CXR-proven RDS.
- Consider in severe meconium aspiration after consultation with neonatologist.
- Ability to confirm placement of ETT with CXR. (Must be in Bethel or SRC.)

Curosurf® Storage

- Curosurf® is stored at 36-46°F.
- If warmed and not opened or used, may be returned to refrigerated storage one time.
- Curosurf® is located in the OB medication refrigerator. If going on a medevac, ask the nurses to get the Curosurf®. It can be stored in a pink thermal bag that is kept next to it in the refrigerator.

Reference:

See this [YouTube video](#) for a demonstration of the Y catheter. Please instill as single dose in supine position to decrease chance of inadvertent loss of ETT placement/position.

Troubleshooting

- If ETT cap is stuck, cut the tube as high as possible and then place the Y cap. This will change the number of the desired depth.
- If having trouble switching caps prior to intubation, hold ETT up to warmer to soften the plastic.

Prior to procedure, discuss with neonatologist. Consider using medications for Rapid Sequence Intubation. If planning to use paralytic, discuss with neonatologist.

Preparation of Curosurf®

- Warm to room temperature and gently invert. Do not shake.
- Choose Curosurf® dose using the [Neonatal Resuscitation Summary](#) using estimated gestational age. If weight is known, calculate dose to be 2.5 mL/kg.
- Draw up total Curosurf® dose using a 20 gauge or larger needle. Draw up 1-2 mL of air into syringe after dose. This helps the surfactant get pushed out of the catheter when administering dose.

Preparation of Equipment and Patient

- Prior to intubation, if possible, check the ETT cap and make sure it comes on and off easily.
- Make sure you have the correct size Y cap for the ETT size.
- Check fit of Y cap on ETT. Attach catheter and feed it down the tube until it is at the tip of the ETT. Look for the number or color that will tell you the depth of the catheter at this point. If time, review YouTube video in reference box for demonstration of how to achieve desired depth.
- Intubate patient with ETT cap on tube.
- Verify placement and secure tube.
- Perform CXR to verify tube placement and rule-out pneumothorax.

Administration of Curosurf®

- Infant should be supine.
- Disconnect Neopuff, bag, or ventilator.
- Remove ETT cap and replace with Y cap. This will change the number of the desired depth.
- Attach the Neopuff or anesthesia bag to the larger port on the Y cap.
- Attach the catheter to the smaller port on the Y cap and advance it until it is at the desired depth.
- Inject the syringe of Curosurf® through the catheter.
- Pull the catheter all the way out but leave attached.
- Bag the baby at a rate of 40-60 breaths/minute for one minute.
- Allow the baby to recover.
- Resume ventilation.
- Do not suction for one hour after administration unless required for obstruction.
- Adjust pressure on Neopuff as lung compliance improves.

Note: After discussion with the neonatologists, surfactant is not to be given in villages without CXR ability. It may be given in SRCs after CXR confirms ETT position.

This resource is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/23/25.

Click [here](#) to see the supplemental resources for this resource.

If comments about this resource, please contact Amy_Carson-Strnad@ykhc.org.



Suspected Neonatal Withdrawal

Signs and Symptoms of Neonatal Withdrawal

- Hyperactivity/excessive alertness
- Crying/irritability/restlessness
- High-pitched cry
- Poor suck/feeding difficulties
- Hyperphagia or excessive suck
- Tremors and/or seizures
- Poor sleeping patterns
- Diaphoresis
- Hyperacusis (sensitivity to noise)
- Vasomotor instability/mottling
- Autonomic dysfunction
- Temperature instability
- Diarrhea or vomiting
- Hypertonia or hypotonia
- Tachypnea, respiratory distress, or apnea
- Hypoglycemia
- Hyperactive reflexes (including Moro)
- Hypertension or tachycardia
- Yawning

Helpful Links

- [Template](#) for prenatal counseling.
- [Brochure](#) for families.
- [Video](#) from ANTHC.

Non-Pharmacologic Interventions (NPI)


- Rooming-in: Parent/caregiver presence to help calm/care for infant. Separation of parents and infant should be avoided unless medically indicated.
- Skin-to-skin to help organize infant feeding behaviors, calming, and sleep.
- Holding by parent/caregiver.
- Safe swaddling with extremities flexed.
- Optimal feeding: Offer feeds when cues and feed until content.
- Non-nutritive sucking with infant's hand, pacifier, gloved finger, etc.
- Quiet, low light environment.
- Rhythmic movement like jiggling or swing.
- Additional support from other family members, etc.
- Limit number of visitors and duration of visits.
- Clustering care/assessments with awake times.
- Safe sleep/fall prevention.
- Parent/caregiver self-care/rest.

Neonatal Drug Testing

- Urine: low sensitivity
- Meconium: high sensitivity, reflects drug use in the second and third trimesters, consider if clinical questions and practical

ESC Assessment

- **Eating:** YES to any (and cannot attribute to etiology other than withdrawal):
 - Takes >10 min to coordinate feeding
 - Breastfeeds <10 min
 - Feeds <10 mL (or other age-appropriate duration/volume)
- **Sleeping:** Sleeps <1 hour (and cannot attribute to etiology other than withdrawal)
- **Consoling:**
 - 1: Consoles on own.
 - 2: Able to console within 10 minutes with caregiver support.
 - 3: Takes >10 minutes to console or cannot stay consoled for at least 10 minutes despite caregiver best efforts.



Feeding Details

- YKHC's [Policy & Procedure on Breastfeeding](#) states, "Mothers should NOT breastfeed or feed expressed breast milk to their infants if...Mother is using an illicit street drug, such as PCP (phencyclidine) or cocaine...(Exception: Narcotic-dependent mothers who are enrolled in a supervised methadone or suboxone program and have a negative screening for HIV infection and other illicit drugs can breastfeed)"
- Infants experiencing withdrawal can have higher caloric requirements. Have a low threshold to increase caloric density of feeds to 22-24 kcal/ounce.

Note: Chronic *in utero* exposure to illicit and/or prescription drugs can lead to habituation. Signs and symptoms of withdrawal worsen as drug level decreases while signs and symptoms of acute toxicity lessen with drug elimination. Withdrawal from tobacco or Iqmiq can manifest similarly.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 12/12/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Amy_Carson-Strnad@ykhc.org.



Village Deliveries (Pediatrics)

Preparation in the Village for the Health Aides

- Turn the heat up until everyone is sweating. May need extra space heaters.
- In the warmest part of the clinic, prepare a table with clean blankets, towels, saran wrap, etc.
- If the clinic has a dryer, instruct the health aides to warm the blankets there prior to birth.
- Ensure the following are prepared and functional: suction, oxygen tanks and tubing, BVM with smallest available mask, bulb suction.
- Seek out extra health aides or former health aides to help.

See [Labor Patient In Village](#) guideline for OB recommendations.

Preparation for Medevac

- Review prenatal history and note risk factors for the baby.
- Coordinate with family medicine hospitalist activating the medevac and LifeMed crew about when to meet at the hangar. The LifeMed hangar is located at 3600 Tower Road.
- Turn over the Tiger Connect role for "Peds Wards on Duty" to another pediatrician or the family medicine hospitalist staying behind.
- Establish roles with LifeMed crew. Discuss doses and equipment based on estimated GA.

What to Bring

- Curosurf if going to an SRC and GA <32 weeks or unknown: located in the OB medication refrigerator. Place in pink thermal case.
- OB & Pediatric Village Delivery Backpack containing OB and pediatric supplies located in the nursery.
- Resources: [Neonatal Resuscitation Summary](#), [Surfactant Administration](#), [Neopuff Set Up Guide](#), [Neonatal Glucose Screening Guideline](#).
- Warm clothing. (There is extra warm gear under the bed in the peds call room.)
- Snacks, drinks, money, motion sickness medication.

Resuscitation

- Resuscitate per NRP algorithm. Remember that CPAP is a great tool for non-invasive respiratory support for transport.

For infants <32 weeks:

- Place infant directly into polyurethane bag without drying. If intubated, bag may cover face/head.
- Attempt IV or UVC access early.
- See [Surfactant Protocol](#), if indicated and in SRC.

Delivery is Imminent

- Set up monitor, Neopuff, and intubation equipment (all carried by LifeMed), using sizes recommended by [Neonatal Resuscitation Summary](#).
- Activate chemical mattress just prior to delivery. Cover with single baby blanket.

For High Risk Deliveries, including GA <32 weeks:

- Discuss with neonatologist early – call (907) 212-3614.
- Activate medevac to Anchorage. Consider direct transfer from village, ramp transfer in Bethel, or further stabilization with NICU team in Bethel, as appropriate.
- Prepare polyurethane bag.

Delivery is not Imminent

- Hospitalist assesses mother, does vaginal exam, obtains cultures, etc.
- LifeMed crew cares for mother.
- Pediatrician should help however possible and otherwise stay out of the way.
- Occasionally a mother will be transported to Bethel dilated and in labor. This decision is made if the benefit of being at a higher level of care outweighs the risks of potential delivery en route.

Prior to Transport

- Inform OB of baby's status and request they call ER Registration to get chart made – should be Peds FIN with baby going to OB unit on Family Medicine service.
- Activate Anchorage team if needed.

Term or Late Preterm Infants

- Admit to YKHC OB for 48 hours.
- Do not obtain blood cultures unless sick or [otherwise indicated](#).

Medications

- Give erythromycin to eyes and vitamin K IM if infant is stable.
- Hepatitis B and HBIg can wait until arrival in Bethel.
- Give ampicillin per Neonatal Resuscitation Summary for all preterm and high risk infants.
- Gentamicin should not be given in the village, as it is high-risk.

Temperature

- Hypothermia in newborns is defined as temp <97.7°F.
- Cold babies do very poorly. It is better to over-prepare (use a polyurethane bag in term babies, etc.) rather than under-prepare.
- The baby pod carried by LifeMed does not have a heat source. It will not generate heat. Avoid placing the baby into it until it has warmed from being outside.
- Check axillary temperature at 5 minutes of life and then Q30 minutes.
- Place a hat and/or saran wrap on the baby as soon as possible.
- Do not remove hat, chemical mattress, or bag until arrived at YKHC.
- You may tear holes in the bag to gain access to the baby for procedures.
- Avoid weighing premature babies, as this frequently contributes to heat loss in the village.

Glucose

- Check glucose as soon as possible and then Q30-60 minutes until stable.
- See [Neonatal Glucose Screening Guideline](#). Goal glucose is >35 in first four hours of life.
- On babies <32 weeks, start D10 maintenance as soon as IV access has been established.
- If unable to get a glucose, have a low threshold to give sugar in preterm or high risk infants.
- If oral dextrose gel unavailable, may give Sweetease, oral glucose, colostrum, formula, or homemade sugar paste. May smear on gums for buccal absorption.

Procedures

Intubation

- Prepare equipment.
- Wipe upper lip and rest of face.
- See [Neonatal Resuscitation Summary](#), especially if considering RSI.
- Intubate and confirm placement with auscultation and ET/CO₂ detector.
- Tape tube with Benzoin and tape.
- Consider using Neopuff to ventilate en route rather than ventilator.

UVC (Always attempt PIV placement first unless infant is very unstable.)

- Use sterile technique.
- Flush catheter and stopcock with sterile saline. NOTE: the syringes for premade saline flushes are not sterile. You will have to use a sterile syringe to draw up flushes from a NS bag.
- If baby is in polyurethane bag, tear a small opening in the plastic.
- Place the UVC just far enough to get blood return.
- Cover skin around umbilicus with Tegaderm. Tape the UVC to the Tegaderm to secure it.


See [Surfactant Administration](#) resource if in SRC.

This resource is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guidelines Committee 7/23/25.

Click [here](#) to see the supplemental resources for this resource.

If comments about this resource, please contact Amy_Carson-Strnad@ykhc.org.





Soft Marker	Aneuploidy Evaluation	Antenatal Management	Follow-up Imaging
Echogenic intracardiac focus	<ul style="list-style-type: none"> • cfDNA or quad screen negative: none • No previous screening: counseling for noninvasive testing for aneuploidy 	Routine care	N/A
Echogenic bowel	<ul style="list-style-type: none"> • cfDNA or quad screen negative: none • No previous screening: counseling for noninvasive testing for aneuploidy 	Evaluation for cystic fibrosis, congenital viral infection, intra-amniotic bleeding	Third-trimester ultrasound examination for reassessment and evaluation of growth
Choroid plexus cyst	<ul style="list-style-type: none"> • cfDNA or quad screen negative: none • No previous screening: counseling for noninvasive testing for aneuploidy 	Routine care	N/A
Single umbilical artery	cfDNA or quad screen negative or no previous screening: none	Consideration for weekly antenatal surveillance beginning at 36 0/7 week of gestation	Third-trimester ultrasound examination for evaluation of growth
Urinary tract dilation	<ul style="list-style-type: none"> • cfDNA or quad screen negative: none • No previous screening: counseling for noninvasive testing for aneuploidy 	Evaluation for persistence, with frequency of evaluation dependent on initial findings	Third-trimester ultrasound examination to determine whether postnatal pediatric urology or nephrology follow-up is needed
Shortened humerus, femur, or both	<ul style="list-style-type: none"> • cfDNA or quad screen negative: none • No previous screening: counseling for noninvasive testing for aneuploidy 	Evaluation for skeletal dysplasias	Third-trimester ultrasound examination for reassessment and evaluation of growth
Thickened nuchal fold	<ul style="list-style-type: none"> • cfDNA negative: none • Quad screen negative: counseling for no further testing vs noninvasive vs invasive testing for aneuploidy • No previous screening: counseling for noninvasive vs invasive testing for aneuploidy 	Routine care	N/A
Absent or hypoplastic nasal bone	<ul style="list-style-type: none"> • cfDNA negative: none • Quad screen negative: counseling for no further testing vs noninvasive vs invasive testing for aneuploidy • No previous screening: counseling for noninvasive vs invasive testing for aneuploidy 	Routine care	N/A

Abbreviations

- cfDNA: cell-free DNA – order in RAVEN as MatemiT21
- CF: cystic fibrosis
- Quad screen: order in RAVEN as AFP Maternal (Quad Screen)

Contact

MFM: Send referral through RAVEN via “Refer to Obstetrics External – Perinatology.”
For non-beneficiaries, place this order AND send a message to Women’s Health Case Manager to ensure it is sent to the correct place.

Source

Society for Maternal-Fetal Medicine. SMFM Consult Series #57: Evaluation and management of isolated soft ultrasound markers for aneuploidy in the second trimester. Am J Obstet Gynecol 2021.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/24/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Someone@ykhc.org.



Abbreviations and Definitions

- **Glucose Screening Test (GST):** fasting or fed plasma glucose value one hour after 50 gram glucose load.
- **Gestational Glucose Tolerance Test (GGTT):** fasting plasma glucose value one hour and two hours after 75 gram glucose load.
- **BGM:** Blood Glucose Monitoring
- **Pre-gestational Diabetes:** patient with diagnosis of diabetes prior to pregnancy.
- **DSMES:** Diabetes Self-Management Education and Support

Block Testing

Monitoring at different times of the day to identify patterns. For example, on some days the patient will check fasting levels, on other days check pre-meal levels, and on other days check levels 1-2 hours post-meal.

Diagnostic Criteria for GDM Utilizing Two Hour 75 g GGTT

Pregnant patient with any of the following:
Fasting glucose ≥ 92 mg/dL
1 hour after oral load, glucose ≥ 180 mg/dL
2 hours after oral load, glucose ≥ 153 mg/dL

BGM Targets

- Fasting glucose < 95 mg/dL
- 2 hour post-prandial glucose < 120 mg/dL
- 1 hour post-prandial glucose < 140 mg/dL


Patients with Suboptimal Participation in Care

- Send letter after two weeks of not sending in sugar logs or two weeks of $< 25\%$ of expected readings.
- Consider admission to monitor blood sugars.
- Consider transfer to ANMC at 32 weeks.

Postpartum Management of All Patients with GDM

- Add diagnosis "History of gestational diabetes" to Problem List.
- At the 6 week postpartum visit, perform two hour 75 gram Oral Glucose Tolerance Test with fasting and two hour blood draws only. NOTE: the criteria are different than in pregnancy.
- As an alternative, at > 12 weeks postpartum, check HgA1C.
- Diabetes screening every three years.

At first prenatal visit, check HgA1C in all patients.



Fetal Monitoring

- **Diet, well-controlled**
 - 28 weeks: kick counts.
 - Normal labor management.
- **Diet, poorly controlled:**
 - 28 weeks: kick counts.
 - 32 weeks: BPP weekly.
 - 38 weeks: Consult OB and consider induction.
- **Insulin-controlled:**
 - 28 weeks: kick counts.
 - 32 weeks: weekly BPP.
 - 32-35 weeks: transfer to Anchorage.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/13/23. Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact David.Compton@ykhc.org.



D&C Prior to Methotrexate?

D&C is NOT necessary prior to treatment with Methotrexate (MTX) for a plateau or abnormally rising hCG level. MTX will treat an abnormal pregnancy in the uterus or any other location.

Typical side effects of MTX

- Less than 30% of patients will experience minor, self-limited side effects from the medication, including nausea, mouth ulcers, and GI cramps.
- Most patients have some lower abdominal pain on the 3-6th day after treatment. This is not a problem if ibuprofen or acetaminophen relieves the pain.

Contraindication to MTX

Absolute contraindications

- Breast Feeding
- Overt or laboratory evidence of immunodeficiency
- Alcoholism, alcoholic liver disease, or other chronic liver disease
- Preexisting blood dyscrasias, such as bone marrow hypoplasia, leukopenia, thrombocytopenia, or significant anemia
- Known sensitivity to MTX
- Active pulmonary disease
- Peptic ulcer disease
- Hepatic, renal, or hematologic dysfunction

Relative contraindications

- Gestational sac larger than 3.5cm
- Embryonic cardiac motion

Single Dose Regimen

- Single dose MTX 50 mg/m² IM on day 1.
- Measure hCG level on post-treatment days 4 and 7.
- Check for 15% hCG decrease between days 4 and 7.
- Then measure hCG level weekly until reaching the nonpregnant level
- If results are less than the expected 15% decrease, readminister MTX 50 mg/m² and repeat hCG measurement on days 4 and 7 after second dose.

If at any time the hCG level rises during the monitoring of weekly hCG levels, consult a GYN Oncologist for further treatment.

Ectopic Pregnancy diagnosed after consultation with HROB or OB/GYN.

Work-up

- Quantitative hCG
- Type and Screen
- CBC
- Comprehensive Metabolic Panel
- Transvaginal Pelvic Ultrasound (US)

Hemodynamically stable?

No → Consult HROB for immediate surgery or transfer.

Yes

Adnexal Mass ≥ 4 cm
Cardiac activity
Pregnancy in location other than a tube

Yes → Consult HROB for immediate surgery or transfer.

No

Platelets,
kidney,
and liver function
normal?

No → Consult HROB for immediate surgery or transfer.

Yes

Is the
hCG >5000 or is
there a large
mass?

No → Single Dose Regimen

Yes → Two Dose Regimen

Two Dose Regimen

- Administer 50 mg/m² on day 1.
- Repeat 50 mg/m² on day 4.
- Measure hCG levels on days 4 and 7, and expect a 15% decrease between days 4 and 7.
- If the decrease is greater than 15%, measure hCG levels weekly until reaching non-pregnant level.
- If less than a 15% decrease in hCG levels, readminister MTX 50 mg/m² on days 7 and 11, measuring hCG levels.
- If hCG levels decrease 15% between days 7 and 11, continue to monitor weekly until non pregnant hCG levels are reached.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 10/21/22. Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact David.Compton@ykhc.org.



Nomenclature


- **Viable:** A pregnancy is viable if it can potentially result in a liveborn baby.
- **Nonviable:** A pregnancy is nonviable if it cannot possibly result in a liveborn baby. Ectopic pregnancies and failed intrauterine pregnancies are nonviable.
- **Intrauterine pregnancy of uncertain viability:** A patient is considered to have this if a transvaginal ultrasound shows an intrauterine gestational sac with no embryonic heartbeat and no findings of definite pregnancy failure.
- **Pregnancy of unknown location:** A patient is considered to have this if there is a positive urine or serum pregnancy test and no intrauterine or ectopic pregnancy on transvaginal ultrasound.

Findings Diagnostic of Pregnancy Failure

- Crown-rump length of $\geq 7\text{mm}$ and no heartbeat.
- Mean sac diameter of $\geq 25\text{mm}$ and no embryo.
- Absence of embryo with heartbeat ≥ 14 days after an US that showed a gestational sac without a yolk sac.
- Absence of embryo with a heartbeat ≥ 11 days after an US that showed a gestational sac with a yolk sac.

Comments

- In a patient with a positive urine or serum pregnancy test, an intrauterine fluid collection with rounded edges containing no yolk sac is most likely a gestational sac; it is certain to be a gestational sac if it contains a yolk sac or embryo.
- Transabdominal imaging without transvaginal scanning may be sufficient for diagnosing early pregnancy failure when an embryo whose crown-rump length is 15 mm or more has no visible cardiac activity.
- Point of care ultrasound performed in the ED or clinic is an ultrasound for the purposes of this guideline. The ultrasound does not need to be performed in Diagnostic Imaging.





Nomenclature

- **Viable:** A pregnancy is viable if it can potentially result in a liveborn baby.
- **Nonviable:** A pregnancy is nonviable if it cannot possibly result in a liveborn baby. Ectopic pregnancies and failed intrauterine pregnancies are nonviable.
- **Intrauterine pregnancy of uncertain viability:** A patient is considered to have this if a transvaginal ultrasound shows an intrauterine gestational sac with no embryonic heartbeat and no findings of definite pregnancy failure.
- **Pregnancy of unknown location:** A patient is considered to have this if there is a positive urine or serum pregnancy test and no intrauterine or ectopic pregnancy on transvaginal ultrasound.


Findings Diagnostic of Pregnancy Failure

- Crown-rump length of $\geq 7\text{mm}$ and no heartbeat.
- Mean sac diameter of $\geq 25\text{mm}$ and no embryo.
- Absence of embryo with heartbeat ≥ 14 days after an US that showed a gestational sac without a yolk sac.
- Absence of embryo with a heartbeat ≥ 11 days after an US that showed a gestational sac with a yolk sac.
- Falling hCG level.

Comments

- In a patient with a positive urine or serum pregnancy test, an intrauterine fluid collection with rounded edges containing no yolk sac is most likely a gestational sac; it is certain to be a gestational sac if it contains a yolk sac or embryo.
- Transabdominal imaging without transvaginal scanning may be sufficient for diagnosing early pregnancy failure when an embryo whose crown-rump length is 15 mm or more has no visible cardiac activity.

Pregnancy of uncertain viability or unknown location



If patient elects wait and see option

- Must be reliable patient who will stay in Bethel.
- Must be followed up every 48 hours for repeat hCG.

If patient elects misoprostol only

- Consult HROB.
- Must be reliable patient who will stay in Bethel.
- Regimen is misoprostol 800 mcg vaginally.
- Follow-up daily.
- Offer ibuprofen for cramping.

If patient elects misoprostol/mifepristone option

- Consult HROB.
- Must be reliable patient who will stay in Bethel.
- Regimen is mifepristone 200 mg oral followed 24-48 hours later with misoprostol 800 mcg placed in posterior fornix of vagina.
- Follow-up 24-48 hours after vaginal misoprostol.
- Offer ibuprofen for cramping.
- Dose can be repeated in 24 hours if uterus is not empty.

If patient elects D&C option

- Consult HROB.
- Consider office-based D&C.
- To schedule procedure, send message via Tiger Connect to OR Charge Nurse on call and OR CRNA on call.
- If on weekend, have patient remain NPO after midnight on Sunday for Monday 0800 procedure.

Following hCG to negative

- Contact GYN CM at 543-6557 or send communication in RAVEN to Women's Health Case Manager Pool.
- CM will follow hCG levels in consultation with HROB.



Maternal GBS Prophylaxis

Use the GBS App

to determine need for prophylaxis and antibiotic of choice for GBS prevention

Web version: <https://www2a.cdc.gov/vaccines/m/gbs3/gbs.html>

or

Download for your smartphone.

Please note: YKHC does not use the neonatal option available here. Please see the [Newborn Early-Onset Sepsis/GBS guideline](#) for more details.

OPT OUT


Default is that patient has HIV screening performed unless she specifically declines the test.

Resources and Abbreviations

EIS: Early Intervention Services (907) 729-2907
BIB: Be in Bethel appointment
HAART: highly-active anti-retroviral therapy

Pre-exposure Prophylaxis for Prevention of HIV (PrEP)


Daily dose of combination emtricitabine and tenofovir to prevent HIV transmission in high risk individuals



This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 10/21/22. Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact David.Compton@ykhc.org.



Any patient with hypertension in pregnancy should have blood pressure monitored for at least two weeks postpartum.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 10/21/22.
Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact David_Compton@ykhc.org.



How to Take a BP

- Patient should be seated for 15 minutes and calm.
- Patient should not chew or smoke.
- The appropriate sized BP cuff should be used.

Severe Features of Preeclampsia

- sBP ≥ 160 OR dBP ≥ 110
- Renal insufficiency
- Pulmonary edema
- Thrombocytopenia (platelets $<100K$)
- Impaired liver function
- IUGR
- Cerebral or visual symptoms
- Severe, unremitting headache

*If random protein to creatinine ratio is between 0.15 and 0.3, obtain 24 hour urine protein.
 - If ≥ 300 , diagnosis is preeclampsia.
 - If <300 , diagnosis is gestational hypertension.

Gestational Hypertension (GH) Diagnostic Criteria

- BP $\geq 140/90$ measured on two occasions at least four hours apart.
- Only one elevated BP is needed to proceed with this guideline.

If patient < 20 weeks, refer to [Chronic Hypertension in Pregnancy guideline](#).

Full Maternal/Fetal Evaluation

- Preeclampsia labs (see box)
- Tests for fetal wellbeing (NST, BPP)
- Ultrasound for growth

Preeclampsia Labs

- CBC
- CMP
- Random urine protein to creatinine ratio

Any signs or symptoms of severe features?

Yes

Preeclampsia/Gestational Hypertension with Severe Features

Treat blood pressure per [Hypertension, Severe](#) guideline.

Admit and consult OB/GYN.

- Magnesium sulfate: 4 grams IV bolus over 20 minutes, then 2 grams/hour IV.
- Check preeclampsia labs.
- Monitor fetal wellbeing.
- Obtain OB ultrasound to evaluate for IUGR or oligohydramnios.
- Monitor for signs and symptoms of magnesium toxicity.

If possible, transfer to Anchorage.
If transfer not possible, consult HROB.

Gestational Hypertension

Random protein to creatinine ratio >0.3 ?

Yes

Preeclampsia

Consult HROB on call.

Consider inpatient monitoring versus transfer to Anchorage.

Yes

Outpatient Monitoring in Bethel

- Daily kick counts.
- Office visit 2 times per week.
- NST twice weekly.
- AFI and preeclampsia labs once a week.
- Ultrasound for growth every 3 weeks.
- Transfer care to Inpatient OB at 38 weeks for delivery or transfer to Anchorage.

No

Any signs or symptoms of severe features?

Yes

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
 Approved by Clinical Guidelines Committee 10/21/22.
 Click [here](#) to see the supplemental resources for this guideline.
 If comments about this guideline, please contact David.Compton@ykhc.org.




BP Technique

- Use the appropriate sized cuff after 5 minutes of rest.
- Patient should be sitting or semi-reclining (not fully reclining).
- Repeat with manual cuff after a minimum of 20 minutes if sBP ≥ 160 or dBP ≥ 110 .

Severe range BP is defined as sBP ≥ 160 or dBP ≥ 110 .

Village Management

- If patient is in a village, initiate the nifedipine regimen (see [Treatment](#) page) and activate medevac to Bethel.
- Do not delay treatment while awaiting disposition.
- If indicated, give steroids per [policy](#) and contact HROB for other recommendations.



This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 11/27/22.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact David_Compton@ykhc.org.



Severe range BP is defined as
sBP \geq 160 or dBP \geq 110.

Anti-hypertensive Medication Regimens

After two severe range BP's have been verified, choose one regimen.
Follow pathway until sBP < 160 and dBP < 110. At that time, return to first page for further management.

Give nifedipine (immediate release) 10 mg PO.

Repeat BP in 20 minutes.

If BP still in severe range, give nifedipine
(immediate release) 20 mg PO.

Repeat BP in 20 minutes.

If BP still in severe range, give nifedipine
(immediate release) 20 mg PO.

Repeat BP in 20 minutes.

If BP still in severe range,
give labetalol 40 mg IV.

Give labetalol 20 mg IV.

Repeat BP in 20 minutes.

If BP still in severe range,
give labetalol 40 mg IV.

Repeat BP in 20 minutes.

If BP still in severe range,
give labetalol 80 mg IV.

Repeat BP in 20 minutes.

If BP still in severe range,
give hydralazine 10 mg IV.

Give hydralazine 5 mg IV.

Repeat BP in 20 minutes.

If BP still in severe range,
give hydralazine 10 mg IV.

Repeat BP in 20 minutes.

If BP still in severe range,
give hydralazine 10 mg IV.

Repeat BP in 20 minutes.

If BP still in severe range,
give labetalol 40 mg IV.

If sBP or dBP continue to be in severe range after pathway is
exhausted, consult an obstetrician for additional recommendations.

Village Management


- If patient is in a village, initiate the nifedipine regimen and activate medevac to Bethel.
- Do not delay treatment while awaiting disposition.
- If indicated, give steroids per [policy](#) and contact HROB for other recommendations.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 11/27/22.

Click [here](#) to see the supplemental resources for this guideline.


If comments about this guideline, please contact David_Compton@ykhc.org.





Lab Criteria for ICP:

- TBA ≥ 10
- Cholic acid ≥ 3
- Total Bilirubin >1
- AST 2x normal
- ALT 2x normal
- Alkaline phosphatase (AP) >300




Intrahepatic Cholestasis of Pregnancy (ICP)

- Abnormal bile acid (BA) metabolism in pregnancy resulting in severe pruritus without rash.
- Mostly genetic etiology.
- 5% incidence in Yup'ik population.
- 5% incidence of stillbirth.
- MUST have elevated bile acids or LFTs.
- 40-70% recurrence in subsequent pregnancies.

Definitions

- ICP – Intrahepatic cholestasis of pregnancy
- Severe Pruritus – Any of the following signs or symptoms:
 - Excoriations
 - Loss of sleep due to pruritus
 - Scratching during appointment
- Pruritus of pregnancy – non-severe pruritus without elevated TBA
- TBA – Total Bile Acids
- Corrected TBA (cTBA) = TBA – ursodeoxycholic acid
- LFT – Liver Function Test



Non-invasive Prenatal Testing (NIPT)

Non-invasive prenatal testing is a way to detect fetal chromosome abnormalities from a maternal blood draw. Our current test is InformaSeq from LabCorp.

Risk Factors for Intrauterine Growth Restriction

Maternal Medical Conditions

- Hypertension
- Renal disease
- Restrictive lung disease
- Diabetes (with microvascular disease)
- Cyanotic heart disease
- Antiphospholipid syndrome
- Auto-immune disease


Other Factors

- Smoking and substance use and abuse
- Severe malnutrition
- Primary placental disease
- Multiple gestation
- Infections (viral, protozoal)
- Genetic disorders
- Exposure to teratogens

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 1/11/23. Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact David_Compton@ykhc.org.



Pregnant patient with symptoms suggestive of labor.



If activating a medevac for village labor

- Do not send a resident.
- Ensure that all Tiger Connect roles are covered by back-up physician.
- Notify ED Doctor On Duty since they are medical control.
- Gather [Village Delivery Backpack](#) from OB and Butterfly/iPad for US.
- Discuss with pediatrician the need to bring surfactant.
- Bring warm clothing (extra gear in peds call room under the bed), snacks, drinks, money, motion sickness medication, etc.
- Coordinate with pediatrician and plan to meet at LifeMed hangar at 3600 Tower Road. Tell LifeMed (LM) Dispatch if delayed more than 20 minutes.

Postpartum and Newborn Care

Mother: Check vital signs and fundal checks q15 min x 1 hour, then q30 min x 2 hours, then q1h x 3 hours. If all is well, may leave clinic to await travel with instructions to return immediately for return of bleeding or passing clots.

Baby: Check vital signs with axillary temperatures q30 minutes x 2 hours, then q1h x 4 hours. Low threshold to check blood glucose after first feed. Ensure vitamin K, erythromycin, and hepatitis B vaccine are given when able. If all is well, may leave clinic with instructions to return immediately for any concerns, especially trouble breathing, fast breathing, pauses in breathing, etc.

Tocolytics

1. Nifedipine (as BP permits)
0 min: 30 mg PO
90 min: 20 mg PO
180 min: 10 mg PO

2. Terbutaline 0.25 mg SC q5 min x 4 PRN
3. Ketorolac 60 mg IM or ibuprofen 800 mg PO

Steroids

- Betamethasone 12 mg IM q24h x 2 (preferred); OR
- Dexamethasone 6 mg IM q12h x 4

Antibiotic (if no allergy)

- Ceftriaxone 1 g IM

Uterotonics

1. Oxytocin 10 units IM, 10-40 units IV bolus (SRC only)
2. Misoprostol 800 mcg PO/PR/SL
3. Methergine 0.2 mg IM q2h

In the village


- Help the crew, follow their instructions, and expect to carry equipment.
- Assess fundal height and Leopold maneuvers; consider dating accuracy versus polyhydramnios if size greater than dates.
- If EGA < 34 weeks, perform a sterile speculum exam, obtain FFN, swab for GBS and GC/CT, and obtain urine sample for culture.
- If low risk for placenta previa (e.g., not noted on prior formal or Butterfly POCUS), check cervix after obtaining cultures.
- Make decision about disposition based on cervical exam, possible complications, and risk/benefit of travel.
- Discuss with HROB if any uncertainty about plan.
- Notify OB charge RN of plan as soon as possible from village clinic or Subregional Center (SRC) so they can prepare.
- If village delivery is anticipated, see [Village Deliveries \(Pediatrics\) Resource](#) for newborn care and preparation.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Guideline Committee 8/23/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact
William_Guerin@ykhc.org.



This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 1/11/23.
Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact David.Compton@ykhc.org.



Differential Diagnosis by Trimester

First

- Aneuploidy
- Fetal Anomaly

Second


- Aneuploidy
- Fetal Anomaly
- Preterm premature rupture of membranes (PPROM)
- Placental abruption
- Fetal growth restriction
- Amniocentesis
- Elevated maternal serum alpha fetoprotein


Third

- Preterm premature rupture of membranes
- Placental abruption
- Fetal growth restriction
- Utero-placental insufficiency
- Preeclampsia
- Maternal vascular diseases
- Fetal anomaly
- Post-term
- Suboptimal maternal hydration

Definition of Oligohydramnios

Amniotic Fluid Volume (AFI) < 5 cm at term
or
Maximum Vertical Pocket < 2 cm.





Bishop Score

Score	Dilatation	Effacement	Station	Position	Consistency
0	closed	0 – 30%	-3	posterior	firm
1	1 – 2 cm	40 – 50%	-2	mid-position	medium
2	3 – 4 cm	60 – 70%	-1, 0	anterior	soft
3	>5 cm	>80%	+1, +2		

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 3/13/23.


Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Ellen_Hodges@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline Postpartum Hemorrhage





BASICS

- Review the chart EVERY visit for incomplete lab or other required testing.
- Review the Problem List EVERY visit for needed testing or intervention items. See risk factors in PowerChart Maternity.
- Patient should see a Bethel provider or CHA/P monthly from first visit to 32 weeks.
- Patient should see a Bethel provider or CHA/P every two weeks after 32 weeks and then weekly at 36 weeks.
- If there is any question of EDC, see guideline or refer to HROB meeting for decision.

First Prenatal

NURSING/CASE MANAGER

- Order First Trimester Transvaginal OB Ultrasound (>6 weeks) for dating.
- Patient to initiate paperwork:
 - Residential Information Sheet.
 - Pregnancy Verification Sheet – use LMP if no EDC from ultrasound.
 - MSAFP and NIPT [consent](#) form.
 - FAS & Drug Assessment Screening questionnaire.
 - 36 Week BIB/Medevac Policy.
- Review TB screening status – patient MUST HAVE a negative Quantiferon or PPD prior to 36 weeks to stay at Prematernal Home. Place PPD if needed.
- Labs: urinalysis, urine culture, blood type and screen, HBsAg, Hepatitis C antibody, CBC, Rubella titer, HIV testing, treponemal testing, HgA1c, 25-OH vitamin D. Draw NIPT if >9 weeks and desired. GC/CT and trichomonas by self swab.
- Set up room for pelvic to do cervical cancer screening if due. (See [guideline](#).)
- Routine patient handouts: WIC handout.

PROVIDER

- Complete Antepartum Intake.
- Prenatal H&P and Prenatal Education.
- Chart review.
- Offer flu vaccine October through the end of the flu season.
- Discuss and sign BIB/Medevac Policy contract.
- Update the Problem List.
- Discuss post-partum birth control plan.
- Refer to HROB meeting if needed.
- Obtain baseline labs as needed.
- Update Risk Factor list in PowerChart Maternity.

PATIENT

- Go to the Medicaid office to file for Medicaid.
- Go to the WIC office to file for WIC.

15-21 Weeks

- If desired, MSAFP must be drawn between 15 and 21 weeks gestation.
- Review TB status.

20 Weeks

- Ultrasound to screen for anomalies: US OB anatomy and cervical length.
 - Only one is needed no matter where it is done.
 - Aim for 20 weeks.
 - If anatomy is incomplete, order US OB follow-up for the next visit to complete the anatomy exam.

24-28 Weeks

NURSING

- Labs: GST, CBC.
- Tdap after 24 weeks.
- GST – 50 g:
 - If result >140 mg/dL, schedule 2 hour GTT ASAP.
 - If the result >190, no GTT; refer directly to diabetes education.
- Attempt to keep the patient until the results of the GST are back.
- Review TB status. Draw Quantiferon if failed to have PPD read.

PROVIDER


- After 28 weeks, ask about preeclampsia symptoms.
- After 24 weeks, ask about preterm labor symptoms and IHCP symptoms.
 - Back pain.
 - Sudden increase in vaginal discharge.
 - Pelvic pressure.
 - Cramps/contractions.
 - Itching.
- Educate patient on fetal movement count.
- Discuss post-partum birth control plan. If patient desires sterilization procedure, complete [Consent for Sterilization](#) form.

32+ Weeks

- Offer RSV vaccine.

36 Weeks/BIB Date

- Labs: CBC, treponemal testing, HIV testing, GBS culture, GC/CT and trichomonas.
- Review TB status. Draw Quantiferon if status unknown.
- Schedule appointments to be seen each week by Bethel provider through 41 weeks.
- Ask about any symptoms of:
 - Rupture of membranes.
 - Preeclampsia.
 - Labor.
 - Itching.



This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/14/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Ellen_Hodges@ykhc.org.



Definition of Preterm Labor

Regular uterine contractions after 20 weeks gestation and before 37 weeks gestation which lead to a progressive **cervical change**.
Diagnosis of preterm labor requires a cervical exam.

Exams

Recheck cervix every two hours as long as there is concern for preterm labor.

Patient presents with signs and symptoms of preterm labor

Sterile speculum vaginal exam to assess for cervical dilation and obtain:
• Wet mount for bacterial vaginosis
• Fetal Fibronectin, GC/CT, trichomonas, GBS PCR, and cervical exam.
• Urinalysis

Patient with bacterial vaginosis?

Yes

Antibiotic Treatment

1. Metronidazole 500 mg PO twice daily x7 days.
- or
2. Clindamycin 300 mg PO twice daily x7 days.

No

Is cervix dilated ≥ 3 cm?

Yes

Refer to [Treatment of Preterm Labor](#) Guideline.

No

Fetal Fibronectin result?

Positive

- Give Bethamethasone 12 mg IM.
- Repeat dose in 24 hours.

Is ultrasound available?

No

Yes

Preterm labor present?

Yes

Refer to [Treatment of Preterm Labor](#) Guideline and OB policy on treating preterm labor.

No

Preterm labor present?

Yes

Follow up on next clinic day for cervical length US.

No

Yes

Is ultrasound available?

No

Is cervical length > 25 mm?

Yes

Routine Care

No

Is cervical length > 25 mm?

Yes

Routine Care

No

Contact HROB on call to discuss plan.

Yes

Stay in Bethel with clinic follow-up in one week.

Preterm Labor Symptoms

- Increased vaginal discharge
- Blood tinged mucus
- Low backache
- Pelvic pressure
- Menstrual-like cramps
- Intestinal cramping with or without diarrhea
- "Not feeling right"
- Loss of cervical mucous/"plug"

There is no need to treat contractions with tocolytics in the absence of cervical change.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/14/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Ellen_Hodges@ykhc.org.



Definition of Preterm Labor

Regular uterine contractions after 20 weeks gestation and before 37 weeks gestation which lead to a progressive **cervical change**. Diagnosis of preterm labor requires a cervical exam.


Contraindications to Tocolysis

- IUFD
- Lethal fetal anomaly
- Non-reassuring fetal assessment
- Severe IUGR
- Chorioamnionitis, relative
- Maternal hemorrhage with hemodynamic instability
- Severe preeclampsia or eclampsia
- PPROM
- Relative contraindication: delivery in Bethel seems inevitable.

Contraindications to Terbutaline

- Diabetes
- HTN
- Suspected placental abruption (relative)

If cervix is greater than 2cm dilated, consider early consult to ANMC On-Call OBGYN L&D for management and possible transfer discussion.



This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/14/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Ellen_Hodges@ykhc.org.



Signs & Symptoms of ROM


- Pool of fluid in vagina on exam
- Leak of fluid with valsalva
- Positive Ferning test
- Positive Amnisure test
- Oligohydramnios

Patient presents to care with complaint of leaking fluid from vagina while pregnant.

- Sterile speculum exam, GC/CT/TV, GBS PCR, GBS Screen, wet prep.
- Ferning Test or Amnisure.
- US OB Growth & AFI.

Village Care

- Health Aide should not do a vaginal exam but should contact Emergency RMT.
- Providers in SRCs and villages should do a speculum exam to look for signs of ROM and then call emergency RMT for consultation.



This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/13/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact David_Compton@ykhc.org.



For more information, see
[Rh Immune Globulin Work-up](#)
Policy & Procedure.

At first prenatal visit, check blood type and antibody screen in all patients.

Rh negative?

No

• Blood Type on newborn after birth only as indicated.
• No further testing of the patient for blood type.

Yes

• Note diagnosis on Problem List.
• Educate the patient.

At 28 weeks

• Obtain labs on RHIG Workup (Antenatal) Power Plan.
• Give RHIG (Rhogam®) 300 mcg IM after antibody screen.

When Patient is in Labor

Obtain blood type and antibody screen on admission.

After Delivery

• Obtain ABO and Rh on newborn.
• Obtain fetal screen on mother.

Newborn Rh
positive?

No

No further workup or
treatment.

Yes

Fetal screen
positive?

Yes

• Give the mother RHIG (Rhogam®) 300 mcg x2
doses (for total 600 mcg)
Immune Globulin.
• Send Kleinhauer-Betke (KB) test.
• Consult OB/GYN.
• Give additional doses based on KB results.

No

Give the mother RHIG (Rhogam®) 300 mcg IM.

Other Situations Which Require anti-D
Immune Globulin

- Miscarriage/Abortion
- Stillbirth
- Ectopic Pregnancy
- Maternal Trauma: consult OB/GYN.
- Threatened abortion
- Maternal hemorrhage in 2nd or 3rd trimester
- External cephalic version
- Amniocentesis

The dose is always 300 mcg at YKDRH due
to blood bank stocking.

This guideline is designed for the general use of most patients but may need to be adapted to meet
the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guidelines Committee 11/27/22.

Click [here](#) to see the supplemental resources for this guideline.


If comments about this guideline, please contact David.Compton@ykhc.org.



Criteria for Trial of Labor after Cesarean Delivery (TOLAC) in Bethel

- Only one previous cesarean delivery.
- No previous obstructive diagnosis.
- One vaginal delivery before or after cesarean delivery.

NOTE: Patients who require induction will not undergo TOLAC in Bethel.



Signs of Impending Uterine Rupture

- Marked variability
- Loss of fetal station
- Atypical pain or abdominal pain
- Vaginal bleeding

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 10/23/24.

If comments about this guideline, please contact Anthony_Markuson@ykhc.org.



Background

- Only 4-9% of those...labeled [penicillin-allergic] are currently allergic. It is important to identify those who are not allergic, because children mislabeled as penicillin-allergic have more medical visits, receive more antibiotic prescriptions, and have longer hospitalizations with more antibiotic-related complications.¹
- Up to 10% of children develop rashes while receiving antibiotics. Most are diagnosed...as allergic to the implicated antibiotic, and most continue to avoid the suspect antibiotic in favor of alternatives, which may be less effective, more toxic, and more expensive.²
- Do not label a patient as allergic to penicillin/amoxicillin unless he or she has true hives, anaphylaxis, or a life-threatening reaction. Please include photos of rashes in RAVEN.
- Children labelled as allergic to penicillin/amoxicillin often carry that label for the rest of their lives.
- Please consult a pediatrician with any questions.

Anaphylaxis

- Acute onset – several minutes to hours from exposure.
- Generalized hives, pruritis or flushing, swelling of lips/tongue/uvula, and at least one of the following:
 - Dyspnea, bronchospasm, stridor
 - Hypotension
 - Evidence of hypoperfusion of end-organs
 - Persistent crampy abdominal pain, and/or vomiting or diarrhea

Hives vs Viral Rash

- True hives are raised, itchy, larger than dime-sized, come and go, move around the body, and change shape and size. True hives are uncomfortable. Ask if the rash bothered the child.
- Keep in mind that many parents refer to any rash as "hives." Get a description every time.
- A viral exanthem is typically diffuse, fine, pinpoint red dots and can be dense, coalesced, larger raised lesions. The rash typically covers the face and chest but can cover the whole body. The rash typically worsens and takes days to clear.

NOTE: If amoxicillin is needed to treat a life threatening infection, consult Allergy & Immunology to discuss possible desensitization. Alaska Asthma, Allergy, & Immunology can be reached at (907) 562-6228.

References

1. Kelso JM. "Provocation challenges to evaluate amoxicillin allergy in children." JAMA Pediatrics 2016;170(6):e160282.
2. Mill C, et al. "Assessing the diagnostic properties of a graded oral provocation challenge for the diagnosis of immediate and nonimmediate reactions to amoxicillin in children." JAMA Pediatrics. 2016;170(6):e160033.

Patient labeled with a penicillin/amoxicillin allergy.

Review history. (See box.)

Was the reaction anaphylaxis (see box) or other life-threatening reaction (eg Stevens-Johnson syndrome, etc.)?

Yes

- Do not give drug or perform trial.
- Update chart, including the Problem List and a comment on the allergy.
- Refer to Allergy & Immunology at 5.

No

What was the reaction?

Rash

Vomiting and/or diarrhea without any other S/Sx anaphylaxis

Other

True hives, skin blistering/peeling, or mucous membrane involvement

Viral-appearing rash or other type of rash

- Do not give drug or perform trial.
- Update chart, including the Problem List and a comment on the allergy.
- Refer to Allergy & Immunology at 5.

- Not a true allergy.**
- Educate and perform Amoxicillin Trial (see box).
 - If patient/family refuses trial, update Problem List.
 - Offer future trial or refer to Allergy & Immunology at age 5 for amoxicillin allergy testing.

- Not a true allergy.**
- Educate and perform Amoxicillin Trial (see box).
 - If patient/family refuses trial, update Problem List.
 - Offer future trial or refer to Allergy & Immunology at age 5 for amoxicillin allergy testing.

Get more history. Consider pediatric consult.

This guideline is designed for patients that are extremely unlikely to be allergic to amoxicillin. There is no upper or lower age limit. Consult pediatrics with any questions.

Amoxicillin Trial Procedure²

Use AMB Amoxicillin Trial Power Plan.

1. Obtain VS. Perform physical exam, including lung exam. Have appropriate dose of EpiPen or epinephrine. Epinephrine (1 mg/mL): 0.01 mg/kg (or 0.01 mL/kg) IM Q5-15 minutes. Per AAP recommendations:
 - 7.5-25 kg: use EpiPen Jr (0.15 mg)
 - ≥ 25 kg: use EpiPen (0.3 mg)
2. Calculate weight-based dose of amoxicillin. Give patient 10% of that dose.
3. Place patient in nearby room and instruct caregiver to notify staff of any changes in status.
4. If no reaction by 20 minutes, give patient remaining 90% of weight-based dose of amoxicillin.
5. Observe another 60 minutes. If no reaction, check VS and physical exam. If all stable, discharge home with regular course of drug.
6. Give patient and family amoxicillin trial education sheet.
7. Update allergy alert in RAVEN. Click the allergy in the banner. Right click over the drug name and choose "cancel." On the "reason" drop-down menu, choose "OK on Retrial."

Notes:

- If patient is on a beta-blocker, stop this for 24 hours prior to procedure, if possible. Beta-blockers can interfere with treatment for anaphylaxis, if it occurs.
- Ensure that patients with asthma have optimal control prior to this procedure.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner. Approved by Clinical Guideline Committee 8/23/23. Click [here](#) to see the supplemental resources for this guideline. If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.





Definitions

- CBE: Clinical Breast Exam
- BSE: Breast Self Exam
- BIRADS: Breast Imaging Reporting and Data System, a system that scores findings on breast imaging

Recommendations for Screening at YKHC

- Mammogram: start at age 40 and rescreen Q2 years until age 74 depending on patient's health.
- Routine CBE or BSE can be done at patient/provider discretion but are not recommended.





Collection Process

- Ages 21-24 collect 1 Thinprep vial in the usual fashion.
- Ages 25-65 collect 2 Thinprep vials in the usual fashion.

Full recommendations on the ASCCP Mobile App, which can be found [here](#).

If History of Hysterectomy

- If indication was HSIL or cancer, screen Q3 years for 20 years.
- Otherwise, no screening required.

Abbreviations/Definitions

- LSIL: low grade squamous intraepithelial lesion
- HSIL: high grade squamous intraepithelial lesion
- ASCUS: atypical squamous cells of undetermined significance
- AGC: atypical glandular cells
- ASC-H: atypical squamous cells cannot exclude high grade


Source: Management of screening and diagnostic findings per ASCCP recommendations.
Consult OB-GYN as needed for more specific recommendations.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 5/28/24. Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact David_Compton@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline Colorectal Cancer Screening





This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 7/23/25.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact clinical_guidelines@ykhc.org.

Please review the [YKHC Low Risk Endoscopy Criteria](#) for questions about the above. If you have further questions related to if patient qualifies or not, Tiger Text or call CRNA on call.





This guideline is not intended for all outpatient pre-op appointments.

Resources for general pre-op visits for other types of surgery include:

- Documentation/requests of surgeon
- <https://www.aafp.org/pubs/afp/issues/2000/0715/p387.html>

Points to Consider

- Did patient already get scoped somewhere else?
- Is the time interval correct?
- Is there appropriate other workup & follow up underway for the condition being evaluated (e.g. dysphagia, anemia)?

It's not necessarily the pre-op provider's job during this visit to extensively evaluate the reason for the procedure. However, please ensure the patient knows why the procedure is being done and the plan for outpatient follow up.

Criteria for Anesthesia at YKHC

The patient must NOT have any of these:

- Age > 75 years.
- BMI > 50. BMI 45-49 needs clearance by anesthesia.
- Uncontrolled HTN (ambulatory SBP > 170 or DBP > 110).
- Congestive heart failure with EF < 40% and/or need for active management in the last year (i.e., diuretics, ER visits).
- Diabetes with A1c > 8 or need for insulin perioperatively.
- Suspected obstructed sleep apnea without sleep study/evaluation completed. (Patients with diagnosed OSA using CPAP are okay.)
- Asthma or COPD classified as severe (based on spirometry) or uncontrolled (based on symptom questionnaire or need for exacerbation treatment within 3 months).
- History of MI or CVA within the last 6 months or PCI within the last 12 months.
- Chronic or paroxysmal arrhythmia (including atrial fibrillation).
- Use of anticoagulant medication (not antiplatelet monotherapy).
- Implanted cardioverter-defibrillator or pacemaker.
- Decompensated cirrhosis.
- ESRD on dialysis.
- Poorly controlled seizures or new seizure diagnosis within 6 months.
- Active pulmonary TB.
- Functional status < 4 METs (unable to climb flight of stairs due to cardiorespiratory symptoms).
- URI, including strep pharyngitis or sinusitis, within 4 weeks; or, pneumonia within 8 weeks. (Time is from resolution of symptoms.)
- New cardiorespiratory symptoms such as chest pain, dyspnea, palpitations, syncope. These must be evaluated appropriately (e.g. Holter, stress test, etc) prior to clearance. Evaluation can be done by YKHC outpatient provider with specialty referral as indicated.

Please review the [YKHC Low Risk Endoscopy Criteria](#) for questions about the above. If you have further questions related to if patient qualifies or not, Tiger Text or call CRNA on call.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/23/25. Click [here](#) to see the supplemental resources for this guideline. If comments about this guideline, please contact clinical_guidelines@ykhc.org.



Additional Testing Requirements Prior to Endoscopy Clearance										
	CBC	BMP	CMP	A1c	HCG	PT/INR	EKG	CXR	MTB	Spirometry
Age > 60							X			
Female of childbearing age					X					
Hypertension		Within 6 months					X			
DM2	Within 6 months	Within 6 months		Within 6 mo if < 8 Within 3 mo if > 8			X			
Hx MI, CVA, CHF	X	X					X			
CKD 3/4	X	X								
Chronic liver disease	X		X			X				
Uncontrolled asthma/COPD								X		X
Clinical concern for active TB								X	X	
Hx bleeding disorder	X					X				
Smoker > 20 yrs	X									
Hx malignancy	X									
RA on MTX or DMARDs	X		X				X			


Medication Management		
Aspirin		Continue pre operatively
Diabetes agents	SGLT2	Stop 3-4 days prior
	GLP1	Hold one dose prior (daily or weekly dose as appropriate)
	All agents other than metformin	Stop one day prior
Oral iron		Stop 5 days prior
ACE or ARB		Hold morning of procedure
Take all other medicines as usual the morning of procedure		
NB: A patient still taking rifampin for TB/LTBI will not clear until off rifampin		

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/23/25.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact clinical_guidelines@ykhc.org.



For adults older than 75, the evidence is insufficient for recommendations. There may be greater absolute risk reduction, but greater risk of side effects.

Statin intensity and doses	
High	Moderate
Atorvastatin 40-80mg Rosuvastatin 20-40mg	Atorvastatin 10-20mg Rosuvastatin 5-10mg Simvastatin 20-40mg Pravastatin 40-80mg

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/11/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact clinical_guidelines@ykhc.org



Adult presenting for general preventive care:

- Annual screening for BP
- Review [Primary Prevention of ASCVD](#) guideline for other recommendations

Correctly measure blood pressure (see Box 1) and review prior BP readings in chart.

BP \geq 140/90?

No

BP \geq 130/80 and comorbidities?

No

Annual BP monitoring.
Encourage healthy lifestyle.

Yes

Yes

- Diagnose and discuss hypertension.
- Address modifiable contributors (see Box 2).
- Discuss lifestyle interventions (Box 3).
- Shared decision making and clinical judgment whether to start medication.

A note on definitions:

There is not consensus on a single reading that defines hypertension. Some groups use 140/90, others use 130/80. Others use ASCVD risk score or presence of comorbidities. Clinical judgment and shared decision making are the way to go.

Box 1: How to measure blood pressure

- Patient should be sitting for 5 minutes. Not fully reclined.
- Use the appropriate sized cuff.
- Verify elevated readings on other arm or by auscultation.

Box 2: Potentially modifiable contributors to hypertension

- Tobacco use
- Obesity
- High sodium diet
- Excess alcohol use
- Stress
- Poor sleep
- Certain medications: OCPs, NSAIDs, steroids, decongestants, some antidepressants

Box 3: Lifestyle interventions for hypertension

- Quit tobacco and alcohol
- Exercise
- Weight loss
- DASH or Mediterranean diet
- Mindfulness based stress reduction

Box 4: Secondary causes of hypertension, and/or conditions that should be co-managed with specialist

- Primary aldosteronism
- Renovascular hypertension
- Obstructive sleep apnea
- Congestive heart failure
- Chronic kidney disease

Pharmacotherapy?

- Review problem list. Should this person's BP be managed by specialist? See Box 4.
- Check BMP, CBC, A1c, lipids, TSH, ECG, UA. Manage abnormalities as indicated.
- Choose a medicine in table 1.

Reassess BP monthly until at goal, then every 3-12 months.

Not at goal?

- Readdress modifiable contributors and lifestyle interventions.
- Assess adherence and barriers/side effects.
- Add agent from another drug class.
- Titrate to max dose tolerated.

Not at goal on 2+ agents?

- Consider secondary causes of hypertension if not done already (see Box 4).
- Consider consult/referral to nephrology or internal medicine.

Severe hypertension \geq 180/120

- Confirm measurement using other arm, different cuff, auscultation
- Inquire for emergent symptoms: Worst headache of your life, acutely altered mental status, acute chest or back pain, stroke symptoms. If present, contact Emergency RMT or ER physician.
- If there are not symptoms of hypertensive emergency, continue to follow this guideline, with closer interval follow up during med titration.

Table 1: Initial pharmacotherapy for hypertension

Class	Drugs and doses	Notes
ARB/ACEi	Losartan 25-100mg Lisinopril 2.5-40mg	<ul style="list-style-type: none"> • Must be on birth control if any chance of pregnancy • Preferred if diabetic, CKD • Check BMP 1 week after starting and every 3-12 months
Calcium channel blocker	Amlodipine 2.5-10mg	<ul style="list-style-type: none"> • May have more potent effect on BP • Common side effect leg swelling
Thiazide diuretic	Chlorthalidone 12.5-25mg HCTZ 12.5-25mg	<ul style="list-style-type: none"> • Warn patients about increased urination • Check BMP 1 week after starting and every 3-12 months

Note: If untreated BP is greater than 160/90, consider starting two agents simultaneously, e.g. ARB + CCB.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.


Approved by Clinical Guideline Committee 8/2/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact clinical_guidelines@ykhc.org.

“Lead, State” and choose capillary or venous.

- At 1st and 2nd year well check.
- If child <6 years with no prior screening or unexplained developmental delays.
- Screen for anemia concurrently.



NOTE: For capillary collections, fingers need to be thoroughly cleaned with soap and water as well as alcohol wipes. Contamination with dirt is the most common reason for high lead levels.

- Mining lead, zinc, silver, or gold ore
- Lead paint in homes or buildings built before 1978
- Firearms and ammunition
- Shooting ranges
- Game meat shot with lead ammunition
- Fishing weights
- Leaded aviation gas
- Marine paint
- Soldering, welding, or craft-making
- Pica or "mouthing" (eating dirt)
- Imported household objects
- Lead or brass pipes/faucets
- Batteries and automobile repair sites


- If venous level >5 or two recorded capillary levels >3.5, Alaska Department of Health will call family to discuss sources.
- For any level >20 or if level remains high after a month, may consult YKHC Office of Environmental Health (543-6420) to consider a home visit.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by Clinical Guideline Committee 9/16/24. Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact Justin Willis@ykhc.org.



General

- Nirsevimab (brand name Beyfortus™) is a monoclonal antibody to prevent RSV.
- Studies show that nirsevimab prevents ~80% of hospitalizations when given to infants <8 months.
- The season will be September 13, 2024 through March 31, 2025. This may be extended if RSV is still being seen at the end of this period.
- At YKHC, dose will be given during the birth hospitalization at 24 hours of life whenever possible.
- NOTE: The AAP recommends that nirsevimab not be given to infants whose birthing parents have received the vaccine (Abrysvo™) during that pregnancy more than 14 days prior to delivery unless the infant is at "substantially increased risk for severe disease." Local experts have agreed that Alaska Native children are at increased risk of severe disease.
 - Thus, Alaska Native infants may be given nirsevimab during the birth hospitalization regardless of birthing parent vaccine status.
 - If the birthing parent received the vaccine during a previous pregnancy but not the most recent pregnancy, the recommendation is to give the infant nirsevimab.
 - Recommendations are still evolving for infants without risk factors for severe disease. If the birthing parent of a non-Alaska Native infant has received Abrysvo™ >14 days prior to delivery, that infant does not meet criteria for nirsevimab at birth but may benefit from administration at 3-6 months of life. This guideline will be updated as recommendations become available.



Allocation of Doses if Limited Supply

In the event of a shortage, nirsevimab will be allocated to highest risk patients first through a tier system. Tiers will be communicated by email whenever there is a change.

- Tier 1:** Lowest supply - only give to highest-need patients: infants <8 months, older patients designated high-risk by previously used palivizumab criteria (this will be noted on the patient's Problem List).
Tier 2: Medium supply or more supply expected soon - give to all Alaska Native infants <12 months as well as all Tier 1 patients.
Tier 3: Full supply - give to any infant who meets criteria per this guideline.

Risk Factors for Severe Disease

Children who meet the following criteria may receive a dose of nirsevimab between 8-19 months.

- Children with chronic lung disease of prematurity who required medical support (chronic corticosteroid therapy, diuretic therapy, or supplemental oxygen) during the 6-month period before the start of their second RSV season.
- Children with severe immunocompromise.
- All American Indian and Alaska Native children – as supply allows. Non AI/AN children without risk factors do not qualify for a second season dose.
- Cystic fibrosis patients who have either 1) manifestations of severe lung disease (previous hospitalization for pulmonary exacerbation in the first year of life or abnormalities on chest imaging that persist when stable) or 2) <10th percentile weight-for-length.

Palivizumab (Synagis®)

- Non-Alaska Native children between 8-23 months with chronic lung disease or congenital heart disease may meet criteria for palivizumab.
- See the [Alaska State Epidemiology Bulletin for the 2022-2023 criteria for palivizumab](#) for more details.

References and Resources


- [Alaska State Epidemiology Bulletin for Nirsevimab](#)
- [Alaska State Epidemiology Bulletin for RSVPreF Vaccine \(Abrysvo\)](#)
- [AAP FAQ for Nirsevimab](#)

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.


Approved by Clinical Guideline Committee 9/16/24.

Click [here](#) to see the supplemental resources for this guideline.


If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by MSEC 11/7/23.
Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact clinical_guidelines@ykhc.org.



Agent	Route of administration	Frequency and duration	Efficacy	Contraindications	Adverse effects
Alendronate	PO (first thing in the morning on empty stomach with glass of plain water, then sit upright x 30 min)	Daily or weekly for 3-5 years. Followed by 2 year break.	<ul style="list-style-type: none"> Reduces risk of serious fracture by 20-70% over 10 years. Number needed to treat is between 17 and 100. 	<ul style="list-style-type: none"> CrCl < 35 ml/min Severe esophagitis or esophageal dysmotility (not just GERD) Inability to sit upright x 30 min 	<ul style="list-style-type: none"> Osteonecrosis of the jaw estimated risk 1-5 in 10,000. Severe esophagitis. Atypical femur fracture. <p>Note: in RCTs there was no significant difference in adverse effects vs placebo.</p>
Zoledronic acid	IV infusion	Annually for 3 years, followed by 2 year break	Same as alendronate, but may be enhanced by increased adherence	CrCl < 35ml/min	<ul style="list-style-type: none"> Same as alendronate Potential difficulty/delay in emergency dental care Infusion reaction
Denosumab	Subcutaneous injection	Every 6 months, discontinuation must be immediately followed by bisphosphonate	Same as bisphosphonates.	Inability to complete Q6 month visits.	<ul style="list-style-type: none"> ONJ estimated risk 3 in 1,000 Rapid bone density loss upon discontinuation if not followed immediately by bisphosphonate



References

- [American Society of Endocrinologists](#)
- [American College of Physicians](#)
- [AAOMS Position Paper on MRONJ 2022](#)

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by MSEC 11/7/23.

Click [here](#) to see the supplemental resources for this guideline.
If comments about this guideline, please contact clinical_guidelines@ykhc.org.



Osteoporosis Treatment Options and Consent

I am considering taking either Zoledronic Acid (Reclast) or Denosumab (Prolia) to lower my risk of serious fracture.

My provider has discussed the following risks, benefits, and alternatives with me.

Purpose of treatment	<ul style="list-style-type: none"> • Reduce risk of serious fracture such as hip, pelvis, or spine fracture. Elders living in the YK Delta might have a higher risk of fracture than others in the US. This might be a 3-4 percent risk over ten years, or a 1 in 25 chance. • Serious fractures can lead to long hospital stays, loss of independence, need for nursing homes, and even increased risk of death.
Benefits of treatment	<ul style="list-style-type: none"> • All of the available medicines at YK are effective at preventing fractures. • All of the medicines could lower the risk of serious fracture by between 20 and 70 percent. • This might reduce a person's risk of serious fracture from 3-4 percent to as low as 1 percent.
Risks of treatment	<ul style="list-style-type: none"> • The medicine isn't effective for me and I have a serious fracture anyway. • Osteonecrosis of the jaw. This means exposed jaw bone that is painful or infected, and could result in multiple surgeries or chronic pain. This is most likely to happen after a pulled tooth or dental procedure. But it can happen spontaneously. The risk of that side effect could be between 0.01% (1 in 10,000) and 0.3% (1 in 300). The risk can be lowered by regular dental care and good oral hygiene. • Other rare side effects: atypical femur fracture, esophagitis, allergic/infusion reaction • If the medicine is not taken exactly as the provider instructs, it may not be effective. One of the medicines (Denosumab) will actually increase the risk of fracture if not taken every 6 months.
Alternatives to treatment	<ul style="list-style-type: none"> • Alendronate This is the same type of medicine as Zoledronic Acid. It is taken as a pill once a week and the person must take it first thing in the morning on empty stomach with glass of plain water, then sit upright for 30 minutes. It is about as effective as Zoledronic Acid, if people can remember to take it correctly. It also has a risk of osteonecrosis of the jaw, but if there are dental problems it is easier to stop this medicine than Zoledronic Acid. • It is very important for all adults to stay active, get at least 90 minutes of weight bearing physical activity per week, consume enough calcium and vitamin D, and learn about preventing falls in the home.

I would like to start treatment for osteoporosis with Zoledronic Acid or Denosumab.

I understand the purpose, benefits, risks, and alternatives that my provider reviewed with me.

Please choose one:

☐ I would like to start the treatment right away. This may have more benefit preventing fractures sooner, but also more risk if I have a dental emergency.

☐ I would like to see a dentist before starting treatment. I understand this may result in a delay before I can start the medicine.

Patient or Surrogate Decision-Maker/POA Signature: _____

Printed Name: _____

Date and Time: _____

Provider Signature: _____

Printed Name: _____

Date and Time: _____

All patients should be verbally referred and encouraged to see a dentist even if they want to immediately start treatment. Dental takes walk in patients every day. Providers are also welcome to consult with a dentist before counseling a patient on their dental risks.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved by MSEC 11/7/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact
clinical_guidelines@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Treatment Protocol Pre-Anesthesia Management

Age	Hb/Hct	Coags	Lytes	BUN/Cr	Glucose	LFTs	EKG	CXR	T&S
30 months – 59 years	No routine testing needed in this age group.								
60 – 74 years							X		

Disease	Hb/Hct	Coags	Lytes	BUN/Cr	Glucose	LFTs	EKG	CXR	T&S
Hypertension			X				X		
Card – moderate	X		X	X			X		
Smoker > 20 years	X								
Malignancy	X								
Lymphoma	X (CBC)							X	
Hepatic	X	X	X			X			
Renal	X	X	X	X					
Bleeding	X (CBC)	X							
Diabetes			X	X	X		X		
Expected blood loss	X								X

Medication	Hb/Hct	Coags	Lytes	BUN/Cr	Glucose	LFTs	EKG	CXR	T&S
Diuretic			X	X					
Antihypertensive			X	X			X		
Cardiac medication			X	X			X		
Steroid			X		X				
Anticoagulant	X	X							

Other

Urine hCG: obtain within 48 hours of surgery in women of childbearing age (13-50).

Drug Levels: draw level on all patients on digoxin or phenytoin.

CXR: obtain if recent change in sputum quality or color, pneumonia in past three months, chronic home oxygen use, planned intrathoracic surgery, or if exam reveals rales, rhonchi, or wheezes.

Surgical Risk Screening for Elective Procedures (including endoscopy)

- Patients who are not to be scheduled at YKHC:
 - Patients with BMI > 45.
 - Severe obstructive sleep apnea.
 - Patients with pending cardiology, pulmonology, or sleep study referrals.
 - Patients younger than 30 months.
 - Patients older than 75 years.
 - Medically unstable patients (for example, uncontrolled diabetes mellitus, uncontrolled hypertension, etc.).
- Preventative antibiotic therapy will be administered within one hour prior to skin incision per protocol pre-operatively based on procedure type and patient's allergies unless otherwise ordered by physician.
- DVT/VTE prevention methods will be implemented using **SCIP Mechanical Prophylaxis Protocol** unless contraindicated or otherwise documented in orders by physician.

Diabetes Management

- Oral agents: Discontinue SGLT2 inhibitors 3-4 days prior to surgery. Discontinue all other oral agents the evening prior to surgery, except Metformin can be taken. No oral agents except Metformin the morning of surgery.
- For patients who take insulin, consult pharmacy.**
 - For patients who take long acting insulin in the morning, take 50% dose of NPH insulin or 75% dose of long-acting insulin (lantus) the morning of surgery.
 - For patients who take long acting insulin at night, take 75% dose of NPH or lantus the night before surgery.
 - For patients who take short acting insulin (regular, aspart), stop this insulin when fasting begins.
- Consume apple or cranberry juice up to two hours prior to arrival to surgery if insulin was given.
- For insulin pumps, set to basal rate and continue throughout pre-operative period.
- Upon arrival to Holding Area, obtain glucose level. Anesthesia will treat results.

Please send a message via Tiger Connect to "OR CRNA on call" with any questions about patient selection, etc.

See YKHC Policy & Procedure on [Patient Selection Criteria for Ambulatory Surgery](#).

This protocol is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.
Approved 6/6/22.

If comments about this protocol, please contact
Jennifer_Lent@ykhc.org.



NPO Guidelines

1. All patients are to be NPO after midnight the night before the procedure. Additionally, patients undergoing endoscopy or with delayed gastric emptying will receive more extensive NPO instructions.
2. Patient may brush his/her teeth but should not swallow toothpaste.
3. Gum and candy of any type are not allowed.
4. All patients will be allowed to eat a full, regular diet (solids) up to eight hours prior to surgery. Patients going to the OR at 0730 who were NPO after midnight are considered to meet this standard.

Estimated Energy Requirements for Various Activities, Based on Duke Activity Status Index*

1 MET	Can you...	
		...take care of yourself?
		...eat, dress, or use the toilet?*
		...walk indoors around the house?
		...walk one or two blocks on level ground at 2-3 mph (3.2-4.8 kph)?
< 4 METs	Can you...	
		...do light work around the house, such as dusting or washing dishes?
≥ 4 METs	Can you...	
		...climb a flight of stairs or walk up a hill?
		...walk on level ground at 4 mph (6.4 kph)?
		...run a short distance?
		...do heavy work around the house, such as scrubbing floors or lifting or moving furniture?
		...participate in moderate recreational activities, such as golf, bowling, dancing, doubles tennis, or throwing a baseball or football?
≥ 10 METs	Can you...	
		...participate in strenuous sports, such as swimming, singles tennis, football, basketball, or skiing?

* MET = metabolic equivalent

Adapted from J AM Coll Cardiol, with permission from Elsevier.

This protocol is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved 6/6/22.

If comments about this protocol, please contact
Jennifer_Lent@ykhc.org.



Initial Visit

- ☐ Review NICU/Nursery course and summarize highlights in note. Update Problem List. Make patient CPP.
- ☐ Enter birth weight and gestational age so that RAVEN Growth Chart will correct for gestational age. (Go to Growth Chart → Enter New → Measurement → Preterm Growth Chart: Change date to DOB, enter gestational age at birth, and enter birth weight.)
- ☐ Check height and weight. Do not discharge to village if insufficient weight gain (at least 25 grams per day for 45 consecutive days), temperature <97.7, or rising bilirubin level.
- ☐ Check bilirubin level if appearing jaundiced. Follow [Jaundice in a Baby <4 weeks](#) guideline and peditools.org.
- ☐ Ensure infant is receiving fortified formula (ie Neosure) if discharged from the NICU on it. Infant should remain on this formula until 6 months corrected gestational age. Ensure that family has formula delivery set up from home health company. Contact Pediatric Case Managers if not.
- ☐ Place order: "Refer to Family, Infant, Toddler Program."
- ☐ If born <34 weeks, place order: "Refer to Child Family Developmental Services External", CFDS Sub-Specialty drop down "NICU Graduate Clinic."
- ☐ Place referrals for any subspecialists per NICU/nursery discharge summary and prescriptions for all medications.
- ☐ Check POC hemoglobin. If less than NICU discharge hemoglobin, consult a pediatrician.
- ☐ Write Vitamin D prescription with 11 refills and ensure receiving 800 IU Vitamin D supplementation. (Poly-vi-sol with iron has 400 IU of Vitamin D per mL)
- ☐ Write iron prescription with 11 refills and ensure receiving iron supplementation (Poly-vi-sol or iron polysaccharide). Needs 2 mg/kg iron supplementation for first year of life. (Note: Poly-vi-sol with iron contains 11 mg of iron per mL.)

To consult the pediatrician on call, send a message through Tiger Connect to Peds Wards on Duty.

General Information

- Soy milk formulas should not be given to preterm infants.
- Physiologic reflux is more common in preterm infants. There is no evidence to support the use of gastric acidity inhibitors. H₂ blockers and PPIs are associated with gastroenteritis, pneumonia, and bone fractures.
- Catch up growth of premature infants occurs for head first (3-8 months), then weight, then length.
- Recommend every member of the household is up to date on pertussis vaccine, COVID, and seasonal influenza vaccines to protect these high-risk infants.

Criteria for Referral to Child Family Developmental Services (CFDS) Birth to Three High Risk Clinic

This is a specialty clinic in Anchorage that follows high-risk infants.

- Birth weight (BW) <1500 grams.
- Gestational age <34 weeks.
- Cardiorespiratory depression at birth
- Apgar score <5 at 5 minutes
- Prolonged hypoxia, acidemia, hypoglycemia, or hypotension requiring pressors.
- Persistent apnea requiring medication.
- Oxygen support for >28 days and X-ray findings consistent with chronic lung disease.
- Extracorporeal membrane oxygenation (ECMO)
- Persistent pulmonary hypertension of the newborn (PPHN)
- Seizure activity
- Intracranial pathology, including intracranial hemorrhage, periventricular leukomalacia, cerebral thrombosis, cerebral infarction, or any developmental/central nervous system (CNS) abnormality
- Other neurological insult, including hypoxic ischemic encephalopathy (HIE), kernicterus, sepsis, CNS infection
- Confirmed prenatal exposures to alcohol, methamphetamines, opiates, or Suboxone.

All Subsequent Visits until Child is 24 Months Old

- ☐ Review and update Problem List.
- ☐ Assess growth based on corrected gestational age. Consult pediatrics if: there is a need to increase/decrease feeding calories, head circumference growth >1.25 cm/week, or infant is crossing major percentile lines.
- ☐ Check POC hemoglobin at all well visits to monitor for anemia of prematurity. Needs at least 2 mg/kg iron supplementation for first year of life. (Note: Poly-vi-sol with iron contains 11 mg/mL of iron.)
- ☐ Review feeding, sleep, and development in detail. Ensure meeting developmental milestones for corrected age.
- ☐ Check on FIT involvement. If family has not been contacted by FIT, reach out to Peds Wards on Duty, who will contact the FIT liaison.
- ☐ Give all vaccines per routine schedule based on chronologic age.
- ☐ Administer ASQ at **9 months**, **18 months**, and **24 months** chronologic age.
- ☐ At 9 months chronological age, ensure infant is scheduled with Audiology. (All former premature infants should have their hearing screened at 9-12 months.)
- ☐ Administer **MCHAT-R** at 18 months and 24 months chronologic age. If score of ≥3 (fail), schedule with pediatrics.
- ☐ Ensure specialty appointments/referrals have been made.
- ☐ If on caffeine, alter dose based on [Caffeine Protocol, Post-NICU Discharge Resource](#).
- ☐ Ensure infant has received nirsevimab (Beyfortus). If unavailable, send email to YKHCsynagis@ykhc.org so infant may be screened for palivizumab (Synagis).
- ☐ Ensure receiving Vitamin D 800 IU supplementation for first year of life (Poly-vi-sol with iron has 400 IU of Vitamin D per mL).


Please see the [Care of Late Preterm Newborns](#) guideline for information about late preterm babies who were cared for at YKDRH and were not admitted to a NICU.

Documentation: Use the autotext ".pednicugrad" for a summary of this checklist for charting purposes.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 3/11/24..

If comments about this guideline, please contact Justin_Willis@ykhc.org.



Aspirin	Age 40-59	Calculate 10 yr ASCVD risk . If risk > 10%, recommend shared decision making. Evidence supports a moderate likelihood that aspirin will prevent non-fatal MI or CVA. There is no evidence of impact on mortality or colorectal cancer. There is increased risk of bleeding (GI, hemorrhagic stroke), of uncertain degree.
	Age > 60, or 40-59 with < 10% ASCVD risk	Recommend against initiation of aspirin for primary prevention.


Statins	Age 40-75 with 1+ risk factor (HTN, dyslipidemia, DM, tobacco use) and 10 yr ASCVD risk > 10%	Recommend statin. There is evidence of moderate mortality benefit. Refer to hyperlipidemia guideline for dosing options.
	Age 40-75 with 1+ risk factor and 10 yr ASCVD risk 7.5 – 10%	Recommend shared decision making. There is evidence of a small mortality benefit and not more than a small risk of harm. Refer to hyperlipidemia guideline for dosing options.
	Age > 76 with no hx CVD and not already taking statin	Inadequate evidence to make recommendation. Inadequate evidence of benefits and harms in this age group.
	LDL > 190	These patients are excluded from above recommendations. Refer to hyperlipidemia guideline .

Hypertension	Annual screening for all adults > 40 and adults 18-40 with increased risk (obesity, Black persons, prior high normal BPs, tobacco use, others.) USPSTF reports insufficient evidence to recommend cutoff at 130/80 vs 140/90. Recommend confirming with ambulatory BP monitoring if possible. Can refer to this State of Alaska program for home BP cuff. See hypertension guideline for medication recommendations.
--------------	---

Tobacco cessation	All adults who use tobacco	Strong recommendation with strong certainty to encourage cessation. "Refer to Nicotine Control- Internal." There is convincing evidence to support to use of NRT, bupropion sustained-release, and varenicline. There is convincing evidence to support combining two types of NRT (e.g. gum/lozenge + patch) or combining NRT with bupropion. There is insufficient evidence of benefit for e-cigarettes.
-------------------	----------------------------	---

Obesity	Adults with BMI > 30 on annual screening	USPSTF recommends referral to multicomponent behavioral intervention for weight loss. At YK, consider: referral to Diabetes nutrition counseling if diabetic, referral to State of Alaska program , or regular clinic visits for discussion of safe and sustainable programs, goal setting, and follow through.
---------	--	--

Diabetes	Adults >35 with BMI > 25 and/or risk factors (hx gestational diabetes, PCOS, family history, dietary concerns)	Screen with A1c every 3 yrs if value remains normal (< 5.7). There is moderate certainty evidence of moderate mortality and morbidity benefit. If A1c > 5.7, refer to Diabetes guideline .
----------	--	---



Symptom Severity Classification for this Guideline

- Mild: no fever, <3 days of symptoms
- Moderate: prolonged fevers and bedrest, hospitalization not required, no abnormal cardiac testing throughout course
- Severe: hospitalized, abnormal cardiac testing, or MIS-C

Note: Providers may use their clinical judgment and perform an ECG if cardiac concerns not addressed by this guideline.

Phone Numbers

Seattle Children's Pediatric Cardiology of Alaska (located in Anchorage):

- Phone: (907) 339-1945
- Fax: (907) 339-1994

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 9/25/23.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact Leslie_Herrmann@ykhc.org.



Table 1: Alcohol Hangover (F10.120)

- Poorly defined but universally understood; occurs the morning after a night of heavy drinking.
- In general, starts <12 hours after a binge of <24 hours.
- Sx: fatigue, thirst, headache, nausea, concentration problems, apathy, loss of appetite, dizziness, vomiting, heart pounding/racing.
- Requirements: HR<130, BP<160/100, RR<24, T<100.4, ambulatory, GCS=15, appropriate history, no tremor, no anxiety, no significant comorbidities.

Table 2: Inpatient Criteria

- CIWA>12, despite treatment with PB/BZD.
- Requiring high-dose sedatives or IV infusion to maintain CIWA<12.
- GCS<8 or hemodynamic instability.
- Persistent hyperthermia (T>100.4 F).
- Respiratory insufficiency (hypoxia, hypercapnia, etc.).
- Marked acid-base disturbance.
- Cardiac disease (heart failure, arrhythmia, evidence of ischemia, etc.).
- Severe electrolyte abnormality.
- Severe renal insufficiency or requiring high volume fluids.
- Evidence of rhabdomyolysis.
- Potentially serious infection (PNA, wounds, etc.).
- Severe GI pathology (GI bleed, pancreatitis, etc.).
- Severe psychomotor agitation (high risk to self or others, gravely disabled, etc.).
- Evidence concerning for Wernicke-Korsakoff Syndrome (oculomotor dysfunction, ataxia, severe malnutrition).
- Withdrawal despite very elevated serum ethanol.

Table 3: Phenobarbital Contraindications

Absolute: Hx allergy, adverse reactions, or porphyria
Relative: current significant sedative level (including EtOH, BZD, or anti-psychotics)

Table 4: Phenobarbital (PB) Protocol

- Phenobarbital 260 mg IV then phenobarbital 130 mg IV every 30-40 minutes until CIWA score ≤ 12. No discharge meds.
OR (for very large/small patients)
- Phenobarbital 4 mg/kg IV (rounded to nearest 130 mg) then phenobarbital 2 mg/kg IV every 30 minutes until CIWA score ≤ 12. No discharge meds.
OR
- Either of the above via IM injection, with subsequent doses very 60-90 minutes.

Adverse Effects:


- Transient asymptomatic hypotension
- Transient ataxia
- Transient lethargy

Table 5: Discharge Criteria

- No inpatient criteria present (Table 2).
- CIWA score <12.
- Awakens to voice or light touch.
- Oriented with no delirium.
- Ambulatory without assistance.
- Taking liquids without vomiting.
- No co-administered sedatives/anti-psychotics.
- No seizures after treatment.
- Likely compliant with important outpatient medications (including antibiotics, etc.).

Please see the Wiki for more information:

[Alcohol Withdrawal in the YK Delta](#)
[Phenobarbital for Alcohol Withdrawal](#)



This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 10/21/22.

If comments about this guideline, please contact Clinical_Guidelines@ykhc.org.



Concern raised by caregiver or teacher for Attention Deficit Hyperactivity Disorder (ADHD) in child aged 6-17 years

Distribute initial Vanderbilt evaluation forms for caregiver and teacher to complete. Click [here](#) for forms.

For children <6 years, consider referral to CFDS and/or psychiatry group.

Schedule appointment with provider trained in ADHD evaluation and management.

Initial Vanderbilt evaluation forms completed by caregiver and teacher. Forms reviewed by provider before, at, or after appointment.¹ Provider conducts medical evaluation at appointment.

Vanderbilt positive for ADHD with comorbid psychiatric conditions

- Address any medical issues (sleep, etc.).
- Refer to psychiatry group for further evaluation and management.

Complicated Patients

Refer complicated patients to psychiatry group. Patients on more than one psychiatric medication may require more frequent visits.

Contact

- If questions about pediatric management, use Tiger Connect Team "Pediatric Outpatient."
- If a patient has run out of medications, send a Tiger Text to role "Psych Med Refills Only (Not for Patients)."
- To consult a psychiatrist, send a message to Tiger Connect role "Psychiatry Attending Physician (Not for patients)."

Vanderbilt positive only for ADHD and physical/situational/organic conditions excluded. No cardiac contraindications (see box).

Start trial of low-dose stimulants for one month and refer to Behavioral Health for concurrent counseling to achieve optimal results.

Distribute follow-up Vanderbilt evaluation forms to be completed by caregiver and teacher after two weeks of new medication or dose.

Schedule follow-up appointment within 30 days of starting medication or changing dose. Review follow-up Vanderbilt forms: do they indicate acceptable control of ADHD symptoms with reasonable side effect profile?

Yes

Monthly weight checks in village clinic or Bethel for three months after any dose adjustment.

May refill stimulant prescription for up to three months at a time once on a stable dose without significant adverse effects.³

- Follow-up ADHD appointment for medication review (with completed Vanderbilt evaluation forms) every 3 months.
- May alternate face-to-face visits with VTC appointments. Must check height, weight, and VS every 6 months.
- May refill one additional month while awaiting an appointment if necessary.

Vanderbilt negative for ADHD

- Perform full WCC.
- Address any medical issues (sleep, etc.).
- Consider checking lead and vitamin D levels and/or referring to Audiology and Optometry.
- Consider referral to Behavioral Health for counseling or to Child and Family Developmental Services (CFDS) clinic in Anchorage for further neurodevelopmental testing.²

Review results with caregiver (may be by phone). Titrate dose as indicated. May consult psychiatrist via Tiger Connect.

No

If any of the following are present, refer to cardiologist prior to starting stimulants:

- Hx congenital heart disease or previous heart surgery
- FHx sudden death suggesting cardiac disease under 40 in a first-degree relative
- SOB on exertion compared to peers
- Syncope on exertion or in response to fright or noise
- Palpitations that are rapid, regular, and start and stop suddenly; fleeting occasional "bumps" do not need investigation
- Chest pain suggestive of cardiac etiology
- S/Sx heart failure
- Heart murmur not c/w benign process
- If BP consistently above the 95th percentile for age and height

Footnotes

1. Scan completed Vanderbilt forms into MultiMedia Manager under "Continuity of Care."
2. To refer to [CFDS](#) or other private psychologist: use "Refer to Other External" order and send a message to the case manager to process the referral.
3. E-prescribe three separate 30 day prescriptions after checking Alaska PDMP. Include the month the medicine is to be filled in the comments or special instructions section.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/14/23.

Click [here](#) for the supplemental resources for this guideline.


If comments about this guideline, please contact Clinical_Guidelines@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Protocol

Care of an Agitated or Aggressive Patient on Inpatient or DES



Code Gray

- Code Gray team is activated by pressing button on panel located at the nursing station. This will activate an overhead page on the hospital PA system.
- Code Gray team will include all available security personnel, behavioral health clinician, charge nurse (or designee), and if possible another nurse. Medical provider will attend if able. Goal is a minimum of six team members at all Code Gray events.
- Charge nurse will determine when patient is calm enough for Code Gray staff to leave unit.
- BH clinician and bedside nurse will document incident in detail, including all interventions attempted, if meds were given, patient response and behaviors, actions if restraint and/or seclusion were applied, and timing of events.

Medications to Treat a Combative Patient (Use "MED Behavioral Health IP Admission" Power Plan.)

- Olanzapine 5-10 mg IM/PO Q10-30 minutes *prn* up to max 24 hour dose 60 mg.
- Haloperidol 2.5-10 mg IM/PO Q10-30 minutes *prn* up to max 24 hour dose 100 mg.
- If multiple classes and/or high doses of medications are used, consider monitoring of vital signs and/or end tidal CO₂ per provider discretion.
- In 24 hours, if a patient receives >30 mg of haloperidol OR >30 mg of olanzapine OR if doses of both add up to >30 mg, notify hospitalist and perform EKG when patient is stable enough to tolerate it.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 7/14/23.


Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact
Thomas_Peterson@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline Intoxicated Patient in the ED



Common Complications of Acute Alcohol Intoxication

- Hypoglycemia
- Electrolyte abnormality
- Hypothermia
- Occult trauma
- Co-ingestion/intoxication
- Gastritis
- Pancreatitis
- Hepatitis
- Occult infection
- Aspiration
- Exacerbation of chronic illness
- Victim of physical/sexual assault

Alcohol Metabolism

- $(\text{Serum Alcohol} - 80) / (20 \text{ to } 30) = \text{Time to sobriety}$
- $\text{BRAC} \times 1000 = \text{Serum Alcohol}$
- Serum alcohol <80 is considered sober.

Deescalation Strategies

- If not immediately dangerous, attempt simple, nonrestrictive strategies:
 - Verbal de-escalation.
 - Reduction of environmental stimuli (a quiet room is much better than a loud hallway).
 - Offer basic needs (ex, food, warm blanket).

Medications

Use caution when giving medications to intoxicated patients, as alcohol can intensify sedation effects.

- Oral vs Intramuscular - If the patient is cooperative, offer oral medications first
 - May give the patient sense of some control.
 - Avoid trauma of being physically restrained for IM shot.
 - Many medications are equally effective in oral form
 - If patient is not cooperative, the oral route is not going to be an option.
 - Benzodiazepines
 - Lorazepam – 0.05-0.1 mg/kg/dose (PO/IM/IV)
 - Midazolam – 0.25-0.5 mg/kg/dose PO; 0.2-0.3 mg/kg IN; 0.1-0.15 mg/kg/dose IM
 - First Generation Antipsychotics
 - Haloperidol – 2-10 mg PO or IM/IV, may repeat IM/IV every 15 min up to 30 mg
 - Second Generation Antipsychotics
 - Risperidone – 0.25-2 mg PO/ODT
 - Olanzapine – 2.5-5 mg PO/ODT
 - Others:
 - Diphenhydramine – 25-50mg PO or IM
 - Ketamine – 1-2 mg/kg/dose IV or 4-5 mg/kg/dose IM
- Rapid onset due to high bioavailability (even when given IM)
No QT prolongation issues
Safe even in overdose (important when you aren't sure of patient weight)
No respiratory depression (rarely, may see laryngospasm)

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 5/28/24.

Click [here](#) to see the supplemental resources for this guideline.

If comments about this guideline, please contact
clinical_guidelines@ykhc.org.



Yukon-Kuskokwim HEALTH CORPORATION

Clinical Guideline

Involuntary Psychiatric Admissions

****This guideline only applies to medically stable patients.****

Patient in village.

Patient in Bethel.

Escorts

- Must be sober and responsible and ensure patient arrives safely. May be TPO/VPO or Trooper.
- If patient is a minor, try to get legal guardian as escort to sign documents and assist with placement if needed.

- Health aide sends RMT for potential Involuntary Psychiatric Admission.
- Provider instructs health aide to contact BH on call if not done already.
- If patient has medical concerns, provider manages as appropriate (e.g. NAC for acetaminophen overdose, hemorrhage control for wounds).
- RMT provider can sign title paperwork or talk to BH about who should sign.
- Patient brought to Bethel with escort. (See box.) BH arranges travel.

Behavioral Health coordinates transfer to Bethel ED, MC-105 (T-47), and escort by Bethel Police.

Contact

- **BH Emergency Services:** x6499
- Tiger Connect BH Emergency (not under roles)
- **Psychiatry:** Tiger Connect role Psych Inpt On Site

Work-up

As there is no medical indication for routine lab work on psychiatric patients, provider may decline to obtain lab work and document reasoning. The following may be considered:

- CBC, CMP
- UDS: result can be followed by DES/inpatient unless newly psychotic patient with concern for acute drug intoxication
- EtOH (BAL or serum)
- APAP/salicylate levels if suspected ingestion or history
- UA if age >65

Definitions

- MC-100: Petition for Order Authorizing Hospitalization for Evaluation. Must be confirmed by a judge.
- MC-105: Notice of Emergency Detention and Application for Evaluation, often referred to as "Title 47." May be completed by law enforcement or physician.

Is patient intoxicated?

Yes

MC-105 (T-47) maintained until alcohol level <80.

No

ED provider evaluates patient.

Is the patient any of the following?

1. Threat to self
2. Threat to others
3. Gravely disabled

Yes

No

- MC-105 (T-47) is maintained.
- ED provider performs appropriate work-up (see box) and determines when patient is medically stable.
- BH or psychiatry provider evaluates patient and completes MC-100.
- If BH evaluation will be delayed and ED / NW provider are in agreement patient warrants admission on psychiatric hold, BH evaluation may occur after admission.

- MC-105 (T-47) is allowed to lapse.
- Behavioral Health evaluates patient and works with provider to determine disposition, which may include home with a safety plan, CRC, or voluntary admission to Inpatient Unit.

Admission

- Hospitalist admits patient to hospital using BH Inpatient Admission order set along with general admission orders.
- Hospitalist writes H&P, addressing both psychiatric and medical conditions.
- When patient is medically stable, hospitalist signs patient over to psychiatry service:
 - Hospitalist documents in a note that patient is cleared for transition to psychiatry service.
 - Hospitalist confirms plan with charge nurse and psychiatry service verbally or via Tiger Connect.
 - Hospitalist places communication order "The patient is transferred to psychiatry service."
 - Hospitalist does not need to round on patient any longer but may choose to remain involved as needed and may be re-consulted for concerns.

Medications to Treat a Combative Patient (Use ED T-47, Psychiatric Disorder Power Plan.)

- Olanzapine 5-10 mg IM/PO Q30 minutes up to max daily dose 60 mg.
- Haloperidol 2.5-10 mg IM/PO Q30 minutes, max daily dose 100 mg.
- Diphenhydramine 25-50 mg IV/IM/PO Q4-6h.
- Lorazepam 2-10 mg IV/IM/PO Q30 minutes, titrate to effect. No max dose. Avoid in intoxicated patients due to risk of respiratory depression.
- Ketamine 0.1-2 mg/kg IV Q10 minutes, max 2 mg/kg total dose.
- Ketamine 1-5 mg/kg IM Q30 minutes, max 5 mg/kg total dose. Consider for temporary control when other medications have failed or if immediate sedation is needed to prevent harm to patient or staff.

CAUTION: There is a risk of respiratory depression with all sedative medications, especially in the setting of alcohol use. Start with 1-2 agents and titrate. Do not add additional medications until prior medications are given time to work. All patients receiving sedative medications must be on continuous pulse-oximetry when they are no longer combative. 1:1 monitoring is required due to ligature risk. Consider ET CO₂ monitoring.

Services at YKHC

- **Behavioral Health (BH):** Masters level clinicians (MSW, LPC, etc.) who provide consultation services and are physically present in the hospital. They field calls from patients, assist Psychiatry in determining whether a patient needs involuntary hospitalization, and coordinate the logistics for where psychiatric patients go. They do not have legal authority to place psychiatric holds and do not have admitting privileges. **It is ultimately a physician's responsibility to determine suitability of psychiatric hold and appropriate disposition.** Non-physician providers may evaluate and treat these patients and maintain existing T-47s. If a new T-47 needs to be initiated, a physician must sign off on it.
- **Psychiatry:** All inpatient psychiatric care (including discharge or transfer to a higher level of care) is provided by a psychiatric physician or an advanced practice psychiatric provider under direct supervision by a psychiatric physician. Psychiatry will manage all patients on the psychiatry service, will be responsible for all patients on Title 47 commitments with the aid of BHES, and will also provide consultation for psychiatric patients on the Inpatient Unit.
- **Inpatient Hospitalists:** Family medicine physicians who admit patients, stabilize medical problems, and transfer to psychiatric service when medically stable. Hospitalist determines whether a patient has medical concerns requiring active ongoing inpatient management (e.g. infection, electrolyte abnormality, alcohol withdrawal). If medical problems, hospitalists remain primary service of record until active medical problems are resolved, writing daily progress notes, placing orders, and billing as usual with psychiatry consulting. If no medical problems, hospitalist may immediately sign patient over to psychiatry service. They can defer all psychiatric management to psychiatry service or collaborate with psychiatry team in rendering diagnoses and ordering medications. This should be communicated clearly both in the note and via direct conversation with psychiatry service.

This guideline is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 5/15/23. Click [here](#) to see the supplemental resources for this guideline. If comments about this guideline, please contact Travis_Nelson@ykhc.org or Kaia_Pearson@ykhc.org.



Non-Psychiatric Medical Needs

At time of transfer to psychiatric service, family medicine hospitalist will write non-psychiatric prescriptions (example: antibiotics, iron, etc.) via Med Rec and non-psychiatric follow-up plans (example: Depo in three months) under Patient Education and Follow-up.

- Family medicine hospitalist has admitted psychiatric patient to the hospital and addressed non-psychiatric medical needs (see box).
- Patient has been medically cleared.
- Psychiatric service assumes care.

Is admission voluntary or involuntary?

Voluntary

Involuntary

Care proceeds per hospital protocols until ready for discharge.

If at any time, patient requests to leave, does psychiatry staff determine that criteria for **MC-105** are met?

No

- If attending psychiatric physician feels patient is still in need of inpatient care, patient should be discharged AMA.
- Discharge process (see box) should be followed.

Yes

Onsite Psych Provider completes MC-105.

Psychiatry onsite provider and attending psychiatric physician complete the **MC-100** and send to courts within 18 hours of request to leave.

Onsite Psych Provider confirms **MC-100** has been completed and sent to court..

Psychiatry on site provider tells the patient their **rights** verbally and provides a written description of the YK commitment process.

Psychiatry staff testify at court per standard process.

Judge orders release.

Attending psychiatric physician is notified of judge order.

Judge orders continued stay.

- Patient must be transferred to a facility with a higher level of care.
- BHES notifies the accepting unit and sends/emails the required documentation (H&P, lab results, etc.).

BHES tracks timing of transfer to the accepting facility and updates psychiatry unit treatment team each morning in daily report.

Once a transfer date is determined, BHES completes the WEKA travel paperwork and arranges for travel.

- Psych RN & BHES complete the transfer process.
- Discharge orders are entered by psychiatric provider.

Definitions

- BHES: Behavioral Health Emergency Services
- MC-100: Petition for Order Authorizing Hospitalization for Evaluation. Must be confirmed by a judge.
- MC-105: Notice of Emergency Detention and Application for Evaluation, often referred to as "Title 47." May be completed by law enforcement or physician. Criteria: Is the patient any of the following?
 1. Threat to self
 2. Threat to others
 3. Gravely disabled
 See [Involuntary Psychiatric Admissions](#) guideline for more details.

If the patient stabilizes at any time per the attending psychiatric physician's assessment (with input from the entire treatment team), psychiatry completes MC-412 and submits to the court. The patient may be discharged after the discharge process and discharge orders are completed.

Discharge Process

- Psych RN & BHES complete the discharge process.
- Psychiatric care coordinator arranges outpatient appointment for day following discharge as well as the day 3 BHES safety check assessment.
- BHES complete discharge safety plan with the patient prior to discharge.
- Discharge orders are entered by psychiatric provider.
- The YK Psychiatry inpatient post discharge intensive outpatient care plan is initiated.

Discharge process (see box) is followed. Psychiatric onsite provider specifies in orders, "as ordered by judge."



Communication	
<input type="checkbox"/> Has YKHC pediatric group been briefed and asked for feedback on concerns or issues?	<input type="checkbox"/> N/A
<input type="checkbox"/> Prior to patient returning, has care conference been scheduled with 1-2 pediatricians to represent group/consensus recommendations? Other key participants include: case managers, SRC providers, health aides, and family members.	<input type="checkbox"/> N/A
<input type="checkbox"/> Has an Informed Consent to Return to Village been customized for this patient and approved by Risk Management (Chris Beltzer as of 4/2025)? [See Peds Folder → Informed Consent to Return to Village for template.]	<input type="checkbox"/> N/A
<input type="checkbox"/> Have the caregivers completed the Informed Consent to Return to Village?	<input type="checkbox"/> N/A
<input type="checkbox"/> If patient is returning to the village against medical advice, have Risk Management, Clinical Director, and appropriate administrators been made aware?	<input type="checkbox"/> N/A
<input type="checkbox"/> Does the patient have a modified code status? If so, have the Expected Home Death Forms been completed? Has the POLST Form been completed?	<input type="checkbox"/> N/A
<input type="checkbox"/> After the care conference: has a detailed note been placed in the chart summarizing the care conference? Has this note been sent by email to the pediatric group, case managers, and SRC providers?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has the YKHC RAVEN Problem List been updated with care plans, follow-up needs, therapeutic parameters, etc.?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has the YKHC chart been updated with a current phone number and email address?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has the family been given the YKHC Follow Up resource ?	<input type="checkbox"/> N/A

Medical Home	
<input type="checkbox"/> Where will primary care occur – village, SRC, Bethel, or Anchorage?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has a clinic appointment been scheduled to establish care at YKHC?	<input type="checkbox"/> N/A
<input type="checkbox"/> Have VTC appointments been set up for patient and family?	<input type="checkbox"/> N/A
<input type="checkbox"/> Have the health aides been notified of the complex needs of this patient?	<input type="checkbox"/> N/A
<input type="checkbox"/> Have the nearest SRC providers been notified of the complex needs of this patient?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has caregiver referral to YKHC Behavioral Health been offered?	<input type="checkbox"/> N/A

Medical Supplies and Equipment for Home	
<input type="checkbox"/> Have all needed medical supplies and emergency equipment been identified? Ex: medications, beds, commodes, syringes, dressings, wheelchair, ambu bag (if no CHA available), suction, pulse-oximeter, oxygen, glucometer, etc.	<input type="checkbox"/> N/A
<input type="checkbox"/> Do the caregivers have needed supplies/equipment?	<input type="checkbox"/> N/A
<input type="checkbox"/> Have caregivers received training on how to use this equipment?	<input type="checkbox"/> N/A
<input type="checkbox"/> Does home have electricity, running water, and a refrigerator?	<input type="checkbox"/> N/A
<input type="checkbox"/> Is there a back-up plan in place if electricity goes down?	<input type="checkbox"/> N/A
<input type="checkbox"/> Have family/caregivers received CPR training?	<input type="checkbox"/> N/A

Medications	
<input type="checkbox"/> Have all current and anticipated prescriptions with refills been ordered on the YKHC RAVEN Medication List?	<input type="checkbox"/> N/A
<input type="checkbox"/> If the patient is at risk for seizures, has the family received Diastat or intranasal midazolam and received the appropriate training?	<input type="checkbox"/> N/A
<input type="checkbox"/> Is there a prescription for electrolyte replacement solution (ex: Pedialyte)?	<input type="checkbox"/> N/A



Gastrostomy Tube

<input type="checkbox"/> Have two caregivers been trained on G-tube care and replacement?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has a CMN been submitted for formula, supplies, and replacement G-tubes?	<input type="checkbox"/> N/A
<input type="checkbox"/> Does family have emergency and replacement supplies, including an extra G-tube and Foley catheters in the same French size and smaller sizes?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has family been counseled that health aides cannot help with G-tube problems, and that they may have to come to Bethel or even Anchorage for G-tube care in the future?	<input type="checkbox"/> N/A

Home Oxygen

<input type="checkbox"/> Has all the necessary paperwork been filled out and are YK Case Managers aware?	<input type="checkbox"/> N/A
<input type="checkbox"/> Does family have all needed supplies – O2, tubing, pulse ox, suction, etc.?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has an Informed Consent to Return to Village been customized and completed?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has the YKHC problem list been updated with “ALERT – emergency care plan”?	<input type="checkbox"/> N/A

Ventriculoperitoneal Shunt

<input type="checkbox"/> Have the nearest SRC providers been informed of return to village?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has an Informed Consent to Return to Village been customized and completed?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has the YKHC problem list been updated with “ALERT – emergency care plan”?	<input type="checkbox"/> N/A

Port-a-cath

<input type="checkbox"/> Have at least two caregivers been trained on and are comfortable with accessing port?	<input type="checkbox"/> N/A
<input type="checkbox"/> Do caregivers have all needed supplies?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has an Informed Consent to Return to Village been customized and completed?	<input type="checkbox"/> N/A
<input type="checkbox"/> Has the YKHC problem list been updated with “ALERT – emergency care plan”?	<input type="checkbox"/> N/A

Baclofen

<input type="checkbox"/> Family has been counselled on the risks and symptoms of withdrawal.	<input type="checkbox"/> N/A
<input type="checkbox"/> Baclofen has been prescribed using the 25 mg/5 mL concentration.	<input type="checkbox"/> N/A
<input type="checkbox"/> Family has baclofen tablets for use if liquid supply runs out. Family knows how to crush and give tablets.	<input type="checkbox"/> N/A



Yukon-Kuskokwim HEALTH CORPORATION

Returning to the Yukon Kuskokwim Delta Region

Name: _____

Birthday: _____

Next Appointment

Date: _____

Location: Village clinic Bethel clinic

Important Phone Numbers

YKHC Travel: (907) 543-6846
 (907) 543-6625
 (833) 543-6625 (for text)

YKHC Pediatric Case Managers: (907) 543-6634

Call before returning home to the YK Delta and anytime for questions about appointments, equipment, paperwork, etc.

Your Village Clinic: _____

After hours, this number will be answered by the YKHC Nurse Triage Line. These nurses are trained to give advice by the phone. If there is an emergency, they can contact the health aide on-call for you. Please be patient; sometimes you will need to be on hold while the nurse helps other patients. Please hold until your call is answered.



Documentation Requirements for Pediatric Nutritional Supplements

The following resource is from the Medicaid Certificate of Medical Necessity.

Medicaid, Medicare, and other insurers have specific requirements for medical provider documentation. If those requirements are not met, nutritional supplements will not be covered.

Use the autotext “..nutritionalsupplementdocumentation.”

Documentation Requirements for the Prescription of Nutritional Supplements:

The following objective documentation is required to show the medical necessity of the nutritional supplement being prescribed.

This information needs to appear in the body of the medical provider's chart notes:

- Diagnosis of the patient including ICD-10 code.
- Product being prescribed. (Example: PediaSure)
- Why product is medically necessary.
- Goal or target weight for the patient.
- Total daily caloric requirement.
- Total daily calories obtained from ingestion (oral) foods.
- Total daily calories to be obtained from nutritional supplement.

Documentation Example

PediaSure is medically necessary for this child.

Diagnosis: dysphagia (R13.10), G-tube dependence

Product: PediaSure

Medical Necessity: Patient has severe dysphagia. He is undergoing oral feeding therapy but is unable to take any degree of sufficient calories by mouth and is thus entirely dependent on a G-tube for nutrition. PediaSure will give him the nutrition he needs to survive.

Goal/Target Weight: currently at target weight of XX kg (XXth percentile for age when corrected for prematurity). Target weight along this trajectory in one year will be XX kg.

Total Daily Caloric Requirement: XX calories/day (usually estimate 100-120 cal/kg/day – adjust based on growth)

Total Calories Obtained from Oral Intake: 0 calories/day

Total Daily Calories to be Obtained from Nutritional Supplement: XX calories/day

For resources and information about nutritional supplements in former premature babies, please see the [ANMC Guideline on Preterm Infant Nutrition through 2 Years Old](#).

This resource is designed for the general use of most patients but may need to be adapted to meet the special needs of a specific patient as determined by the medical practitioner.

Approved by Clinical Guideline Committee 11/27/22.

If comments about this resource, please contact Tamara_Hill@ykhc.org.